

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 April 2003 (24.04.2003)

PCT

(10) International Publication Number
WO 03/033724 A2

(51) International Patent Classification⁷: **C12Q**

(21) International Application Number: PCT/US02/33244

(22) International Filing Date: 15 October 2002 (15.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/977,868 15 October 2001 (15.10.2001) US
09/982,212 18 October 2001 (18.10.2001) US

(71) Applicant: **MOLECULAR STAGING, INC.** [US/US];
300 George Street, Suite 701, New Haven, CT 06511 (US).

(72) Inventors: **DEAN, Frank, B.**; 21 Pearl Street, Guilford, CT 06437 (US). **LASKEN, Roger, S.**; 364 Central Avenue, First Floor, New Haven, CT 06515 (US). **FANG, Linhua**; 6 Svea Avenue, Second Floor, Branford, CT 06405 (US). **FARUQI, Fawad, A.**; 985 Durham Road, Guilford, CT 06437 (US). **ALSMADI, Osama, A.**; 310 Glenwood

Drive, Guilford, CT 06437 (US). **DRISCOLL, Mark, D.**; 14 Bruce Lane, Wallingford, CT 06492 (US). **HOSONO, Seiyu**; 140 New Street, Apt. 2315, Mamaroneck, NY 10543 (US). **WISNIEWSKI, Michele**; 913 Aspen Glen Drive, Hamden, CT 06518 (US). **SONG, Wanmin**; 55F Briarwood Lane, Branford, CT 06405 (US).


(74) Agents: **HODGES, Robert, A.** et al.; Needle & Rosenberg, P.C., 127 Peachtree Street, NE, The Candler Building, Atlanta, GA 30308-1811 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: NUCLEIC ACID AMPLIFICATION



(57) Abstract: Disclosed are compositions and a method for amplification of nucleic acid sequences of interest. The disclosed method generally involves replication of a target sequence such that, during replication, the replicated strands are displaced from the target sequence by strand displacement replication of another replicated strand. In one form of the disclosed method, the target sample is not subjected to denaturing conditions. It was discovered that the target nucleic acids, genomic DNA, for example, need not be denatured for efficient multiple displacement amplification. The primers used can be hexamer primers. The primers can also each contain at least one modified nucleotide such that the primers are nuclease resistant. The primers can also each contain at least one modified nucleotide such that the melting temperature of the primer is altered relative to a primer of the same sequence without the modified nucleotide(s). The DNA polymerase can be ϕ 29 DNA polymerase.



WO 03/033724 A2



European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished upon receipt of that report*

NUCLEIC ACID AMPLIFICATION

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of copending Application No. 09/982,212, filed October 18, 2001, which is a continuation of copending Application No. 09/977,868, filed October 15, 2001, both of which are hereby incorporated herein
5 by reference in their entirety.

FIELD OF THE INVENTION

The disclosed invention is generally in the field of nucleic acid amplification.

BACKGROUND OF THE INVENTION

10 A number of methods have been developed for exponential amplification of nucleic acids. These include the polymerase chain reaction (PCR), ligase chain reaction (LCR), self-sustained sequence replication (3SR), nucleic acid sequence based amplification (NASBA), strand displacement amplification (SDA), and amplification with Q β replicase (Birkenmeyer and Mushahwar, *J. Virological Methods*, **35**:117-126
15 (1991); Landegren, *Trends Genetics* **9**:199-202 (1993)).

Fundamental to most genetic analysis is availability of genomic DNA of adequate quality and quantity. Since DNA yield from human samples is frequently limiting, much effort has been invested in general methods for propagating and archiving genomic DNA. Methods include the creation of EBV-transformed cell lines
20 or whole genome amplification (WGA) by random or degenerate oligonucleotide-primed PCR. Whole genome PCR, a variant of PCR amplification, involves the use of random or partially random primers to amplify the entire genome of an organism in the same PCR reaction. This technique relies on having a sufficient number of primers of random or partially random sequence such that pairs of primers will hybridize
25 throughout the genomic DNA at moderate intervals. Replication initiated at the primers can then result in replicated strands overlapping sites where another primer can hybridize. By subjecting the genomic sample to multiple amplification cycles, the genomic sequences will be amplified. Whole genome PCR has the same disadvantages as other forms of PCR. However, WGA methods suffer from high cost or insufficient
30 coverage and inadequate average DNA size (Telenius *et al.*, *Genomics*. **13**:718-725 (1992); Cheung and Nelson, *Proc Natl Acad Sci U S A.* **93**:14676-14679 (1996); Zhang *et al.*, *Proc Natl Acad Sci U S A.* **89**:5847-5851 (1992)).

Another form of nucleic acid amplification, involving strand displacement, has been described in U.S. Patent No. 6,124,120 to Lizardi. In one form of the method, two sets of primers are used that are complementary to opposite strands of nucleotide sequences flanking a target sequence. Amplification proceeds by replication initiated at each primer and continuing through the target nucleic acid sequence, with the growing strands encountering and displacing previously replicated strands. In another form of the method a random set of primers is used to randomly prime a sample of genomic nucleic acid. The primers in the set are collectively, and randomly, complementary to nucleic acid sequences distributed throughout nucleic acid in the sample.

Amplification proceeds by replication initiating at each primer and continuing so that the growing strands encounter and displace adjacent replicated strands. In another form of the method concatenated DNA is amplified by strand displacement synthesis with either a random set of primers or primers complementary to linker sequences between the concatenated DNA. Synthesis proceeds from the linkers, through a section of the concatenated DNA to the next linker, and continues beyond, with the growing strands encountering and displacing previously replicated strands.

BRIEF SUMMARY OF THE INVENTION

Disclosed are compositions and a method for amplification of nucleic acid sequences of interest. The method is based on strand displacement replication of the nucleic acid sequences by multiple primers. The disclosed method, referred to as multiple displacement amplification (MDA), improves on prior methods of strand displacement replication. The disclosed method generally involves bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence.

Replication of the target sequence results in replicated strands such that, during replication, the replicated strands are displaced from the target sequence by strand displacement replication of another replicated strand.

In some forms of the disclosed method, a genomic sample is prepared by exposing cells to alkaline conditions, thereby lysing the cells and resulting in a cell lysate; reducing the pH of the cell lysate to make the pH of the cell lysate compatible with DNA replication; and incubating the cell lysate under conditions that promote replication of the genome of the cells by multiple displacement amplification. It has

been discovered that alkaline lysis can cause less damage to genomic DNA and that alkaline lysis is compatible with multiple displacement amplification. The alkaline conditions can be, for example, those that cause a substantial number of cells to lyse or those that cause a sufficient number of cells to lyse. The number of lysed cells can be
5 considered sufficient if the genome can be sufficiently amplified in the disclosed method. The amplification is sufficient if enough amplification product is produced to permit some use of the amplification product, such as detection of sequences or other analysis. The reduction in pH is generally into the neutral range of pH 9.0 to pH 6.0.

In some embodiments, the cells are not lysed by heat and/or the nucleic acids in
10 the cell lysate is not denatured by heating. Those of skill in the art will understand that different cells under different conditions will be lysed at different temperatures and so can determine temperatures and times at which the cells will not be lysed by heat. In general, the cells are not subjected to heating above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions used. In
15 some embodiments, the cells and/or cell lysate are not subjected to heating substantially above the temperature at which the cells grow. In other embodiments, the cells and/or cell lysate are not subjected to heating substantially above the temperature of the amplification reaction (where the genome is replicated). The disclosed multiple displacement amplification reaction is generally conducted at a substantially constant
20 temperature (that is, the amplification reaction is substantially isothermic), and this temperature is generally below the temperature at which the nucleic acids would be substantially or significantly denatured.

In some embodiments, the cell lysate is not subjected to purification prior to the amplification reaction. In the context of the disclosed method, purification generally
25 refers to the separation of nucleic acids from other material in the cell lysate. It has been discovered that multiple displacement amplification can be performed on unpurified and partially purified samples. It is commonly thought that amplification reactions cannot be efficiently performed using unpurified nucleic acid. In particular, PCR is very sensitive to contaminants.

30 In some forms of the disclosed method, the target sample is not subjected to denaturing conditions. It was discovered that the target nucleic acids, genomic DNA, for example, need not be denatured for efficient multiple displacement amplification. It

was discovered that elimination of a denaturation step and denaturation conditions has additional advantages such as reducing sequence bias in the amplified products. In another embodiment, the primers can be hexamer primers. It was discovered that such short, 6 nucleotide primers can still prime multiple strand displacement replication efficiently. Such short primers are easier to produce as a complete set of primers of random sequence (random primers) than longer primers because there are fewer separate species of primers in a pool of shorter primers. In another embodiment, the primers can each contain at least one modified nucleotide such that the primers are nuclease resistant. In another embodiment, the primers can each contain at least one modified nucleotide such that the melting temperature of the primer is altered relative to a primer of the same sequence without the modified nucleotide(s). For these last two embodiments, it is preferred that the primers are modified RNA. In another embodiment, the DNA polymerase can be ϕ 29 DNA polymerase. It was discovered that ϕ 29 DNA polymerase produces greater amplification in multiple displacement amplification. The combination of two or more of the above features also yields improved results in multiple displacement amplification. In a preferred embodiment, for example, the target sample is not subjected to denaturing conditions, the primers are hexamer primers and contain modified nucleotides such that the primers are nuclease resistant, and the DNA polymerase is ϕ 29 DNA polymerase. The above features are especially useful in whole genome strand displacement amplification (WGSDA).

In some forms of the disclosed method, the method includes labeling of the replicated strands (that is, the strands produced in multiple displacement amplification) using terminal deoxynucleotidyl transferase. The replicated strands can be labeled by, for example, the addition of modified nucleotides, such as biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, bromodeoxyuridine triphosphate (BrdUTP), or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates, to the 3' ends of the replicated strands. The replicated strands can also be labeled by incorporating modified nucleotides during replication. Probes replicated in this manner are particularly useful for hybridization, including use in microarray formats.

In one form of the disclosed method, referred to as whole genome strand displacement amplification (WGSDA), a random set of primers is used to randomly prime a sample of genomic nucleic acid (or another sample of nucleic acid of high

complexity). By choosing a sufficiently large set of primers of random or partially random sequence, the primers in the set will be collectively, and randomly, complementary to nucleic acid sequences distributed throughout nucleic acid in the sample. Amplification proceeds by replication with a highly processive polymerase initiating at each primer and continuing until spontaneous termination. A key feature of this method is the displacement of intervening primers during replication by the polymerase. In this way, multiple overlapping copies of the entire genome can be synthesized in a short time. The method has advantages over the polymerase chain reaction since it can be carried out under isothermal conditions. Other advantages of whole genome strand displacement amplification include a higher level of amplification than whole genome PCR (up to five times higher), amplification is less sequence-dependent than PCR, and there are no re-annealing artifacts or gene shuffling artifacts as can occur with PCR (since there are no cycles of denaturation and re-annealing). In preferred embodiments of WGSDA, the target sample is not subjected to denaturing conditions, the primers are hexamer primers and contain modified nucleotides such that the primers are nuclease resistant, the DNA polymerase is ϕ 29 DNA polymerase, or any combination of these features.

In another form of the method, referred to as multiple strand displacement amplification (MSDA), two sets of primers are used, a right set and a left set. Primers in the right set of primers each have a portion complementary to nucleotide sequences flanking one side of a target nucleotide sequence and primers in the left set of primers each have a portion complementary to nucleotide sequences flanking the other side of the target nucleotide sequence. The primers in the right set are complementary to one strand of the nucleic acid molecule containing the target nucleotide sequence and the primers in the left set are complementary to the opposite strand. The 5' end of primers in both sets are distal to the nucleic acid sequence of interest when the primers are hybridized to the flanking sequences in the nucleic acid molecule. Preferably, each member of each set has a portion complementary to a separate and non-overlapping nucleotide sequence flanking the target nucleotide sequence. Amplification proceeds by replication initiated at each primer and continuing through the target nucleic acid sequence. In another form of MSDA, referred to as linear MSDA, amplification is

performed with a set of primers complementary to only one strand, thus amplifying only one of the strands.

In another form of the method, referred to as gene specific strand displacement amplification (GS-MSDA), target DNA is first digested with a restriction endonuclease. The digested fragments are then ligated end-to-end to form DNA circles. These circles can be monomers or concatemers. Two sets of primers are used for amplification, a right set and a left set. Primers in the right set of primers each have a portion complementary to nucleotide sequences flanking one side of a target nucleotide sequence and primers in the left set of primers each have a portion complementary to nucleotide sequences flanking the other side of the target nucleotide sequence. The primers in the right set are complementary to one strand of the nucleic acid molecule containing the target nucleotide sequence and the primers in the left set are complementary to the opposite strand. The primers are designed to cover all or part of the sequence needed to be amplified. Preferably, each member of each set has a portion complementary to a separate and non-overlapping nucleotide sequence flanking the target nucleotide sequence. Amplification proceeds by replication initiated at each primer and continuing through the target nucleic acid sequence. In one form of GS-MSDA, referred to as linear GS-MSDA, amplification is performed with a set of primers complementary to only one strand, thus amplifying only one of the strands. In another form of GS-MSDA, cDNA sequences can be circularized to form single stranded DNA circles. Amplification is then performed with a set of primers complementary to the single-stranded circular cDNA.

A key feature of this method is the displacement of intervening primers during replication. Once the nucleic acid strands elongated from the right set of primers reaches the region of the nucleic acid molecule to which the left set of primers hybridizes, and vice versa, another round of priming and replication will take place. This allows multiple copies of a nested set of the target nucleic acid sequence to be synthesized in a short period of time. By using a sufficient number of primers in the right and left sets, only a few rounds of replication are required to produce hundreds of thousands of copies of the nucleic acid sequence of interest. The disclosed method has advantages over the polymerase chain reaction since it can be carried out under isothermal conditions. No thermal cycling is needed because the polymerase at the

head of an elongating strand (or a compatible strand-displacement protein) will displace, and thereby make available for hybridization, the strand ahead of it. Other advantages of multiple strand displacement amplification include the ability to amplify very long nucleic acid segments (on the order of 50 kilobases) and rapid amplification of shorter segments (10 kilobases or less). In multiple strand displacement amplification, single priming events at unintended sites will not lead to artifactual amplification at these sites (since amplification at the intended site will quickly outstrip the single strand replication at the unintended site). In preferred embodiments of MSDA, the target sample is not subjected to denaturing conditions, the primers are hexamer primers and contain modified nucleotides such that the primers are nuclease resistant, the DNA polymerase is ϕ 29 DNA polymerase, or any combination of these features.

In preferred embodiments of WGSDA, the target sample is not subjected to denaturing conditions, the primers are hexamer primers and contain modified nucleotides such that the primers are nuclease resistant, the DNA polymerase is ϕ 29 DNA polymerase, or any combination of these features.

Following amplification, the amplified sequences can be used for any purpose, such as uses known and established for PCR amplified sequences. For example, amplified sequences can be detected using any of the conventional detection systems for nucleic acids such as detection of fluorescent labels, enzyme-linked detection systems, antibody-mediated label detection, and detection of radioactive labels. A preferred form of labeling involves labeling of the replicated strands (that is, the strands produced in multiple displacement amplification) using terminal deoxynucleotidyl transferase. The replicated strands can be labeled by, for example, the addition of modified nucleotides, such as biotinylated nucleotides, fluorescent nucleotides, 5-methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates, to the 3' ends of the replicated strands.

In the disclosed method amplification takes place not in cycles, but in a continuous, isothermal replication. This makes amplification less complicated and much more consistent in output. Strand displacement allows rapid generation of multiple copies of a nucleic acid sequence or sample in a single, continuous, isothermal reaction. DNA that has been produced using the disclosed method can then be used for

any purpose or in any other method desired. For example, PCR can be used to further amplify any specific DNA sequence that has been previously amplified by the whole genome strand displacement method.

Genetic analysis must frequently be carried out with DNA derived from
5 biological samples, such as blood, tissue culture cells, buccal swabs, mouthwash, stool, tissues slices, biopsy aspiration, and archeological samples such as bone or mummified tissue. In some cases, the samples are too small to extract a sufficient amount of pure DNA and it is necessary to carry out DNA-based assays directly from the unprocessed sample. Furthermore, it is time consuming to isolate pure DNA, and so the disclosed
10 method, which can amplify the genome directly from biological samples, represents a substantial improvement.

The disclosed method has several distinct advantages over current methodologies. The genome can be amplified directly from whole blood or cultured cells with simple cell lysis techniques such as KOH treatment. PCR and other DNA
15 amplification methods are severely inhibited by cellular contents and so purification of DNA is needed prior to amplification and assay. For example, heme present in lysed blood cells inhibits PCR. In contrast, the disclosed form of whole genome amplification can be carried out on crude lysates with no need to physically separate DNA by miniprep extraction and precipitation procedures, or with column or spin
20 cartridge methods.

Bacteria, fungi, and viruses may all be involved in nosocomial infections. Identification of nosocomial pathogens at the sub-species level requires sophisticated discriminatory techniques. Such techniques utilize traditional as well as molecular methods for typing. Some traditional techniques are antimicrobial susceptibility
25 testing, determination of the ability to utilize biochemical substrates, and serotyping. A major limitation of these techniques is that they take several days to complete, since they require pure bacterial cultures. Because such techniques are long, and the bacteria may even be non-viable in the clinical samples, there is a need to have a quick and reliable method for bacterial species identification.

30 Some of the DNA-based molecular methods for the identification of bacterial species are macrorestriction analysis (MRA) followed by pulsed-field gel electrophoresis (PFGE), amplified fragment length polymorphism (AFLP) analysis, and

arbitrarily primed PCR (AP-PCR) (Tenover et al., J. Clin. Microbiol. 32:407-415 (1994), and Pruckler et al., J. Clin. Microbiol. 33:2872-2875 (1995)). These molecular techniques are labor-intensive and difficult to standardize among different laboratories.

The disclosed method provides a useful alternative method for the identification
5 of bacterial strains by amplification of microbial DNA for analysis. Unlike PCR (Lantz et al., Biotechnol. Annu. Rev. 5:87-130 (2000)), the disclosed method is rapid, non-biased, reproducible, and capable of amplifying large DNA segments from bacterial, viral or fungal genomes.

The disclosed method can be used, for example, to obtain enough DNA from
10 unculturable organisms for sequencing or other studies. Most microorganisms cannot be propagated outside their native environment, and therefore their nucleic acids cannot be sequenced. Many unculturable organisms live under extreme conditions, which makes their genetic complement of interest to investigators. Other microorganisms live in communities that play a vital role in certain ecosystems. Individual organisms or
15 entire communities of organisms can be amplified and sequenced, individually or together.

Recombinant proteins may be purified from a large biomass grown up from bacterial or yeast strains harboring desired expression vectors. A high degree of purity may be desired for the isolated recombinant protein, requiring a sensitive procedure for
20 the detection of trace levels of protein or DNA contaminants. The disclosed method is a DNA amplification reaction that is highly robust even in the presence of low levels of DNA template, and can be used to monitor preparations of recombinant protein for trace amounts of contaminating bacterial or yeast genomic DNA.

Amplification of forensic material for RFLP-based testing is one useful
25 application for the disclosed method.

Also disclosed is a method for amplifying and repairing damaged DNA. This method is useful, for example, for amplifying degraded genomic DNA. The method involves substantially denaturing a damaged DNA sample (generally via exposure to heat and alkaline conditions), removal or reduction of the denaturing conditions (such
30 as by reduction of the pH and temperature of the denatured DNA sample), and replicating the DNA. The damaged DNA is repaired during replication by increasing the average length of the damaged DNA. For example, the average length of DNA

fragments can be increase from, for example, 2 kb in the damaged DNA sample to, for example, 10 kb or greater for the replicated DNA. This repair method can result in an overall improvement in amplification of damaged DNA by increasing the average length of the product, increasing the quality of the amplification products by 3-fold (by, for example, increasing the marker representation in the sample), and improving the genotyping of amplified products by lowering the frequency of allelic dropout; all compared to the results when amplifying damaged DNA by other methods. The removal of denaturing conditions can allow denatured strands of damaged DNA to hybridize to other denatured damaged DNA. The replication can be multiple displacement amplification. Substantial denaturation and transient denaturation of the DNA samples generally is carried out such that the DNA is not further damaged. This method can generally be combined or used with any of the disclosed amplification methods.

It is an object of the disclosed invention to provide a method of amplifying a target nucleic acid sequence in a continuous, isothermal reaction.

It is another object of the disclosed invention to provide a method of amplifying an entire genome or other highly complex nucleic acid sample in a continuous, isothermal reaction.

It is another object of the disclosed invention to provide a method of amplifying a target nucleic acid sequence where multiple copies of the target nucleic acid sequence are produced in a single amplification cycle.

It is another object of the disclosed invention to provide a method of amplifying a concatenated DNA in a continuous, isothermal reaction.

It is another object of the disclosed invention to provide a kit for amplifying a target nucleic acid sequence in a continuous, isothermal reaction.

It is another object of the disclosed invention to provide a kit for amplifying an entire genome or other highly complex nucleic acid sample in a continuous, isothermal reaction.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph of DNA synthesis (in μg) versus time (in hours) using different amounts of nucleic acid for amplification in the disclosed method.

Figure 2 is a graph of the effect of incubation time at 95°C on template DNA length.

Figure 3 is a graph of the effect of template incubation at 95°C on the rate and yield of MDA.

5 Figure 4 is a graph of the effect of template incubation at 95°C on the average size of DNA product strands.

Figure 5 is a graph showing a comparison of the effect of template incubation at 95°C versus no incubation at 95°C on locus representation in DNA amplified by MDA.

10 Figures 6A, 6B, and 6C are graphs showing the effect of amplification on gene representation bias for three different amplification procedures, MDA, DOP-PCR, and PEP.

Figure 7 is a graph showing amplification of c-jun sequences using nested primers.

15 Figure 8 is a graph the relative representation of eight loci for DNA from five different amplification reactions. The Y-axis is the locus representation, expressed as a percent, relative to input genomic DNA, which is calculated as the yield of quantitative PCR product from 1 µg of amplified DNA divided by the yield from 1 µg of genomic DNA control.

20 Figure 9 is a graph showing a comparison of the percent representation for 8 loci for DNA amplified in a reaction containing 100% dTTP and DNA amplified in a reaction containing 30% dTTP / 70% AAdUTP.

25 Figure 10 is a graph showing the amplification of c-jun sequences using circularized genomic template. The Y-axis is the locus representation, expressed as a percent, relative to input genomic DNA, which is calculated as the yield of quantitative PCR product from 1 µg of amplified DNA divided by the yield from 1 µg of genomic DNA control.

Figure 11 is a graph showing a comparison of the percent representation for 8 loci in DNA amplified using c-jun specific primers and circularized DNA target.

30 Figure 12 is a graph of percent locus representation of different DNA samples exposed to different treatments (control or repair treatments).

Figure 13 is a graph of percent locus representation of 40 samples with or without repair treatment.

DETAILED DESCRIPTION OF THE INVENTION

The disclosed method makes use of certain materials and procedures which allow amplification of target nucleic acid sequences and whole genomes or other highly complex nucleic acid samples. These materials and procedures are described in detail
5 below.

Materials

A. Target Sequence

The target sequence, which is the object of amplification, can be any nucleic acid. The target sequence can include multiple nucleic acid molecules, such as in the
10 case of whole genome amplification, multiple sites in a nucleic acid molecule, or a single region of a nucleic acid molecule. For multiple strand displacement amplification, generally the target sequence is a single region in a nucleic acid molecule or nucleic acid sample. For whole genome amplification, the target sequence is the entire genome or nucleic acid sample. A target sequence can be in any nucleic acid
15 sample of interest. The source, identity, and preparation of many such nucleic acid samples are known. It is preferred that nucleic acid samples known or identified for use in amplification or detection methods be used for the method described herein. The nucleic acid sample can be, for example, a nucleic acid sample from one or more cells, tissue, or bodily fluids such as blood, urine, semen, lymphatic fluid, cerebrospinal fluid,
20 or amniotic fluid, or other biological samples, such as tissue culture cells, buccal swabs, mouthwash, stool, tissues slices, biopsy aspiration, and archeological samples such as bone or mummified tissue. Types of useful target samples include blood samples, urine samples, semen samples, lymphatic fluid samples, cerebrospinal fluid samples, amniotic fluid samples, biopsy samples, needle aspiration biopsy samples, cancer
25 samples, tumor samples, tissue samples, cell samples, cell lysate samples, a crude cell lysate samples, forensic samples, archeological samples, infection samples, nosocomial infection samples, production samples, drug preparation samples, biological molecule production samples, protein preparation samples, lipid preparation samples, and/or carbohydrate preparation samples.

30 For multiple strand displacement amplification, preferred target sequences are those which are difficult to amplify using PCR due to, for example, length or composition. For whole genome amplification, preferred target sequences are nucleic

acid samples from a single cell. For multiple strand displacement amplification of concatenated DNA the target is the concatenated DNA. The target sequence can be either one or both strands of cDNA. The target sequences for use in the disclosed method are preferably part of nucleic acid molecules or samples that are complex and non-repetitive (with the exception of the linkers in linker-concatenated DNA and sections of repetitive DNA in genomic DNA).

Target nucleic acids can include damaged DNA and damaged DNA samples. For example, preparation of genomic DNA samples can result in damage to the genomic DNA (for example, degradation and fragmentation). This can make amplification of the genome or sequences in it both more difficult and provide less reliable results (by, for example, resulting in amplification of many partial and fragmented genomic sequences. Damaged DNA and damaged DNA samples are thus useful for the disclosed method of amplifying damaged DNA. Any degraded, fragmented or otherwise damaged DNA or sample containing such DNA can be used in the disclosed method.

1. Target Sequences for Multiple Strand Displacement Amplification

Although multiple sites in a nucleic acid sample can be amplified simultaneously in the same MSDA reaction, for simplicity, the following discussion will refer to the features of a single nucleic acid sequence of interest which is to be amplified. This sequence is referred to below as a target sequence. It is preferred that a target sequence for MSDA include two types of target regions, an amplification target and a hybridization target. The hybridization target includes the sequences in the target sequence that are complementary to the primers in a set of primers. The amplification target is the portion of the target sequence which is to be amplified. For this purpose, the amplification target is preferably downstream of, or flanked by the hybridization target(s). There are no specific sequence or structural requirements for choosing a target sequence. The hybridization target and the amplification target within the target sequence are defined in terms of the relationship of the target sequence to the primers in a set of primers. The primers are designed to match the chosen target sequence. Although preferred, it is not required that sequences to be amplified and the sites of hybridization of the primers be separate since sequences in and around the sites where the primers hybridize will be amplified.

In multiple strand displacement amplification of circularized DNA, the circular DNA fragments are the amplification targets. The hybridization targets include the sequences that are complementary to the primers used for amplification. One form of circular DNA for amplification is circularized cDNA.

5 In multiple strand displacement amplification of linker-concatenated DNA, the DNA fragments joined by the linkers are the amplification targets and the linkers are the hybridization target. The hybridization targets (that is, the linkers) include the sequences that are complementary to the primers used for amplification. One form of concatenated DNA for amplification is concatenated cDNA.

10 B. Primers

Primers for use in the disclosed amplification method are oligonucleotides having sequence complementary to the target sequence. This sequence is referred to as the complementary portion of the primer. The complementary portion of a primer can be any length that supports specific and stable hybridization between the primer and the
15 target sequence under the reaction conditions. Generally, for reactions at 37°C, this can be 10 to 35 nucleotides long or 16 to 24 nucleotides long. For whole genome amplification, the primers can be from 5 to 60 nucleotides long, and in particular, can be 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and/or 20 nucleotides long.

For some forms of the disclosed method, such as those using primers or random
20 or degenerate sequence (that is, use of a collection of primers having a variety of sequences), primer hybridization need not be specific. In such cases the primers need only be effective in priming synthesis. For example, in whole genome amplification specificity of priming is not essential since the goal generally is to amplify all sequences equally. Sets of random or degenerate primers can be composed of primers
25 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and/or 20 nucleotides long or more. Primers six nucleotides long are referred to as hexamer primers. Preferred primers for whole genome amplification are random hexamer primers, for example, random hexamer primers where every possible six nucleotide sequence is represented in the set of primers. Similarly, sets of random primers of other particular lengths, or of a
30 mixture of lengths preferably contain every possible sequence the length of the primer, or, in particular, the length of the complementary portion of the primer. Use of random primers is described in U.S. Patent No. 5,043,272 and U.S. Patent No. 6,214,587.

The disclosed primers can have one or more modified nucleotides. Such primers are referred to herein as modified primers. Modified primers have several advantages. First, some forms of modified primers, such as RNA/ 2'-O-methyl RNA chimeric primers, have a higher melting temperature (T_m) than DNA primers. This increases the stability of primer hybridization and will increase strand invasion by the primers. This will lead to more efficient priming. Also, since the primers are made of RNA, they will be exonuclease resistant. Such primers, if tagged with minor groove binders at their 5' end, will also have better strand invasion of the template dsDNA. In addition, RNA primers can also be very useful for WGA from biological samples such as cells or tissue. Since the biological samples contain endogenous RNA, this RNA can be degraded with RNase to generate a pool of random oligomers, which can then be used to prime the polymerase for amplification of the DNA. This eliminates any need to add primers to the reaction. Alternatively, DNase digestion of biological samples can generate a pool of DNA oligo primers for RNA dependent DNA amplification.

Chimeric primers can also be used. Chimeric primers are primers having at least two types of nucleotides, such as both deoxyribonucleotides and ribonucleotides, ribonucleotides and modified nucleotides, or two different types of modified nucleotides. One form of chimeric primer is peptide nucleic acid/nucleic acid primers. For example, 5'-PNA-DNA-3' or 5'-PNA-RNA-3' primers may be used for more efficient strand invasion and polymerization invasion. The DNA and RNA portions of such primers can have random or degenerate sequences. Other forms of chimeric primers are, for example, 5'-(2'-O-Methyl) RNA-RNA-3' or 5'-(2'-O-Methyl) RNA-DNA-3'.

Many modified nucleotides (nucleotide analogs) are known and can be used in oligonucleotides. A nucleotide analog is a nucleotide which contains some type of modification to either the base, sugar, or phosphate moieties. Modifications to the base moiety would include natural and synthetic modifications of A, C, G, and T/U as well as different purine or pyrimidine bases, such as uracil-5-yl, hypoxanthin-9-yl (I), and 2-aminoadenin-9-yl. A modified base includes but is not limited to 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine,

5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Additional base modifications can be found for example in U.S. Pat. No. 3,687,808, Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and Sanghvi, Y. S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S. T. and Lebleu, B. ed., CRC Press, 1993. Certain nucleotide analogs, such as 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine can increase the stability of duplex formation. Other modified bases are those that function as universal bases. Universal bases include 3-nitropyrrole and 5-nitroindole. Universal bases substitute for the normal bases but have no bias in base pairing. That is, universal bases can base pair with any other base. Primers composed, either in whole or in part, of nucleotides with universal bases are useful for reducing or eliminating amplification bias against repeated sequences in a target sample. This would be useful, for example, where a loss of sequence complexity in the amplified products is undesirable. Base modifications often can be combined with for example a sugar modification, such as 2'-O-methoxyethyl, to achieve unique properties such as increased duplex stability. There are numerous United States patents such as 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, which detail and describe a range of base modifications. Each of these patents is herein incorporated by reference.

Nucleotide analogs can also include modifications of the sugar moiety. Modifications to the sugar moiety would include natural modifications of the ribose and deoxyribose as well as synthetic modifications. Sugar modifications include but are not limited to the following modifications at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C1 to C10, alkyl or C2 to C10 alkenyl

and alkynyl. 2' sugar modifications also include but are not limited to $-O[(CH_2)_n O]_m$ CH_3 , $-O(CH_2)_n OCH_3$, $-O(CH_2)_n NH_2$, $-O(CH_2)_n CH_3$, $-O(CH_2)_n -ONH_2$, and $-O(CH_2)_n ON[(CH_2)_n CH_3]_2$, where n and m are from 1 to about 10.

Other modifications at the 2' position include but are not limited to: C1 to C10
5 lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH_3 ,
OCN, Cl, Br, CN, CF_3 , OCF_3 , $SOCH_3$, $SO_2 CH_3$, ONO_2 , NO_2 , N_3 , NH_2 ,
heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted
silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving
the pharmacokinetic properties of an oligonucleotide, or a group for improving the
10 pharmacodynamic properties of an oligonucleotide, and other substituents having
similar properties. Similar modifications may also be made at other positions on the
sugar, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5'
linked oligonucleotides and the 5' position of 5' terminal nucleotide. Modified sugars
would also include those that contain modifications at the bridging ring oxygen, such as
15 CH_2 and S. Nucleotide sugar analogs may also have sugar mimetics such as cyclobutyl
moieties in place of the pentofuranosyl sugar. There are numerous United States
patents that teach the preparation of such modified sugar structures such as 4,981,957;
5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785;
5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;
20 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, each of which is herein
incorporated by reference in its entirety.

Nucleotide analogs can also be modified at the phosphate moiety. Modified
phosphate moieties include but are not limited to those that can be modified so that the
linkage between two nucleotides contains a phosphorothioate, chiral phosphorothioate,
25 phosphorodithioate, phosphotriester, aminoalkylphosphotriester, methyl and other alkyl
phosphonates including 3'-alkylene phosphonate and chiral phosphonates,
phosphinates, phosphoramidates including 3'-amino phosphoramidate and
aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates,
thionoalkylphosphotriesters, and boranophosphates. It is understood that these
30 phosphate or modified phosphate linkages between two nucleotides can be through a
3'-5' linkage or a 2'-5' linkage, and the linkage can contain inverted polarity such as
3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also

included. Numerous United States patents teach how to make and use nucleotides containing modified phosphates and include but are not limited to, 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

It is understood that nucleotide analogs need only contain a single modification, but may also contain multiple modifications within one of the moieties or between different moieties.

Nucleotide substitutes are molecules having similar functional properties to nucleotides, but which do not contain a phosphate moiety, such as peptide nucleic acid (PNA). Nucleotide substitutes are molecules that will recognize and hybridize to complementary nucleic acids in a Watson-Crick or Hoogsteen manner, but which are linked together through a moiety other than a phosphate moiety. Nucleotide substitutes are able to conform to a double helix type structure when interacting with the appropriate target nucleic acid.

Nucleotide substitutes are nucleotides or nucleotide analogs that have had the phosphate moiety and/or sugar moieties replaced. Nucleotide substitutes do not contain a standard phosphorus atom. Substitutes for the phosphate can be for example, short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts. Numerous United States patents disclose how to make and use these types of phosphate replacements and include but are not limited to 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225;

5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

5 It is also understood in a nucleotide substitute that both the sugar and the phosphate moieties of the nucleotide can be replaced, by for example an amide type linkage (aminoethylglycine) (PNA). United States patents 5,539,082; 5,714,331; and 5,719,262 teach how to make and use PNA molecules, each of which is herein incorporated by reference. (See also Nielsen *et al.*, *Science* 254:1497-1500 (1991)).

10 Primers can be comprised of nucleotides and can be made up of different types of nucleotides or the same type of nucleotides. For example, one or more of the nucleotides in a primer can be ribonucleotides, 2'-O-methyl ribonucleotides, or a mixture of ribonucleotides and 2'-O-methyl ribonucleotides; about 10% to about 50% of the nucleotides can be ribonucleotides, 2'-O-methyl ribonucleotides, or a mixture of ribonucleotides and 2'-O-methyl ribonucleotides; about 50% or more of the nucleotides
15 can be ribonucleotides, 2'-O-methyl ribonucleotides, or a mixture of ribonucleotides and 2'-O-methyl ribonucleotides; or all of the nucleotides are ribonucleotides, 2'-O-methyl ribonucleotides, or a mixture of ribonucleotides and 2'-O-methyl ribonucleotides. The nucleotides can be comprised of bases (that is, the base portion of the nucleotide) and can (and normally will) comprise different types of bases. For
20 example, one or more of the bases can be universal bases, such as 3-nitropyrrole or 5-nitroindole; about 10% to about 50% of the bases can be universal bases; about 50% or more of the bases can be universal bases; or all of the bases can be universal bases.

Primers may, but need not, also contain additional sequence at the 5' end of the primer that is not complementary to the target sequence. This sequence is referred to as
25 the non-complementary portion of the primer. The non-complementary portion of the primer, if present, serves to facilitate strand displacement during DNA replication. The non-complementary portion of the primer can also include a functional sequence such as a promoter for an RNA polymerase. The non-complementary portion of a primer may be any length, but is generally 1 to 100 nucleotides long, and preferably 4 to 8
30 nucleotides long. The use of a non-complementary portion is not preferred when random or partially random primers are used for whole genome amplification.

1. Primers for Whole Genome Strand Displacement Amplification

In the case of whole genome strand displacement amplification, it is preferred that a set of primers having random or partially random nucleotide sequences be used. In a nucleic acid sample of significant or substantial complexity, which is the preferred target sequence for WGSDA, specific nucleic acid sequences present in the sample need not be known and the primers need not be designed to be complementary to any particular sequence. Rather, the complexity of the nucleic acid sample results in a large number of different hybridization target sequences in the sample which will be complementary to various primers of random or partially random sequence. The complementary portion of primers for use in WGSDA can be fully randomized, have only a portion that is randomized, or be otherwise selectively randomized.

The number of random base positions in the complementary portion of primers are preferably from 20% to 100% of the total number of nucleotides in the complementary portion of the primers. More preferably the number of random base positions are from 30% to 100% of the total number of nucleotides in the complementary portion of the primers. Most preferably the number of random base positions are from 50% to 100% of the total number of nucleotides in the complementary portion of the primers. Sets of primers having random or partially random sequences can be synthesized using standard techniques by allowing the addition of any nucleotide at each position to be randomized. It is also preferred that the sets of primers are composed of primers of similar length and/or hybridization characteristics.

2. Primers for Multiple Strand Displacement Amplification

In the case of multiple strand displacement amplification, the complementary portion of each primer is designed to be complementary to the hybridization target in the target sequence. In a set of primers, it is preferred that the complementary portion of each primer be complementary to a different portion of the target sequence. It is more preferred that the primers in the set be complementary to adjacent sites in the target sequence. It is also preferred that such adjacent sites in the target sequence are also adjacent to the amplification target in the target sequence.

It is preferred that, when hybridized to a target sequence, the primers in a set of primers are separated from each other. It is preferred that, when hybridized, the

primers in a set of primers are separated from each other by at least 5 bases. It is more preferred that, when hybridized, the primers in a set of primers are separated from each other by at least 10 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by at least 20 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by at least 30 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by at least 40 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by at least 50 bases.

It is preferred that, when hybridized, the primers in a set of primers are separated from each other by no more than about 500 bases. It is more preferred that, when hybridized, the primers in a set of primers are separated from each other by no more than about 400 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by no more than about 300 bases. It is still more preferred that, when hybridized, the primers in a set of primers are separated from each other by no more than about 200 bases. Any combination of the preferred upper and lower limits of separation described above are specifically contemplated, including all intermediate ranges. The primers in a set of primers need not, when hybridized, be separated from each other by the same number of bases. It is preferred that, when hybridized, the primers in a set of primers are separated from each other by about the same number of bases.

The optimal separation distance between primers will not be the same for all DNA polymerases, because this parameter is dependent on the net polymerization rate. A processive DNA polymerase will have a characteristic polymerization rate which may range from 5 to 300 nucleotides per second, and may be influenced by the presence or absence of accessory ssDNA binding proteins and helicases. In the case of a non-processive polymerase, the net polymerization rate will depend on the enzyme concentration, because at higher concentrations there are more re-initiation events and thus the net polymerization rate will be increased. An example of a processive polymerase is ϕ 29 DNA polymerase, which proceeds at 50 nucleotides per second. An example of a non-processive polymerase is Vent exo(-) DNA polymerase, which will

give effective polymerization rates of 4 nucleotides per second at low concentration, or 16 nucleotides per second at higher concentrations.

To obtain an optimal yield in an MSDA reaction, the primer spacing is preferably adjusted to suit the polymerase being used. Long primer spacing is preferred when using a polymerase with a rapid polymerization rate. Shorter primer spacing is preferred when using a polymerase with a slower polymerization rate. The following assay can be used to determine optimal spacing with any polymerase. The assay uses sets of primers, with each set made up of 5 left primers and 5 right primers. The sets of primers are designed to hybridize adjacent to the same target sequence with each of the different sets of primers having a different primer spacing. The spacing is varied systematically between the sets of primers in increments of 25 nucleotides within the range of 25 nucleotides to 400 nucleotides (the spacing of the primers within each set is the same). A series of reactions are performed in which the same target sequence is amplified using the different sets of primers. The spacing that produces the highest experimental yield of DNA is the optimal primer spacing for the specific DNA polymerase, or DNA polymerase plus accessory protein combination being used.

DNA replication initiated at the sites in the target sequence where the primers hybridize will extend to and displace strands being replicated from primers hybridized at adjacent sites. Displacement of an adjacent strand makes it available for hybridization to another primer and subsequent initiation of another round of replication. The region(s) of the target sequence to which the primers hybridize is referred to as the hybridization target of the target sequence.

A set of primers can include any desired number of primers of different nucleotide sequence. For MSDA, it is preferred that a set of primers include a plurality of primers. It is more preferred that a set of primers include 3 or more primers. It is still more preferred that a set of primers include 4 or more, 5 or more, 6 or more, or 7 or more primers. In general, the more primers used, the greater the level of amplification that will be obtained. There is no fundamental upper limit to the number of primers that a set of primers can have. However, for a given target sequence, the number of primers in a set of primers will generally be limited to the number of hybridization sites available in the target sequence. For example, if the target sequence is a 10,000 nucleotide DNA molecule and 20 nucleotide primers are used, there are 500 non-

overlapping 20 nucleotide sites in the target sequence. Even more primers than this could be used if overlapping sites are either desired or acceptable. It is preferred that a set of primers include no more than about 300 primers. It is preferred that a set of primers include no more than about 200 primers. It is still more preferred that a set of primers include no more than about 100 primers. It is more preferred that a set of primers include no more than about 50 primers. It is most preferred that a set of primers include from 7 to about 50 primers. Any combination of the preferred upper and lower limits for the number of primers in a set of primers described above are specifically contemplated, including all intermediate ranges.

10 A preferred form of primer set for use in MSDA includes two sets of primers, referred to as a right set of primers and a left set of primers. The right set of primers and left set of primers are designed to be complementary to opposite strands of a target sequence. It is preferred that the complementary portions of the right set of primers are each complementary to the right hybridization target, and that each is complementary to a different portion of the right hybridization target. It is preferred that the complementary portions of the left set of primers are each complementary to the left hybridization target, and that each is complementary to a different portion of the left hybridization target. The right and left hybridization targets flank opposite ends of the amplification target in a target sequence. It is preferred that a right set of primers and a left set of primers each include a preferred number of primers as described above for a set of primers. Specifically, it is preferred that a right or left set of primers include a plurality of primers. It is more preferred that a right or left set of primers include 3 or more primers. It is still more preferred that a right or left set of primers include 4 or more, 5 or more, 6 or more, or 7 or more primers. It is preferred that a right or left set of primers include no more than about 200 primers. It is more preferred that a right or left set of primers include no more than about 100 primers. It is most preferred that a right or left set of primers include from 7 to about 100 primers. Any combination of the preferred upper and lower limits for the number of primers in a set of primers described above are specifically contemplated, including all intermediate ranges. It is also preferred that, for a given target sequence, the right set of primers and the left set of primers include the same number of primers. It is also preferred that, for a given

target sequence, the right set of primers and the left set of primers are composed of primers of similar length and/or hybridization characteristics.

Where the target sequence(s) are present in mixed sample--for example, a nosocomial sample containing both human and non-human nucleic acids—the primers used can be specific for the nucleic acids of interest. Thus, to detect pathogen (that is, non-human) nucleic acids in a patient sample, primers specific to pathogen nucleic acids can be used. If human nucleic acids are to be detected, then primers specific to human nucleic acids can be used. In this context, primers specific for particular target nucleic acids or target sequences or a particular class of target nucleic acids or target sequences refer to primers that support amplification of the target nucleic acids and target sequences but do not support substantial amplification of non-target nucleic acids or sequences that are in the relevant sample.

3. Detection Tags

The non-complementary portion of a primer can include sequences to be used to further manipulate or analyze amplified sequences. An example of such a sequence is a detection tag, which is a specific nucleotide sequence present in the non-complementary portion of a primer. Detection tags have sequences complementary to detection probes. Detection tags can be detected using their cognate detection probes. Detection tags become incorporated at the ends of amplified strands. The result is amplified DNA having detection tag sequences that are complementary to the complementary portion of detection probes. If present, there may be one, two, three, or more than three detection tags on a primer. It is preferred that a primer have one, two, three or four detection tags. Most preferably, a primer will have one detection tag. Generally, it is preferred that a primer have 10 detection tags or less. There is no fundamental limit to the number of detection tags that can be present on a primer except the size of the primer. When there are multiple detection tags, they may have the same sequence or they may have different sequences, with each different sequence complementary to a different detection probe. It is preferred that a primer contain detection tags that have the same sequence such that they are all complementary to a single detection probe. For some multiplex detection methods, it is preferable that primers contain up to six detection tags and that the detection tag portions have different sequences such that each of the detection tag portions is complementary to a

different detection probe. A similar effect can be achieved by using a set of primers where each has a single different detection tag. The detection tags can each be any length that supports specific and stable hybridization between the detection tags and the detection probe. For this purpose, a length of 10 to 35 nucleotides is preferred, with a
5 detection tag portion 15 to 20 nucleotides long being most preferred.

4. Address Tag

Another example of a sequence that can be included in the non-complementary portion of a primer is an address tag. An address tag has a sequence complementary to an address probe. Address tags become incorporated at the ends of amplified strands.
10 The result is amplified DNA having address tag sequences that are complementary to the complementary portion of address probes. If present, there may be one, or more than one, address tag on a primer. It is preferred that a primer have one or two address tags. Most preferably, a primer will have one address tag. Generally, it is preferred that a primer have 10 address tags or less. There is no fundamental limit to the number
15 of address tags that can be present on a primer except the size of the primer. When there are multiple address tags, they may have the same sequence or they may have different sequences, with each different sequence complementary to a different address probe. It is preferred that a primer contain address tags that have the same sequence such that they are all complementary to a single address probe. The address tag portion
20 can be any length that supports specific and stable hybridization between the address tag and the address probe. For this purpose, a length between 10 and 35 nucleotides long is preferred, with an address tag portion 15 to 20 nucleotides long being most preferred.

C. Lysis Solution

25 In the disclosed method, the cells can be exposed to alkaline conditions by mixing the cells with a lysis solution. A lysis solution is generally a solution that can raise the pH of a cell solution sufficiently to cause cell lysis. Denaturing solutions can be used as lysis solutions so long as the denaturing solution can have the effects required of lysis solutions. In some embodiments, the lysis solution can comprise a
30 base, such as an aqueous base. Useful bases include potassium hydroxide, sodium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium

bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine, methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, and DBU (1,8-diazobicyclo[5,4,0]undec-7-ene). Useful formulations of lysis solution include lysis solution comprising 400 mM KOH, lysis solution comprising 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA, and lysis solution consisting of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

In some embodiments, the lysis solution can comprise a plurality of basic agents. As used herein, a basic agent is a compound, composition or solution that results in alkaline conditions. In some embodiments, the lysis solution can comprise a buffer. Useful buffers include phosphate buffers, "Good" buffers (such as BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, and TRICINE), sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, and Bis-tris propane. The lysis solution can comprise a plurality of buffering agents. As used herein, a buffering agent is a compound, composition or solution that acts as a buffer. An alkaline buffering agent is a buffering agent that results in alkaline conditions. In some embodiments, the lysis solution can comprise a combination of one or more bases, basic agents, buffers and buffering agents.

The amount of lysis solution mixed with the cells can be that amount that causes a substantial number of cells to lyse or those that cause a sufficient number of cells to lyse. Generally, this volume will be a function of the pH of the cell/lysis solution mixture. Thus, the amount of lysis solution to mix with cells can be determined generally from the volume of the cells and the alkaline concentration of the lysis buffer. For example, a smaller volume of a lysis solution with a stronger base and/or higher concentration of base would be needed to create sufficient alkaline conditions than the volume needed of a lysis solution with a weaker base and/or lower concentration of base. The lysis solution can be formulated such that the cells are mixed with an equal volume of the lysis solution (to produce the desired alkaline conditions).

For example, lysis solutions can be solutions that have a pH of from about 9.0 to about 13.0, from about 9.5 to about 13.0, from about 10.0 to about 13.0, from about

10.5 to about 13.0, from about 11.0 to about 13.0, from about 11.5 to about 13.0, from
about 12.0 to about 13.0, from about 9.0 to about 12.0, from about 9.5 to about 12.0,
from about 10.0 to about 12.0, from about 10.5 to about 12.0, from about 11.0 to about
12.0, from about 11.5 to about 12.0, from about 9.0 to about 11.5, from about 9.5 to
5 about 11.5, from about 10.0 to about 11.5, from about 10.5 to about 11.5, from about
11.0 to about 11.5, from about 9.0 to about 11.0, from about 9.5 to about 11.0, from
about 10.0 to about 11.0, from about 10.5 to about 11.0, from about 9.0 to about 10.5,
from about 9.5 to about 10.5, from about 10.0 to about 10.5, from about 9.0 to about
10.0, from about 9.5 to about 10.0, from about 9.0 to about 9.5, about 9.0, about 9.5,
10 about 10.0, about 10.5, about 11.0, about 11.5, about 12.0, about 12.5, or about 13.0.

Lysis solutions can have, for example, component concentrations of from about
10 mM to about 1 M, from about 10 mM to about 900 mM, from about 10 mM to about
800 mM, from about 10 mM to about 700 mM, from about 10 mM to about 600 mM,
from about 10 mM to about 500 mM, from about 10 mM to about 400 mM, from about
15 10 mM to about 300 mM, from about 10 mM to about 200 mM, from about 10 mM to
about 100 mM, from about 10 mM to about 90 mM, from about 10 mM to about 80
mM, from about 10 mM to about 70 mM, from about 10 mM to about 60 mM, from
about 10 mM to about 50 mM, from about 10 mM to about 40 mM, from about 10 mM
to about 30 mM, from about 10 mM to about 20 mM, from about 20 mM to about 1 M,
20 from about 20 mM to about 900 mM, from about 20 mM to about 800 mM, from about
20 mM to about 700 mM, from about 20 mM to about 600 mM, from about 20 mM to
about 500 mM, from about 20 mM to about 400 mM, from about 20 mM to about 300
mM, from about 20 mM to about 200 mM, from about 20 mM to about 100 mM, from
about 20 mM to about 90 mM, from about 20 mM to about 80 mM, from about 20 mM
25 to about 70 mM, from about 20 mM to about 60 mM, from about 20 mM to about 50
mM, from about 20 mM to about 40 mM, from about 20 mM to about 30 mM, from
about 30 mM to about 1 M, from about 30 mM to about 900 mM, from about 30 mM to
about 800 mM, from about 30 mM to about 700 mM, from about 30 mM to about 600
mM, from about 30 mM to about 500 mM, from about 30 mM to about 400 mM, from
30 about 30 mM to about 300 mM, from about 30 mM to about 200 mM, from about 30
mM to about 100 mM, from about 30 mM to about 90 mM, from about 30 mM to about
80 mM, from about 30 mM to about 70 mM, from about 30 mM to about 60 mM, from

about 30 mM to about 50 mM, from about 30 mM to about 40 mM, from about 40 mM to about 1 M, from about 40 mM to about 900 mM, from about 40 mM to about 800 mM, from about 40 mM to about 700 mM, from about 40 mM to about 600 mM, from about 40 mM to about 500 mM, from about 40 mM to about 400 mM, from about 40 mM to about 300 mM, from about 40 mM to about 200 mM, from about 40 mM to about 100 mM, from about 40 mM to about 90 mM, from about 40 mM to about 80 mM, from about 40 mM to about 70 mM, from about 40 mM to about 60 mM, from about 40 mM to about 50 mM, from about 50 mM to about 1 M, from about 50 mM to about 900 mM, from about 50 mM to about 800 mM, from about 50 mM to about 700 mM, from about 50 mM to about 600 mM, from about 50 mM to about 500 mM, from about 50 mM to about 400 mM, from about 50 mM to about 300 mM, from about 50 mM to about 200 mM, from about 50 mM to about 100 mM, from about 50 mM to about 90 mM, from about 50 mM to about 80 mM, from about 50 mM to about 70 mM, from about 50 mM to about 60 mM, from about 60 mM to about 1 M, from about 60 mM to about 900 mM, from about 60 mM to about 800 mM, from about 60 mM to about 700 mM, from about 60 mM to about 600 mM, from about 60 mM to about 500 mM, from about 60 mM to about 400 mM, from about 60 mM to about 300 mM, from about 60 mM to about 200 mM, from about 60 mM to about 100 mM, from about 60 mM to about 90 mM, from about 60 mM to about 80 mM, from about 60 mM to about 70 mM, from about 70 mM to about 1 M, from about 70 mM to about 900 mM, from about 70 mM to about 800 mM, from about 70 mM to about 700 mM, from about 70 mM to about 600 mM, from about 70 mM to about 500 mM, from about 70 mM to about 400 mM, from about 70 mM to about 300 mM, from about 70 mM to about 200 mM, from about 70 mM to about 100 mM, from about 70 mM to about 90 mM, from about 70 mM to about 80 mM, from about 80 mM to about 1 M, from about 80 mM to about 900 mM, from about 80 mM to about 800 mM, from about 80 mM to about 700 mM, from about 80 mM to about 600 mM, from about 80 mM to about 500 mM, from about 80 mM to about 400 mM, from about 80 mM to about 300 mM, from about 80 mM to about 200 mM, from about 80 mM to about 100 mM, from about 80 mM to about 90 mM, from about 90 mM to about 1 M, from about 90 mM to about 900 mM, from about 90 mM to about 800 mM, from about 90 mM to about 700 mM, from about 90 mM to about 600 mM, from about 90 mM to about 500 mM, from about 90 mM to

about 400 mM, from about 90 mM to about 300 mM, from about 90 mM to about 200 mM, from about 90 mM to about 100 mM, from about 100 mM to about 1 M, from about 100 mM to about 900 mM, from about 100 mM to about 800 mM, from about 100 mM to about 700 mM, from about 100 mM to about 600 mM, from about 100 mM to about 500 mM, from about 100 mM to about 400 mM, from about 100 mM to about 300 mM, from about 100 mM to about 200 mM, from about 200 mM to about 1 M, from about 200 mM to about 900 mM, from about 200 mM to about 800 mM, from about 200 mM to about 700 mM, from about 200 mM to about 600 mM, from about 200 mM to about 500 mM, from about 200 mM to about 400 mM, from about 200 mM to about 300 mM, from about 300 mM to about 1 M, from about 300 mM to about 900 mM, from about 300 mM to about 800 mM, from about 300 mM to about 700 mM, from about 300 mM to about 600 mM, from about 300 mM to about 500 mM, from about 300 mM to about 400 mM, from about 400 mM to about 1 M, from about 400 mM to about 900 mM, from about 400 mM to about 800 mM, from about 400 mM to about 700 mM, from about 400 mM to about 600 mM, from about 400 mM to about 500 mM, from about 500 mM to about 1 M, from about 500 mM to about 900 mM, from about 500 mM to about 800 mM, from about 500 mM to about 700 mM, from about 500 mM to about 600 mM, from about 600 mM to about 1 M, from about 600 mM to about 900 mM, from about 600 mM to about 800 mM, from about 600 mM to about 700 mM, from about 700 mM to about 1 M, from about 700 mM to about 900 mM, from about 700 mM to about 800 mM, from about 800 mM to about 1 M, from about 800 mM to about 900 mM, from about 900 mM to about 1 M, about 10 mM, about 20 mM, about 30 mM, about 40 mM, about 50 mM, about 60 mM, about 70 mM, about 80 mM, about 90 mM, about 100 mM, about 200 mM, about 300 mM, about 400 mM, about 500 mM, about 600 mM, about 700 mM, about 800 mM, about 900 mM, or about 1 M.

The lysis solution can be composed of multiple solutions and/or components that can be added to cells separately or combined in different combinations prior to addition to cells. Thus, for example, a solution of 400 mM KOH and 10 mM EDTA and a solution of 100 mM dithiothreitol can be added to the cells separately. Similarly, the disclosed kits can be composed of multiple solutions and/or components to be

combined to form a lysis solution prior to addition to cells or for separate addition to cells.

D. Stabilization Solution

In the disclosed method, the pH of the cell lysate can be reduced to form a stabilized cell lysate. A stabilization solution is generally a solution that can reduce the pH of a cell lysate exposed to alkaline conditions as described elsewhere herein. In some embodiments, the stabilization solution can comprise an acid. Useful acids include hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, and any carboxylic acid (ethanoic, propanoic, butanoic, etc., including both linear or branched chain carboxylic acids). In some embodiments, the stabilization solution can comprise a buffer. Useful buffers include Tris-HCl, HEPES, "Good" buffers (such as BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, and TRICINE), sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, and Bis-tris propane. Useful formulations of stabilization solutions include stabilization solution comprising 800 mM Tris-HCl; stabilization solution comprising 800 mM Tris-HCl at pH 4.1, and stabilization solution consisting of 800 mM Tris-HCl, pH 4.1.

In some embodiments, the stabilization solution can comprise a plurality of acidic agents. As used herein, an acidic agent is a compound, composition or solution that forms an acid in solution. In some embodiments, the stabilization solution can comprise a plurality of buffering agents. An acidic buffering agent is a buffering agent that forms an acid in solution. In some embodiments, the stabilization solution can comprise a combination of one or more acids, acidic agents, buffers and buffering agents.

A stabilized cell lysate is a cell lysate the pH of which is in the neutral range (from about pH 6.0 to about pH 9.0). Useful stabilized cell lysates have a pH that allows replication of nucleic acids in the cell lysate. For example, the pH of the stabilized cell lysate is usefully at a pH at which the DNA polymerase can function. The pH of the cell lysate can be reduced by mixing the cell lysate with a stabilization solution.

The amount of stabilization solution mixed with the cell lysate can be that amount that causes a reduction in pH to the neutral range (or other desired pH value). Generally, this volume will be a function of the pH of the cell lysate/stabilization solution mixture. Thus, the amount of stabilization solution to mix with the cell lysate
5 can be determined generally from the volume of the cell lysate, its pH and buffering capacity, and the acidic concentration of the stabilization buffer. For example, a smaller volume of a stabilization solution with a stronger acid and/or higher concentration of acid would be needed to reduce the pH sufficiently than the volume needed of a stabilization solution with a weaker acid and/or lower concentration of
10 acid. The stabilization solution can be formulated such that the cell lysate is mixed with an equal volume of the stabilization solution (to produce the desired pH).

For example, stabilization solutions can be solutions that have a pH of from about 1.0 to about 6.0, from about 2.0 to about 6.0, from about 3.0 to about 6.0, from about 3.5 to about 6.0, from about 4.0 to about 6.0, from about 4.5 to about 6.0, from
15 about 5.0 to about 6.0, from about 5.5 to about 6.0, from about 1.0 to about 5.5, from about 2.0 to about 5.5, from about 3.0 to about 5.5, from about 3.5 to about 5.5, from about 4.0 to about 5.5, from about 4.5 to about 5.5, from about 5.0 to about 5.5, from about 1.0 to about 5.0, from about 2.0 to about 5.0, from about 3.0 to about 5.0, from about 3.5 to about 5.0, from about 4.0 to about 5.0, from about 4.5 to about 5.0, from
20 about 1.0 to about 4.5, from about 2.0 to about 4.5, from about 3.0 to about 4.5, from about 3.5 to about 4.5, from about 4.0 to about 4.5, from about 1.0 to about 4.0, from about 2.0 to about 4.0, from about 3.0 to about 4.0, from about 3.5 to about 4.0, from about 1.0 to about 3.5, from about 2.0 to about 3.5, from about 3.0 to about 3.5, from about 1.0 to about 3.0, from about 2.0 to about 3.0, from about 1.0 to about 2.5, from
25 about 2.0 to about 2.5, from about 1.0 to about 2.0, about 1.0, about 2.0, about 2.5, about 3.0, about 3.5, about 4.0, about 4.5, about 5.0, about 5.5, or about 6.0.

Stabilization solutions can have, for example, component concentrations of from about 100 mM to about 1 M, from about 100 mM to about 900 mM, from about 100 mM to about 800 mM, from about 100 mM to about 700 mM, from about 100 mM
30 to about 600 mM, from about 100 mM to about 500 mM, from about 100 mM to about 400 mM, from about 100 mM to about 300 mM, from about 100 mM to about 200 mM, from about 200 mM to about 1 M, from about 200 mM to about 900 mM, from about

200 mM to about 800 mM, from about 200 mM to about 700 mM, from about 200 mM to about 600 mM, from about 200 mM to about 500 mM, from about 200 mM to about 400 mM, from about 200 mM to about 300 mM, from about 300 mM to about 1 M, from about 300 mM to about 900 mM, from about 300 mM to about 800 mM, from about 300 mM to about 700 mM, from about 300 mM to about 600 mM, from about 300 mM to about 500 mM, from about 300 mM to about 400 mM, from about 400 mM to about 1 M, from about 400 mM to about 900 mM, from about 400 mM to about 800 mM, from about 400 mM to about 700 mM, from about 400 mM to about 600 mM, from about 400 mM to about 500 mM, from about 500 mM to about 1 M, from about 500 mM to about 900 mM, from about 500 mM to about 800 mM, from about 500 mM to about 700 mM, from about 500 mM to about 600 mM, from about 600 mM to about 1 M, from about 600 mM to about 900 mM, from about 600 mM to about 800 mM, from about 600 mM to about 700 mM, from about 700 mM to about 1 M, from about 700 mM to about 900 mM, from about 700 mM to about 800 mM, from about 800 mM to about 1 M, from about 800 mM to about 900 mM, from about 900 mM to about 1 M, about 100 mM, about 200 mM, about 300 mM, about 400 mM, about 500 mM, about 600 mM, about 700 mM, about 800 mM, about 900 mM, or about 1 M.

The stabilization solution can be composed of multiple solutions and/or components that can be added to cell lysates separately or combined in different combinations prior to addition to cell lysates. Thus, for example, a solution of a buffer and a solution of an acid can be added to the cells separately. Similarly, the disclosed kits can be composed of multiple solutions and/or components to be combined to form a stabilization solution prior to addition to cell lysates or for separate addition to cell lysates.

E. Denaturing Solution

In some forms of the disclosed method, the DNA samples can be exposed to denaturing conditions by mixing the sample with a denaturing solution. A denaturing solution is generally a solution that can raise the pH of a sample sufficiently to cause, in combination with other conditions such as heating, substantial denaturation of DNA in the DNA sample. Substantial denaturation refers to denaturation of 90% or more of the nucleotides in 90% or more of the DNA molecules in a sample. In this context, denaturation of nucleotides refers to unpaired nucleotides whether physically denatured

by treatment or already unpaired in the sample. Lysis solutions can be used as denaturing solutions so long as the lysis solution has the effects required of denaturing solutions.

In some embodiments, the denaturing solution can comprises a base, such as an aqueous base. Useful bases include potassium hydroxide, sodium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine, methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, and DBU (1,8-diazobicyclo[5,4,0]undec-7-ene). Useful formulations of denaturing solution include denaturing solution comprising about 150 mM to about 500 mM NaOH, denaturing solution comprising about 150 mM to about 500 mM NaOH, and denaturing solution consisting of about 150 mM to about 500 mM NaOH.

In some embodiments, the denaturing solution can comprise a plurality of basic agents. As used herein, a basic agent is a compound, composition or solution that results in denaturing conditions. In some embodiments, the denaturing solution can comprise a buffer. Useful buffers include phosphate buffers, "Good" buffers (such as BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, and TRICINE), sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, and Bis-tris propane. The denaturing solution can comprise a plurality of buffering agents. As used herein, a buffering agent is a compound, composition or solution that acts as a buffer. An alkaline buffering agent is a buffering agent that results in alkaline conditions. In some embodiments, the denaturing solution can comprise a combination of one or more bases, basic agents, buffers and buffering agents.

The amount of denaturing solution mixed with the DNA samples can be that amount that causes, in combination with other conditions such as heating, substantial denaturation of DNA in the DNA sample. Generally, this volume will be a function of the pH, ionic strength, and temperature of the sample/denaturing solution mixture. Thus, the amount of denaturing solution to mix with DNA samples can be determined

generally from the volume of the DNA sample, the alkaline concentration of the denaturing buffer, and the temperature to which the resulting mixture will be heated. For example, at a given temperature, a smaller volume of a denaturing solution with a stronger base and/or higher concentration of base would be needed to create sufficient
5 denaturing conditions than the volume needed of a denaturing solution with a weaker base and/or lower concentration of base. The denaturing solution can be formulated such that the DNA samples are mixed with, for example, one tenth volume of the denaturing solution (to produce the desired denaturing conditions).

For example, denaturing solutions can be solutions that have a pH of from about
10 9.0 to about 13.0, from about 9.5 to about 13.0, from about 10.0 to about 13.0, from about 10.5 to about 13.0, from about 11.0 to about 13.0, from about 11.5 to about 13.0, from about 12.0 to about 13.0, from about 9.0 to about 12.0, from about 9.5 to about 12.0, from about 10.0 to about 12.0, from about 10.5 to about 12.0, from about 11.0 to about 12.0, from about 11.5 to about 12.0, from about 9.0 to about 11.5, from about 9.5
15 to about 11.5, from about 10.0 to about 11.5, from about 10.5 to about 11.5, from about 11.0 to about 11.5, from about 9.0 to about 11.0, from about 9.5 to about 11.0, from about 10.0 to about 11.0, from about 10.5 to about 11.0, from about 9.0 to about 10.5, from about 9.5 to about 10.5, from about 10.0 to about 10.5, from about 9.0 to about 10.0, from about 9.5 to about 10.0, from about 9.0 to about 9.5, about 9.0, about 9.5,
20 about 10.0, about 10.5, about 11.0, about 11.5, about 12.0, about 12.5, or about 13.0.

Denaturing solutions can have, for example, component concentrations of from about 10 mM to about 1 M, from about 10 mM to about 900 mM, from about 10 mM to about 800 mM, from about 10 mM to about 700 mM, from about 10 mM to about 600 mM, from about 10 mM to about 500 mM, from about 10 mM to about 400 mM, from
25 about 10 mM to about 300 mM, from about 10 mM to about 200 mM, from about 10 mM to about 100 mM, from about 10 mM to about 90 mM, from about 10 mM to about 80 mM, from about 10 mM to about 70 mM, from about 10 mM to about 60 mM, from about 10 mM to about 50 mM, from about 10 mM to about 40 mM, from about 10 mM to about 30 mM, from about 10 mM to about 20 mM, from about 20 mM to about 1 M,
30 from about 20 mM to about 900 mM, from about 20 mM to about 800 mM, from about 20 mM to about 700 mM, from about 20 mM to about 600 mM, from about 20 mM to about 500 mM, from about 20 mM to about 400 mM, from about 20 mM to about 300

mM, from about 20 mM to about 200 mM, from about 20 mM to about 100 mM, from about 20 mM to about 90 mM, from about 20 mM to about 80 mM, from about 20 mM to about 70 mM, from about 20 mM to about 60 mM, from about 20 mM to about 50 mM, from about 20 mM to about 40 mM, from about 20 mM to about 30 mM, from about 30 mM to about 1 M, from about 30 mM to about 900 mM, from about 30 mM to about 800 mM, from about 30 mM to about 700 mM, from about 30 mM to about 600 mM, from about 30 mM to about 500 mM, from about 30 mM to about 400 mM, from about 30 mM to about 300 mM, from about 30 mM to about 200 mM, from about 30 mM to about 100 mM, from about 30 mM to about 90 mM, from about 30 mM to about 80 mM, from about 30 mM to about 70 mM, from about 30 mM to about 60 mM, from about 30 mM to about 50 mM, from about 30 mM to about 40 mM, from about 40 mM to about 1 M, from about 40 mM to about 900 mM, from about 40 mM to about 800 mM, from about 40 mM to about 700 mM, from about 40 mM to about 600 mM, from about 40 mM to about 500 mM, from about 40 mM to about 400 mM, from about 40 mM to about 300 mM, from about 40 mM to about 200 mM, from about 40 mM to about 100 mM, from about 40 mM to about 90 mM, from about 40 mM to about 80 mM, from about 40 mM to about 70 mM, from about 40 mM to about 60 mM, from about 40 mM to about 50 mM, from about 50 mM to about 1 M, from about 50 mM to about 900 mM, from about 50 mM to about 800 mM, from about 50 mM to about 700 mM, from about 50 mM to about 600 mM, from about 50 mM to about 500 mM, from about 50 mM to about 400 mM, from about 50 mM to about 300 mM, from about 50 mM to about 200 mM, from about 50 mM to about 100 mM, from about 50 mM to about 90 mM, from about 50 mM to about 80 mM, from about 50 mM to about 70 mM, from about 50 mM to about 60 mM, from about 60 mM to about 1 M, from about 60 mM to about 900 mM, from about 60 mM to about 800 mM, from about 60 mM to about 700 mM, from about 60 mM to about 600 mM, from about 60 mM to about 500 mM, from about 60 mM to about 400 mM, from about 60 mM to about 300 mM, from about 60 mM to about 200 mM, from about 60 mM to about 100 mM, from about 60 mM to about 90 mM, from about 60 mM to about 80 mM, from about 60 mM to about 70 mM, from about 70 mM to about 1 M, from about 70 mM to about 900 mM, from about 70 mM to about 800 mM, from about 70 mM to about 700 mM, from about 70 mM to about 600 mM, from about 70 mM to about 500 mM, from about 70 mM to

about 400 mM, from about 70 mM to about 300 mM, from about 70 mM to about 200 mM, from about 70 mM to about 100 mM, from about 70 mM to about 90 mM, from about 70 mM to about 80 mM, from about 80 mM to about 1 M, from about 80 mM to about 900 mM, from about 80 mM to about 800 mM, from about 80 mM to about 700 mM, from about 80 mM to about 600 mM, from about 80 mM to about 500 mM, from about 80 mM to about 400 mM, from about 80 mM to about 300 mM, from about 80 mM to about 200 mM, from about 80 mM to about 100 mM, from about 80 mM to about 90 mM, from about 90 mM to about 1 M, from about 90 mM to about 900 mM, from about 90 mM to about 800 mM, from about 90 mM to about 700 mM, from about 90 mM to about 600 mM, from about 90 mM to about 500 mM, from about 90 mM to about 400 mM, from about 90 mM to about 300 mM, from about 90 mM to about 200 mM, from about 90 mM to about 100 mM, from about 100 mM to about 1 M, from about 100 mM to about 900 mM, from about 100 mM to about 800 mM, from about 100 mM to about 700 mM, from about 100 mM to about 600 mM, from about 100 mM to about 500 mM, from about 100 mM to about 400 mM, from about 100 mM to about 300 mM, from about 100 mM to about 200 mM, from about 200 mM to about 1 M, from about 200 mM to about 900 mM, from about 200 mM to about 800 mM, from about 200 mM to about 700 mM, from about 200 mM to about 600 mM, from about 200 mM to about 500 mM, from about 200 mM to about 400 mM, from about 200 mM to about 300 mM, from about 300 mM to about 1 M, from about 300 mM to about 900 mM, from about 300 mM to about 800 mM, from about 300 mM to about 700 mM, from about 300 mM to about 600 mM, from about 300 mM to about 500 mM, from about 300 mM to about 400 mM, from about 400 mM to about 1 M, from about 400 mM to about 900 mM, from about 400 mM to about 800 mM, from about 400 mM to about 700 mM, from about 400 mM to about 600 mM, from about 400 mM to about 500 mM, from about 500 mM to about 1 M, from about 500 mM to about 900 mM, from about 500 mM to about 800 mM, from about 500 mM to about 700 mM, from about 500 mM to about 600 mM, from about 600 mM to about 1 M, from about 600 mM to about 900 mM, from about 600 mM to about 800 mM, from about 600 mM to about 700 mM, from about 700 mM to about 1 M, from about 700 mM to about 900 mM, from about 700 mM to about 800 mM, from about 800 mM to about 1 M, from about 800 mM to about 900 mM, from about 900 mM to about 1 M, about 10 mM,

about 20 mM, about 30 mM, about 40 mM, about 50 mM, about 60 mM, about 70 mM, about 80 mM, about 90 mM, about 100 mM, about 200 mM, about 300 mM, about 400 mM, about 500 mM, about 600 mM, about 700 mM, about 800 mM, about 900 mM, or about 1 M.

5 The denaturing solution can be composed of multiple solutions and/or components that can be added to DNA samples separately or combined in different combinations prior to addition to DNA samples. Thus, for example, a solution of a buffer and a solution of a base can be added to the samples separately. Similarly, the disclosed kits can be composed of multiple solutions and/or components to be
10 combined to form a denaturing solution prior to addition to DNA samples or for separate addition to samples.

F. Nucleic Acid Fingerprints

 The disclosed method can be used to produce replicated strands that serve as a nucleic acid fingerprint of a complex sample of nucleic acid. Such a nucleic acid
15 fingerprint can be compared with other, similarly prepared nucleic acid fingerprints of other nucleic acid samples to allow convenient detection of differences between the samples. The nucleic acid fingerprints can be used both for detection of related nucleic acid samples and comparison of nucleic acid samples. For example, the presence or identity of specific organisms can be detected by producing a nucleic acid fingerprint of
20 the test organism and comparing the resulting nucleic acid fingerprint with reference nucleic acid fingerprints prepared from known organisms. Changes and differences in gene expression patterns can also be detected by preparing nucleic acid fingerprints of mRNA from different cell samples and comparing the nucleic acid fingerprints. The replicated strands can also be used to produce a set of probes or primers that is specific
25 for the source of a nucleic acid sample. The replicated strands can also be used as a library of nucleic acid sequences present in a sample. Nucleic acid fingerprints can be made up of, or derived from, whole genome amplification of a sample such that the entire relevant nucleic acid content of the sample is substantially represented, or from multiple strand displacement amplification of selected target sequences within a
30 sample.

 Nucleic acid fingerprints can be stored or archived for later use. For example, replicated strands produced in the disclosed method can be physically stored, either in

solution, frozen, or attached or adhered to a solid-state substrate such as an array. Storage in an array is useful for providing an archived probe set derived from the nucleic acids in any sample of interest. As another example, informational content of, or derived from, nucleic acid fingerprints can also be stored. Such information can be stored, for example, in or as computer readable media. Examples of informational content of nucleic acid fingerprints include nucleic acid sequence information (complete or partial); differential nucleic acid sequence information such as sequences present in one sample but not another; hybridization patterns of replicated strands to, for example, nucleic acid arrays, sets, chips, or other replicated strands. Numerous other data that is or can be derived from nucleic acid fingerprints and replicated strands produced in the disclosed method can also be collected, used, saved, stored, and/or archived.

Nucleic acid fingerprints can also contain or be made up of other information derived from the information generated in the disclosed method, and can be combined with information obtained or generated from any other source. The informational nature of nucleic acid fingerprints produced using the disclosed method lends itself to combination and/or analysis using known bioinformatics systems and methods.

Nucleic acid fingerprints of nucleic acid samples can be compared to a similar nucleic acid fingerprint derived from any other sample to detect similarities and differences in the samples (which is indicative of similarities and differences in the nucleic acids in the samples). For example, a nucleic acid fingerprint of a first nucleic acid sample can be compared to a nucleic acid fingerprint of a sample from the same type of organism as the first nucleic acid sample, a sample from the same type of tissue as the first nucleic acid sample, a sample from the same organism as the first nucleic acid sample, a sample obtained from the same source but at time different from that of the first nucleic acid sample, a sample from an organism different from that of the first nucleic acid sample, a sample from a type of tissue different from that of the first nucleic acid sample, a sample from a strain of organism different from that of the first nucleic acid sample, a sample from a species of organism different from that of the first nucleic acid sample, or a sample from a type of organism different from that of the first nucleic acid sample.

The same type of tissue is tissue of the same type such as liver tissue, muscle tissue, or skin (which may be from the same or a different organism or type of organism). The same organism refers to the same individual, animal, or cell. For example, two samples taken from a patient are from the same organism. The same source is similar but broader, referring to samples from, for example, the same organism, the same tissue from the same organism, the same DNA molecule, or the same DNA library. Samples from the same source that are to be compared can be collected at different times (thus allowing for potential changes over time to be detected). This is especially useful when the effect of a treatment or change in condition is to be assessed. Samples from the same source that have undergone different treatments can also be collected and compared using the disclosed method. A different organism refers to a different individual organism, such as a different patient, a different individual animal. Different organism includes a different organism of the same type or organisms of different types. A different type of organism refers to organisms of different types such as a dog and cat, a human and a mouse, or *E. coli* and *Salmonella*. A different type of tissue refers to tissues of different types such as liver and kidney, or skin and brain. A different strain or species of organism refers to organisms differing in their species or strain designation as those terms are understood in the art.

20 **G. Solid-State Detectors**

Solid-state detectors are solid-state substrates or supports to which address probes or detection molecules have been coupled. A preferred form of solid-state detector is an array detector. An array detector is a solid-state detector to which multiple different address probes or detection molecules have been coupled in an array, grid, or other organized pattern.

Solid-state substrates for use in solid-state detectors can include any solid material to which oligonucleotides can be coupled. This includes materials such as acrylamide, cellulose, nitrocellulose, glass, gold, polystyrene, polyethylene vinyl acetate, polypropylene, polymethacrylate, polyethylene, polyethylene oxide, glass, polysilicates, polycarbonates, teflon, fluorocarbons, nylon, silicon rubber, polyanhydrides, polyglycolic acid, polylactic acid, polyorthoesters, functionalized silane, polypropylfumerate, collagen, glycosaminoglycans, and polyamino acids.

Solid-state substrates can have any useful form including thin films or membranes, beads, bottles, dishes, fibers, optical fibers, woven fibers, chips, compact disks, shaped polymers, particles and microparticles. A chip is a rectangular or square small piece of material. Preferred forms for solid-state substrates are thin films, beads, or chips.

5 Address probes immobilized on a solid-state substrate allow capture of the products of the disclosed amplification method on a solid-state detector. Such capture provides a convenient means of washing away reaction components that might interfere with subsequent detection steps. By attaching different address probes to different regions of a solid-state detector, different amplification products can be captured at
10 different, and therefore diagnostic, locations on the solid-state detector. For example, in a multiplex assay, address probes specific for numerous different amplified nucleic acids (each representing a different target sequence amplified via a different set of primers) can be immobilized in an array, each in a different location. Capture and detection will occur only at those array locations corresponding to amplified nucleic
15 acids for which the corresponding target sequences were present in a sample.

 Methods for immobilization of oligonucleotides to solid-state substrates are well established. Oligonucleotides, including address probes and detection probes, can be coupled to substrates using established coupling methods. For example, suitable attachment methods are described by Pease *et al.*, *Proc. Natl. Acad. Sci. USA*
20 **91**(11):5022-5026 (1994), and Khrapko *et al.*, *Mol Biol (Mosk) (USSR)* 25:718-730 (1991). A method for immobilization of 3'-amine oligonucleotides on casein-coated slides is described by Stimpson *et al.*, *Proc. Natl. Acad. Sci. USA* **92**:6379-6383 (1995). A preferred method of attaching oligonucleotides to solid-state substrates is described by Guo *et al.*, *Nucleic Acids Res.* 22:5456-5465 (1994). Examples of nucleic acid chips
25 and arrays, including methods of making and using such chips and arrays, are described in U.S. Patent No. 6,287,768, U.S. Patent No. 6,288,220, U.S. Patent No. 6,287,776, U.S. Patent No. 6,297,006, and U.S. Patent No. 6,291,193.

H. Solid-State Samples

 Solid-state samples are solid-state substrates or supports to which target
30 sequences or MDA products (that is, replicated strands) have been coupled or adhered. Target sequences are preferably delivered in a target sample or assay sample. A preferred form of solid-state sample is an array sample. An array sample is a solid-state

sample to which multiple different target sequences have been coupled or adhered in an array, grid, or other organized pattern.

Solid-state substrates for use in solid-state samples can include any solid material to which target sequences can be coupled or adhered. This includes materials
5 such as acrylamide, cellulose, nitrocellulose, glass, gold, polystyrene, polyethylene vinyl acetate, polypropylene, polymethacrylate, polyethylene, polyethylene oxide, glass, polysilicates, polycarbonates, teflon, fluorocarbons, nylon, silicon rubber, polyanhydrides, polyglycolic acid, polylactic acid, polyorthoesters, functionalized silane, polypropylfumerate, collagen, glycosaminoglycans, and polyamino acids.
10 Solid-state substrates can have any useful form including thin films or membranes, beads, bottles, dishes, slides, fibers, optical fibers, woven fibers, chips, compact disks, shaped polymers, particles and microparticles. A chip is a rectangular or square small piece of material. Preferred forms for solid-state substrates are thin films, beads, or chips.

15 Target sequences immobilized on a solid-state substrate allow formation of target-specific amplified nucleic acid localized on the solid-state substrate. Such localization provides a convenient means of washing away reaction components that might interfere with subsequent detection steps, and a convenient way of assaying multiple different samples simultaneously. Amplified nucleic acid can be
20 independently formed at each site where a different sample is adhered. For immobilization of target sequences or other oligonucleotide molecules to form a solid-state sample, the methods described above can be used. Nucleic acids produced in the disclosed method can be coupled or adhered to a solid-state substrate in any suitable way. For example, nucleic acids generated by multiple strand displacement can be
25 attached by adding modified nucleotides to the 3' ends of nucleic acids produced by strand displacement replication using terminal deoxynucleotidyl transferase, and reacting the modified nucleotides with a solid-state substrate or support thereby attaching the nucleic acids to the solid-state substrate or support.

A preferred form of solid-state substrate is a glass slide to which up to 256
30 separate target samples have been adhered as an array of small dots. Each dot is preferably from 0.1 to 2.5 mm in diameter, and most preferably around 2.5 mm in diameter. Such microarrays can be fabricated, for example, using the method described

by Schena *et al.*, *Science* 270:487-470 (1995). Briefly, microarrays can be fabricated on poly-L-lysine-coated microscope slides (Sigma) with an arraying machine fitted with one printing tip. The tip is loaded with 1 μ l of a DNA sample (0.5 mg/ml) from, for example, 96-well microtiter plates and deposited \sim 0.005 μ l per slide on multiple
5 slides at the desired spacing. The printed slides can then be rehydrated for 2 hours in a humid chamber, snap-dried at 100°C for 1 minute, rinsed in 0.1% SDS, and treated with 0.05% succinic anhydride prepared in buffer consisting of 50% 1-methyl-2-pyrrolidinone and 50% boric acid. The DNA on the slides can then be denatured in, for example, distilled water for 2 minutes at 90°C immediately before use. Microarray
10 solid-state samples can scanned with, for example, a laser fluorescent scanner with a computer-controlled XY stage and a microscope objective. A mixed gas, multiline laser allows sequential excitation of multiple fluorophores.

I. Detection Labels

To aid in detection and quantitation of nucleic acids amplified using the
15 disclosed method, detection labels can be directly incorporated into amplified nucleic acids or can be coupled to detection molecules. As used herein, a detection label is any molecule that can be associated with amplified nucleic acid, directly or indirectly, and which results in a measurable, detectable signal, either directly or indirectly. Many such labels for incorporation into nucleic acids or coupling to nucleic acid probes are
20 known to those of skill in the art. Examples of detection labels suitable for use in the disclosed method are radioactive isotopes, fluorescent molecules, phosphorescent molecules, enzymes, antibodies, and ligands.

Examples of suitable fluorescent labels include fluorescein isothiocyanate (FITC), 5,6-carboxymethyl fluorescein, Texas red, nitrobenz-2-oxa-1,3-diazol-4-yl
25 (NBD), coumarin, dansyl chloride, rhodamine, amino-methyl coumarin (AMCA), Eosin, Erythrosin, BODIPY[®], Cascade Blue[®], Oregon Green[®], pyrene, lissamine, xanthenes, acridines, oxazines, phycoerythrin, macrocyclic chelates of lanthanide ions such as quantum dye[™], fluorescent energy transfer dyes, such as thiazole orange-ethidium heterodimer, and the cyanine dyes Cy3, Cy3.5, Cy5, Cy5.5 and Cy7.
30 Examples of other specific fluorescent labels include 3-Hydroxypyrene 5,8,10-Tri Sulfonic acid, 5-Hydroxy Tryptamine (5-HT), Acid Fuchsin, Alizarin Complexon, Alizarin Red, Allophycocyanin, Aminocoumarin, Anthroyl Stearate, Astrazon Brilliant

Red 4G, Astrazon Orange R, Astrazon Red 6B, Astrazon Yellow 7 GLL, Atabrine,
 Auramine, Auroposphine, Auroposphine G, BAO 9 (Bisaminophenyloxadiazole),
 BCECF, Berberine Sulphate, Bisbenzamide, Blancophor FFG Solution, Blancophor
 SV, Bodipy F1, Brilliant Sulphoflavin FF, Calcien Blue, Calcium Green, Calcofluor
 5 RW Solution, Calcofluor White, Calcophor White ABT Solution, Calcophor White
 Standard Solution, Carbostyryl, Cascade Yellow, Catecholamine, Chinacrine,
 Coriphosphine O, Coumarin-Phalloidin, CY3.1 8, CY5.1 8, CY7, Dans (1-Dimethyl
 Amino Naphaline 5 Sulphonic Acid), Dansa (Diamino Naphtyl Sulphonic Acid),
 Dansyl NH-CH₃, Diamino Phenyl Oxydiazole (DAO), Dimethylamino-5-Sulphonic
 10 acid, Dipyrrometheneboron Difluoride, Diphenyl Brilliant Flavine 7GFF, Dopamine,
 Erythrosin ITC, Euchrysin, FIF (Formaldehyde Induced Fluorescence), Flazo Orange,
 Fluo 3, Fluorescamine, Fura-2, Genacryl Brilliant Red B, Genacryl Brilliant Yellow
 10GF, Genacryl Pink 3G, Genacryl Yellow 5GF, Gloxalic Acid, Granular Blue,
 Haematoporphyrin, Indo-1, Intrawhite Cf Liquid, Leucophor PAF, Leucophor SF,
 15 Leucophor WS, Lissamine Rhodamine B200 (RD200), Lucifer Yellow CH, Lucifer
 Yellow VS, Magdala Red, Marina Blue, Maxilon Brilliant Flavin 10 GFF, Maxilon
 Brilliant Flavin 8 GFF, MPS (Methyl Green Pyronine Stilbene), Mithramycin, NBD
 Amine, Nitrobenzoxadidole, Noradrenaline, Nuclear Fast Red, Nuclear Yellow,
 Nylosan Brilliant Flavin E8G, Oxadiazole, Pacific Blue, Pararosanine (Feulgen),
 20 Phorwite AR Solution, Phorwite BKL, Phorwite Rev, Phorwite RPA, Phosphine 3R,
 Phthalocyanine, Phycoerythrin R, Polyazaindacene Pontochrome Blue Black,
 Porphyrin, Primuline, Procion Yellow, Pyronine, Pyronine B, Pyrozal Brilliant Flavin
 7GF, Quinacrine Mustard, Rhodamine 123, Rhodamine 5 GLD, Rhodamine 6G,
 Rhodamine B, Rhodamine B 200, Rhodamine B Extra, Rhodamine BB, Rhodamine
 25 BG, Rhodamine WT, Serotonin, Sevron Brilliant Red 2B, Sevron Brilliant Red 4G,
 Sevron Brilliant Red B, Sevron Orange, Sevron Yellow L, SITS (Primuline), SITS
 (Stilbene Isothiosulphonic acid), Stilbene, Snarf 1, sulpho Rhodamine B Can C, Sulpho
 Rhodamine G Extra, Tetracycline, Thiazine Red R, Thioflavin S, Thioflavin TCN,
 Thioflavin 5, Thiolyte, Thiozol Orange, Tinopol CBS, True Blue, Ultralite, Uranine B,
 30 Uvitex SFC, Xylene Orange, and XRITC.

Preferred fluorescent labels are fluorescein (5-carboxyfluorescein-N-
 hydroxysuccinimide ester), rhodamine (5,6-tetramethyl rhodamine), and the cyanine

dyes Cy3, Cy3.5, Cy5, Cy5.5 and Cy7. The absorption and emission maxima, respectively, for these fluors are: FITC (490 nm; 520 nm), Cy3 (554 nm; 568 nm), Cy3.5 (581 nm; 588 nm), Cy5 (652 nm; 672 nm), Cy5.5 (682 nm; 703 nm) and Cy7 (755 nm; 778 nm), thus allowing their simultaneous detection. Other examples of fluorescein dyes include 6-carboxyfluorescein (6-FAM), 2',4',1,4,-
5 tetrachlorofluorescein (TET), 2',4',5',7',1,4-hexachlorofluorescein (HEX), 2',7'-dimethoxy-4', 5'-dichloro-6-carboxyrhodamine (JOE), 2'-chloro-5'-fluoro-7',8'-fused phenyl-1,4-dichloro-6-carboxyfluorescein (NED), and 2'-chloro-7'-phenyl-1,4-dichloro-6-carboxyfluorescein (VIC). Fluorescent labels can be obtained from a variety of
10 commercial sources, including Amersham Pharmacia Biotech, Piscataway, NJ; Molecular Probes, Eugene, OR; and Research Organics, Cleveland, Ohio.

Additional labels of interest include those that provide for signal only when the probe with which they are associated is specifically bound to a target molecule, where such labels include: "molecular beacons" as described in Tyagi & Kramer, *Nature*
15 *Biotechnology* (1996) 14:303 and EP 0 070 685 B1. Other labels of interest include those described in U.S. Pat. No. 5,563,037; WO 97/17471 and WO 97/17076.

Labeled nucleotides are a preferred form of detection label since they can be directly incorporated into the amplification products during synthesis. Examples of detection labels that can be incorporated into amplified nucleic acids include nucleotide
20 analogs such as BrdUrd (5-bromodeoxyuridine, Hoy and Schimke, *Mutation Research* **290**:217-230 (1993)), aminoallyldeoxyuridine (Henegariu *et al.*, *Nature Biotechnology* **18**:345-348 (2000)), 5-methylcytosine (Sano *et al.*, *Biochim. Biophys. Acta* **951**:157-165 (1988)), bromouridine (Wansick *et al.*, *J. Cell Biology* **122**:283-293 (1993)) and nucleotides modified with biotin (Langer *et al.*, *Proc. Natl. Acad. Sci. USA* **78**:6633
25 (1981)) or with suitable haptens such as digoxigenin (Kerkhof, *Anal. Biochem.* **205**:359-364 (1992)). Suitable fluorescence-labeled nucleotides are Fluorescein-isothiocyanate-dUTP, Cyanine-3-dUTP and Cyanine-5-dUTP (Yu *et al.*, *Nucleic Acids Res.*, **22**:3226-3232 (1994)). A preferred nucleotide analog detection label for DNA is BrdUrd (bromodeoxyuridine, BrdUrd, BrdU, BUdR, Sigma-Aldrich Co). Other
30 preferred nucleotide analogs for incorporation of detection label into DNA are AA-dUTP (aminoallyl-deoxyuridine triphosphate, Sigma-Aldrich Co.), and 5-methyl-dCTP (Roche Molecular Biochemicals). A preferred nucleotide analog for incorporation of

detection label into RNA is biotin-16-UTP (biotin-16-uridine-5'-triphosphate, Roche Molecular Biochemicals). Fluorescein, Cy3, and Cy5 can be linked to dUTP for direct labelling. Cy3.5 and Cy7 are available as avidin or anti-digoxigenin conjugates for secondary detection of biotin- or digoxigenin-labelled probes.

5 Detection labels that are incorporated into amplified nucleic acid, such as biotin, can be subsequently detected using sensitive methods well-known in the art. For example, biotin can be detected using streptavidin-alkaline phosphatase conjugate (Tropix, Inc.), which is bound to the biotin and subsequently detected by chemiluminescence of suitable substrates (for example, chemiluminescent substrate
10 CSPD: disodium, 3-(4-methoxyspiro-[1,2,-dioxetane-3-2'-(5'-chloro)tricyclo [3.3.1.1^{3,7}]decane]-4-yl) phenyl phosphate; Tropix, Inc.). Labels can also be enzymes, such as alkaline phosphatase, soybean peroxidase, horseradish peroxidase and polymerases, that can be detected, for example, with chemical signal amplification or by using a substrate to the enzyme which produces light (for example, a
15 chemiluminescent 1,2-dioxetane substrate) or fluorescent signal.

 Molecules that combine two or more of these detection labels are also considered detection labels. Any of the known detection labels can be used with the disclosed probes, tags, and method to label and detect nucleic acid amplified using the disclosed method. Methods for detecting and measuring signals generated by detection
20 labels are also known to those of skill in the art. For example, radioactive isotopes can be detected by scintillation counting or direct visualization; fluorescent molecules can be detected with fluorescent spectrophotometers; phosphorescent molecules can be detected with a spectrophotometer or directly visualized with a camera; enzymes can be detected by detection or visualization of the product of a reaction catalyzed by the
25 enzyme; antibodies can be detected by detecting a secondary detection label coupled to the antibody. As used herein, detection molecules are molecules which interact with amplified nucleic acid and to which one or more detection labels are coupled.

J. Detection Probes

 Detection probes are labeled oligonucleotides having sequence complementary
30 to detection tags on amplified nucleic acids. The complementary portion of a detection probe can be any length that supports specific and stable hybridization between the detection probe and the detection tag. For this purpose, a length of 10 to 35 nucleotides

is preferred, with a complementary portion of a detection probe 16 to 20 nucleotides long being most preferred. Detection probes can contain any of the detection labels described above. Preferred labels are biotin and fluorescent molecules. A particularly preferred detection probe is a molecular beacon. Molecular beacons are detection probes labeled with fluorescent moieties where the fluorescent moieties fluoresce only when the detection probe is hybridized (Tyagi and Kramer, *Nature Biotechnol.* **14**:303-309 (1995)). The use of such probes eliminates the need for removal of unhybridized probes prior to label detection because the unhybridized detection probes will not produce a signal. This is especially useful in multiplex assays.

10 **K. Address Probes**

An address probe is an oligonucleotide having a sequence complementary to address tags on primers. The complementary portion of an address probe can be any length that supports specific and stable hybridization between the address probe and the address tag. For this purpose, a length of 10 to 35 nucleotides is preferred, with a complementary portion of an address probe 12 to 18 nucleotides long being most preferred. An address probe can contain a single complementary portion or multiple complementary portions. Preferably, address probes are coupled, either directly or via a spacer molecule, to a solid-state support. Such a combination of address probe and solid-state support are a preferred form of solid-state detector.

20 **L. Oligonucleotide Synthesis**

Primers, detection probes, address probes, and any other oligonucleotides can be synthesized using established oligonucleotide synthesis methods. Methods to produce or synthesize oligonucleotides are well known in the art. Such methods can range from standard enzymatic digestion followed by nucleotide fragment isolation (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Edition (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) Chapters 5, 6) to purely synthetic methods, for example, by the cyanoethyl phosphoramidite method. Solid phase chemical synthesis of DNA fragments is routinely performed using protected nucleoside cyanoethyl phosphoramidites (S. L. Beaucage *et al.* (1981) *Tetrahedron Lett.* **22**:1859). In this approach, the 3'-hydroxyl group of an initial 5'-protected nucleoside is first covalently attached to the polymer support (R. C. Pless *et al.* (1975) *Nucleic Acids Res.* **2**:773 (1975)). Synthesis of the oligonucleotide then

proceeds by deprotection of the 5'-hydroxyl group of the attached nucleoside, followed by coupling of an incoming nucleoside-3'-phosphoramidite to the deprotected hydroxyl group (M. D. Matteucci et al. (1981) J. Am. Chem. Soc. 103:3185). The resulting phosphite triester is finally oxidized to a phosphotriester to complete the internucleotide bond (R. L. Letsinger et al. (1976) J. Am. Chem. Soc. 98:3655). Alternatively, the synthesis of phosphorothioate linkages can be carried out by sulfurization of the phosphite triester. Several chemicals can be used to perform this reaction, among them 3H-1,2-benzodithiole-3-one, 1,1-dioxide (R.P. Iyer, W. Egan, J.B. Regan, and S.L. Beaucage, J. Am. Chem. Soc., 1990, 112, 1253-1254). The steps of deprotection, coupling and oxidation are repeated until an oligonucleotide of the desired length and sequence is obtained. Other methods exist to generate oligonucleotides such as the H-phosphonate method (Hall et al, (1957) J. Chem. Soc., 3291-3296) or the phosphotriester method as described by Ikuta *et al.*, *Ann. Rev. Biochem.* **53**:323-356 (1984), (phosphotriester and phosphite-triester methods), and Narang *et al.*, *Methods Enzymol.*, **65**:610-620 (1980), (phosphotriester method). Protein nucleic acid molecules can be made using known methods such as those described by Nielsen *et al.*, *Bioconjug. Chem.* **5**:3-7 (1994). Other forms of oligonucleotide synthesis are described in U.S. Patent No. 6,294,664 and U.S. Patent No. 6,291,669.

The nucleotide sequence of an oligonucleotide is generally determined by the sequential order in which subunits of subunit blocks are added to the oligonucleotide chain during synthesis. Each round of addition can involve a different, specific nucleotide precursor, or a mixture of one or more different nucleotide precursors. In general, degenerate or random positions in an oligonucleotide can be produced by using a mixture of nucleotide precursors representing the range of nucleotides that can be present at that position. Thus, precursors for A and T can be included in the reaction for a particular position in an oligonucleotide if that position is to be degenerate for A and T. Precursors for all four nucleotides can be included for a fully degenerate or random position. Completely random oligonucleotides can be made by including all four nucleotide precursors in every round of synthesis. Degenerate oligonucleotides can also be made having different proportions of different nucleotides. Such

oligonucleotides can be made, for example, by using different nucleotide precursors, in the desired proportions, in the reaction.

Many of the oligonucleotides described herein are designed to be complementary to certain portions of other oligonucleotides or nucleic acids such that stable hybrids can be formed between them. The stability of these hybrids can be calculated using known methods such as those described in Lesnick and Freier, *Biochemistry* **34**:10807-10815 (1995), McGraw *et al.*, *Biotechniques* **8**:674-678 (1990), and Rychlik *et al.*, *Nucleic Acids Res.* **18**:6409-6412 (1990).

Hexamer oligonucleotides were synthesized on a Perseptive Biosystems 8909 Expedite Nucleic Acid Synthesis system using standard β -cyanoethyl phosphoramidite coupling chemistry on mixed dA+dC+dG+dT synthesis columns (Glen Research, Sterling, VA). The four phosphoramidites were mixed in equal proportions to randomize the bases at each position in the oligonucleotide. Oxidation of the newly formed phosphites were carried out using the sulfurizing reagent 3H-1,2-benzothiole-3-one-1,1-dioxide (Glen Research) instead of the standard oxidizing reagent after the first and second phosphoramidite addition steps. The thio-phosphitylated oligonucleotides were deprotected using 30% ammonium hydroxide (3.0 ml) in water at 55°C for 16 hours, concentrated in an OP 120 Savant Oligo Prep deprotection unit for 2 hours, and desalted with PD10 Sephadex columns using the protocol provided by the manufacturer.

M. DNA polymerases

DNA polymerases useful in multiple displacement amplification must be capable of displacing, either alone or in combination with a compatible strand displacement factor, a hybridized strand encountered during replication. Such polymerases are referred to herein as strand displacement DNA polymerases. It is preferred that a strand displacement DNA polymerase lack a 5' to 3' exonuclease activity. Strand displacement is necessary to result in synthesis of multiple copies of a target sequence. A 5' to 3' exonuclease activity, if present, might result in the destruction of a synthesized strand. It is also preferred that DNA polymerases for use in the disclosed method are highly processive. The suitability of a DNA polymerase for use in the disclosed method can be readily determined by assessing its ability to carry out strand displacement replication. Preferred strand displacement DNA

polymerases are bacteriophage ϕ 29 DNA polymerase (U.S. Patent Nos. 5,198,543 and 5,001,050 to Blanco *et al.*), Bst large fragment DNA polymerase (Exo(-) Bst; Aliotta *et al.*, *Genet. Anal. (Netherlands)* **12**:185-195 (1996)) and exo(-)Bca DNA polymerase (Walker and Linn, *Clinical Chemistry* **42**:1604-1608 (1996)). Other useful

5 polymerases include phage M2 DNA polymerase (Matsumoto *et al.*, *Gene* **84**:247 (1989)), phage ϕ PRD1 DNA polymerase (Jung *et al.*, *Proc. Natl. Acad. Sci. USA* **84**:8287 (1987)), exo(-)VENT® DNA polymerase (Kong *et al.*, *J. Biol. Chem.* **268**:1965-1975 (1993)), Klenow fragment of DNA polymerase I (Jacobsen *et al.*, *Eur. J. Biochem.* **45**:623-627 (1974)), T5 DNA polymerase (Chatterjee *et al.*, *Gene* **97**:13-19

10 (1991)), Sequenase (U.S. Biochemicals), PRD1 DNA polymerase (Zhu and Ito, *Biochim. Biophys. Acta.* **1219**:267-276 (1994)), and T4 DNA polymerase holoenzyme (Kaboord and Benkovic, *Curr. Biol.* **5**:149-157 (1995)). ϕ 29 DNA polymerase is most preferred.

Strand displacement can be facilitated through the use of a strand displacement

15 factor, such as helicase. It is considered that any DNA polymerase that can perform strand displacement replication in the presence of a strand displacement factor is suitable for use in the disclosed method, even if the DNA polymerase does not perform strand displacement replication in the absence of such a factor. Strand displacement factors useful in strand displacement replication include BMRF1 polymerase accessory

20 subunit (Tsurumi *et al.*, *J. Virology* **67**(12):7648-7653 (1993)), adenovirus DNA-binding protein (Zijderveld and van der Vliet, *J. Virology* **68**(2):1158-1164 (1994)), herpes simplex viral protein ICP8 (Boehmer and Lehman, *J. Virology* **67**(2):711-715 (1993); Skalter and Lehman, *Proc. Natl. Acad. Sci. USA* **91**(22):10665-10669 (1994)); single-stranded DNA binding proteins (SSB; Rigler and Romano, *J. Biol. Chem.* **270**:8910-8919 (1995)); phage T4 gene 32 protein (Villemain and Giedroc, *Biochemistry* **35**:14395-14404 (1996); and calf thymus helicase (Siegel *et al.*, *J. Biol. Chem.* **267**:13629-13635 (1992)).

25

The ability of a polymerase to carry out strand displacement replication can be determined by using the polymerase in a strand displacement replication assay such as

30 those described in Examples 1 and 5. The assay in the examples can be modified as appropriate. For example, a helicase can be used instead of SSB. Such assays should be performed at a temperature suitable for optimal activity for the enzyme being used,

for example, 32°C for ϕ 29 DNA polymerase, from 46°C to 64°C for exo(-) Bst DNA polymerase, or from about 60°C to 70°C for an enzyme from a hyperthermophilic organism. For assays from 60°C to 70°C, primer length may be increased to provide a melting temperature appropriate for the assay temperature. Another useful assay for
5 selecting a polymerase is the primer-block assay described in Kong *et al.*, *J. Biol. Chem.* **268**:1965-1975 (1993). The assay consists of a primer extension assay using an M13 ssDNA template in the presence or absence of an oligonucleotide that is hybridized upstream of the extending primer to block its progress. Enzymes able to displace the blocking primer in this assay are expected to be useful for the disclosed
10 method.

N. Kits

The materials described above can be packaged together in any suitable combination as a kit useful for performing the disclosed method. Kit components in a given kit can be designed and adapted for use together in the disclosed method. For
15 example, disclosed are kits for amplifying genomic DNA, the kit comprising a lysis solution, a stabilization solution, a set of primers, and a DNA polymerase. The components of such a kit are described elsewhere herein. In some forms of the kit, the lysis solution can comprise potassium hydroxide, for example, 400 mM KOH. Some useful forms of lysis solution can comprise 400 mM KOH, 100 mM dithiothreitol, and
20 10 mM EDTA. In some forms of the kit, the stabilization solution can comprise Tris-HCl at pH 4.1. Some useful forms of stabilization solution can comprise 800 mM Tris-HCl, pH 4.1. In some forms of the kit, the set of primers can comprise random hexamer primers. In some forms of the kit, the DNA polymerase can be ϕ 29 DNA polymerase. In some forms of the kit, the kit can further comprise deoxynucleotide
25 triphosphates. In some forms of the kit, the kit can further comprise one or more detection probes. Detection probes are described elsewhere herein. In some forms of the kit, the detection probes can each comprise a complementary portion, where the complementary portion is complementary to a nucleic acid sequence of interest. In some forms of the kit, the kit can further comprise denaturing solution. In some forms
30 of the kit, the kit can further comprise reaction mix.

Some useful kits can comprise a lysis solution, a stabilization solution, a set of primers, a ϕ 29 DNA polymerase, 1M dithiothreitol, 1X Phosphate-Buffered Saline, pH

7.5, and control DNA template; where the lysis solution comprises 400 mM KOH and 10 mM EDTA, the stabilization solution comprises 800 mM Tris-HCl, pH 4, and the set of primers comprises a reaction mix; where the reaction mix comprises 150 mM Tris-HCl, 200 mM KCl, 40 mM MgCl₂, 20 mM (NH₄)₂SO₄, 4 mM deoxynucleotide triphosphates, and 0.2 mM random hexamer primers.

Any of the components that can be present in a kit that can be used together can be combined in a single component of the kit. Thus, a reaction mix can include, for example, buffers, deoxynucleotide triphosphates and primers. Similarly, components and solutions can be divided into constituent parts or sub-solutions. The kits can be used for any purpose, generally for nucleic acid amplification. In some forms, the kit can be designed to detect nucleic acid sequences of interest in a genome or other nucleic acid sample. In some forms, the kit can be designed to assess a disease, condition or predisposition of an individual based on a nucleic acid sequences of interest.

O. Mixtures

Disclosed are mixtures formed by performing, or formed during the course of performing, any form of the disclosed method. For example, disclosed are mixtures comprising, for example, cells and lysis solution; cell lysate and stabilization solution; stabilized cell lysate and one or more primers; stabilized cell lysate and DNA polymerase; stabilized cell lysate, one or more primers, and DNA polymerase; stabilized cell lysate and replicated strands; stabilized cell lysate, one or more primers, and replicated strands; stabilized cell lysate, DNA polymerase, and replicated strands; stabilized cell lysate, one or more primers, DNA polymerase, and replicated strands; stabilized cell lysate and one or more detection probes; stabilized cell lysate, one or more primers, one or more detection probes, and replicated strands; stabilized cell lysate, DNA polymerase, one or more detection probes, and replicated strands; and stabilized cell lysate, one or more primers, DNA polymerase, one or more detection probes, and replicated strands.

Whenever the method involves mixing or bringing into contact, for example, compositions or components or reagents, performing the method creates a number of different mixtures. For example, if the method includes three mixing steps, after each one of these steps a unique mixture is formed if the steps are performed sequentially.

In addition, a mixture is formed at the completion of all of the steps regardless of how the steps were performed. The present disclosure contemplates these mixtures, obtained by the performance of the disclosed method as well as mixtures containing any disclosed reagent, composition, or component, for example, disclosed herein.

5

Uses

The disclosed method and compositions are applicable to numerous areas including, but not limited to, analysis of nucleic acids present in cells (for example, analysis of genomic DNA in cells), disease detection, mutation detection, gene discovery, gene mapping (molecular haplotyping), and agricultural research.

10

Particularly useful is whole genome amplification. Other uses include, for example, detection of nucleic acids in cells and on genomic DNA arrays; molecular haplotyping; mutation detection; detection of inherited diseases such as cystic fibrosis, muscular dystrophy, diabetes, hemophilia, sickle cell anemia; assessment of predisposition for cancers such as prostate cancer, breast cancer, lung cancer, colon cancer, ovarian cancer, testicular cancer, pancreatic cancer.

15

Method

The disclosed method is based on strand displacement replication of the nucleic acid sequences by multiple primers. The method can be used to amplify one or more specific sequences (multiple strand displacement amplification), an entire genome or other DNA of high complexity (whole genome strand displacement amplification), or concatenated DNA (multiple strand displacement amplification of concatenated DNA). The disclosed method generally involves hybridization of primers to a target nucleic acid sequence and replication of the target sequence primed by the hybridized primers such that replication of the target sequence results in replicated strands complementary to the target sequence. During replication, the growing replicated strands displace other replicated strands from the target sequence (or from another replicated strand) via strand displacement replication. As used herein, a replicated strand is a nucleic acid strand resulting from elongation of a primer hybridized to a target sequence or to another replicated strand. Strand displacement replication refers to DNA replication where a growing end of a replicated strand encounters and displaces another strand from the template strand (or from another replicated strand). Displacement of replicated strands by other replicated strands is a hallmark of the disclosed method

20

25

30

which allows multiple copies of a target sequence to be made in a single, isothermal reaction.

Nucleic acids for amplification are often obtained from cellular samples. This generally requires disruption of the cell (to make the nucleic acid accessible) and
5 purification of the nucleic acids prior to amplification. It also generally requires the inactivation of protein factors such as nucleases that could degrade the DNA, or of factors such as histones that could bind to DNA strands and impede their use as a template for DNA synthesis by a polymerase. There are a variety of techniques used to break open cells, such as sonication, enzymatic digestion of cell walls, heating, and
10 exposure to lytic conditions. Lytic conditions typically involve use of non-physiological pH and/or solvents. Many lytic techniques can result in damage to nucleic acids in cells, including, for example, breakage of genomic DNA. In particular, use of heating to lyse cells can damage genomic DNA and reduce the amount and quality of amplification products of genomic DNA. It has been discovered that alkaline
15 lysis can cause less damage to genomic DNA and can thus result in higher quality amplification results. Alkaline lysis also inactivates protein factors such as nucleases, histones, or other factors that could impede the amplification of DNA within the sample. In addition, it is a useful property of alkaline lysis that reducing the pH does not reactivate the protein factors, but that such protein factors remain inactivated when
20 the pH of the solution is brought back within a neutral range.

In some forms of the disclosed method, a genomic sample is prepared by exposing cells to alkaline conditions, thereby lysing the cells and resulting in a cell lysate; reducing the pH of the cell lysate to make the pH of the cell lysate compatible with DNA replication; and incubating the cell lysate under conditions that promote
25 replication of the genome of the cells by multiple displacement amplification. Alkaline conditions are conditions where the pH is greater than 9.0. Particularly useful alkaline conditions for the disclosed method are conditions where the pH is greater than 10.0. The alkaline conditions can be, for example, those that cause a substantial number of cells to lyse, those that cause a significant number of cells to lyse, or those that cause a
30 sufficient number of cells to lyse. The number of lysed cells can be considered sufficient if the genome can be sufficiently amplified in the disclosed method. The amplification is sufficient if enough amplification product is produced to permit some

use of the amplification product, such as detection of sequences or other analysis. The reduction in pH is generally into the neutral range of pH 9.0 to pH 6.0.

The cells can be exposed to alkaline conditions by mixing the cells with a lysis solution. The amount of lysis solution mixed with the cells can be that amount that
5 causes a substantial number of cells to lyse or those that cause a sufficient number of cells to lyse. Generally, this volume will be a function of the pH of the cell/lysis solution mixture. Thus, the amount of lysis solution to mix with cells can be determined generally from the volume of the cells and the alkaline concentration of the lysis buffer. For example, a smaller volume of a lysis solution with a stronger base and/or higher
10 concentration of base would be needed to create sufficient alkaline conditions than the volume needed of a lysis solution with a weaker base and/or lower concentration of base. The lysis solution can be formulated such that the cells are mixed with an equal volume of the lysis solution (to produce the desired alkaline conditions).

In some embodiments, the lysis solution can comprise a base, such as an
15 aqueous base. Useful bases include potassium hydroxide, sodium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine, methylamine, N-
20 methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, and DBU (1,8-diazobicyclo[5,4,0]undec-7-ene). Useful formulations of lysis solution include lysis solution comprising 400 mM KOH, lysis solution comprising 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA, and lysis
25 solution consisting of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

In some embodiments, the lysis solution can comprise a plurality of basic agents. As used herein, a basic agent is a compound, composition or solution that results in alkaline conditions. In some embodiments, the lysis solution can comprise a buffer. Useful buffers include phosphate buffers, "Good" buffers (such as BES,
30 BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, and TRICINE), sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, and Bis-tris propane. The lysis solution can comprise a

plurality of buffering agents. As used herein, a buffering agent is a compound, composition or solution that acts as a buffer. An alkaline buffering agent is a buffering agent that results in alkaline conditions. In some embodiments, the lysis solution can comprise a combination of one or more bases, basic agents, buffers and buffering
5 agents.

The pH of the cell lysate can be reduced to form a stabilized cell lysate. A stabilized cell lysate is a cell lysate the pH of which is in the neutral range (from about pH 6.0 to about pH 9.0). Useful stabilized cell lysates have a pH that allows replication of nucleic acids in the cell lysate. For example, the pH of the stabilized cell lysate is
10 usefully at a pH at which the DNA polymerase can function. The pH of the cell lysate can be reduced by mixing the cell lysate with a stabilization solution. The stabilization solution comprises a solution that can reduce the pH of a cell lysate exposed to alkaline conditions as described elsewhere herein.

The amount of stabilization solution mixed with the cell lysate can be that
15 amount that causes a reduction in pH to the neutral range (or other desired pH value). Generally, this volume will be a function of the pH of the cell lysate/stabilization solution mixture. Thus, the amount of stabilization solution to mix with the cell lysate can be determined generally from the volume of the cell lysate, its pH and buffering capacity, and the acidic concentration of the stabilization buffer. For example, a
20 smaller volume of a stabilization solution with a stronger acid and/or higher concentration of acid would be needed to reduce the pH sufficiently than the volume needed of a stabilization solution with a weaker acid and/or lower concentration of acid. The stabilization solution can be formulated such that the cell lysate is mixed with an equal volume of the stabilization solution (to produce the desired pH).

25 In some embodiments, the stabilization solution can comprise an acid. Useful acids include hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, and any carboxylic acid (ethanoic, propanoic,
30 butanoic, etc., including both linear or branched chain carboxylic acids). In some embodiments, the stabilization solution can comprise a buffer. Useful buffers include Tris-HCl, HEPES, "Good" buffers (such as BES, BICINE, CAPS, EPPS, HEPES,

MES, MOPS, PIPES, TAPS, TES, and TRICINE), sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, and Bis-tris propane. Useful formulations of stabilization solutions include stabilization solution comprising 800 mM Tris-HCl; stabilization solution comprising 800 mM Tris-HCl at
5 pH 4.1; and stabilization solution consisting of 800 mM Tris-HCl, pH 4.1.

In some embodiments, the stabilization solution can comprise a plurality of acidic agents. As used herein, an acidic agent is a compound, composition or solution that forms an acid in solution. In some embodiments, the stabilization solution can comprise a plurality of buffering agents. An acidic buffering agent is a buffering agent
10 that forms an acid in solution. In some embodiments, the stabilization solution can comprise a combination of one or more acids, acidic agents, buffers and buffering agents.

In some embodiments, the pH of the cell lysate can be reduced to about pH 9.0 or below, to about pH 8.5 or below, to about pH 8.0 or below, to about pH 7.5 or
15 below, to about pH 7.2 or below, or to about pH 7.0 or below. In some embodiments, the pH of the cell lysate can be reduced to the range of about pH 9.0 to about pH 6.0, to the range of about pH 9.0 to about pH 6.5, to the range of about pH 9.0 to about pH 6.8, to the range of about pH 9.0 to about pH 7.0, to the range of about pH 9.0 to about pH 7.2, to the range of about pH 9.0 to about pH 7.5, to the range of about pH 9.0 to about
20 pH 8.0, to the range of about pH 9.0 to about pH 8.5, to the range of about pH 8.5 to about pH 6.0, to the range of about pH 8.5 to about pH 6.5, to the range of about pH 8.5 to about pH 6.8, to the range of about pH 8.5 to about pH 7.0, to the range of about pH 8.5 to about pH 7.2, to the range of about pH 8.5 to about pH 7.5, to the range of about pH 8.5 to about pH 8.0, to the range of about pH 8.0 to about pH 6.0, to the range of
25 about pH 8.0 to about pH 6.5, to the range of about pH 8.0 to about pH 6.8, to the range of about pH 8.0 to about pH 7.0, to the range of about pH 8.0 to about pH 7.2, to the range of about pH 8.0 to about pH 7.5, to the range of about pH 7.5 to about pH 6.0, to the range of about pH 7.5 to about pH 6.5, to the range of about pH 7.5 to about pH 6.8, to the range of about pH 7.5 to about pH 7.0, to the range of about pH 7.5 to about pH
30 7.2, to the range of about pH 7.2 to about pH 6.0, to the range of about pH 7.2 to about pH 6.5, to the range of about pH 7.2 to about pH 6.8, to the range of about pH 7.2 to about pH 7.0, to the range of about pH 7.0 to about pH 6.0, to the range of about pH 7.0

to about pH 6.5, to the range of about pH 7.0 to about pH 6.8, to the range of about pH 6.8 to about pH 6.0, to the range of about pH 6.8 to about pH 6.5, or to the range of about pH 6.5 to about pH 6.0. In some embodiments, the pH of the cell lysate can be reduced to any range having any combination of endpoints from about pH 6.0 to about pH 9.0 All such endpoints and ranges are specifically and separately contemplated.

In some embodiments, the cells are not lysed by heat. Those of skill in the art will understand that different cells under different conditions will be lysed at different temperatures and so can determine temperatures and times at which the cells will not be lysed by heat. In general, the cells are not subjected to heating above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions used. As used herein, substantial cell lysis refers to lysis of 90% or greater of the cells exposed to the alkaline conditions. Significant cell lysis refers to lysis of 50% or more of the cells exposed to the alkaline conditions. Sufficient cell lysis refers to lysis of enough of the cells exposed to the alkaline conditions to allow synthesis of a detectable amount of amplification products by multiple strand displacement amplification. In general, the alkaline conditions used in the disclosed method need only cause sufficient cell lysis. It should be understood that alkaline conditions that could cause significant or substantial cell lysis need not result in significant or substantial cell lysis when the method is performed.

In some embodiments, the cells are not subjected to heating substantially or significantly above the temperature at which the cells grow. As used herein, the temperature at which the cells grow refers to the standard temperature, or highest of different standard temperatures, at which cells of the type involved are cultured. In the case of animal cells, the temperature at which the cells grow refers to the body temperature of the animal. In other embodiments, the cells are not subjected to heating substantially or significantly above the temperature of the amplification reaction (where the genome is replicated).

In some embodiments, the cell lysate is not subjected to purification prior to the amplification reaction. In the context of the disclosed method, purification generally refers to the separation of nucleic acids from other material in the cell lysate. It has been discovered that multiple displacement amplification can be performed on unpurified and partially purified samples. It is commonly thought that amplification

reactions cannot be efficiently performed using unpurified nucleic acid. In particular, PCR is very sensitive to contaminants.

Forms of purification include centrifugation, extraction, chromatography, precipitation, filtration, and dialysis. Partially purified cell lysate includes cell lysates subjected to centrifugation, extraction, chromatography, precipitation, filtration, and dialysis. Partially purified cell lysate generally does not include cell lysates subjected to nucleic acid precipitation or dialysis. As used herein, separation of nucleic acid from other material refers to physical separation such that the nucleic acid to be amplified is in a different container or container from the material. Purification does not require separation of all nucleic acid from all other materials. Rather, what is required is separation of some nucleic acid from some other material. As used herein in the context of nucleic acids to be amplified, purification refers to separation of nucleic acid from other material. In the context of cell lysates, purification refers to separation of nucleic acid from other material in the cell lysate. As used herein, partial purification refers to separation of nucleic acid from some, but not all, of other material with which the nucleic acid is mixed. In the context of cell lysates, partial purification refers to separation of nucleic acid from some, but not all, of the other material in the cell lysate.

Unless the context clearly indicates otherwise, reference herein to a lack of purification, lack of one or more types of purification or separation operations or techniques, or exclusion of purification or one or more types of purification or separation operations or techniques does not encompass the exposure of cells to alkaline conditions (or the results thereof) the reduction of pH of a cell lysate (or the results thereof). That is, to the extent that the alkaline conditions and pH reduction of the disclosed method produce an effect that could be considered "purification" or "separation," such effects are excluded from the definition of purification and separation when those terms are used in the context of processing and manipulation of cell lysates and stabilized cell lysates (unless the context clearly indicates otherwise).

As used herein, substantial purification refers to separation of nucleic acid from at least a substantial portion of other material with which the nucleic acid is mixed. In the context of cell lysates, substantial purification refers to separation of nucleic acid from at least a substantial portion of the other material in the cell lysate. A substantial portion refers to 90% of the other material involved. Specific levels of purification can

be referred to as a percent purification (such as 95% purification and 70% purification). A percent purification refers to purification that results in separation from nucleic acid of at least the designated percent of other material with which the nucleic acid is mixed.

Denaturation of nucleic acid molecules to be amplified is common in
5 amplification techniques. This is especially true when amplifying genomic DNA. In particular, PCR uses multiple denaturation cycles. Denaturation is generally used to make nucleic acid strands accessible to primers. It was discovered that the target nucleic acids, genomic DNA, for example, need not be denatured for efficient multiple displacement amplification. It was also discovered that elimination of a denaturation
10 step and denaturation conditions has additional advantages such as reducing sequence bias in the amplified products. In some embodiments, the nucleic acids in the cell lysate are not denatured by heating. In some embodiments, the cell lysate is not subjected to heating substantially or significantly above the temperature at which the cells grow. In other embodiments, the cell lysate is not subjected to heating
15 substantially or significantly above the temperature of the amplification reaction (where the genome is replicated). The disclosed multiple displacement amplification reaction is generally conducted at a substantially constant temperature (that is, the amplification reaction is substantially isothermic), and this temperature is generally below the temperature at which the nucleic acids would be notably denatured. As used herein,
20 notable denaturation refers to denaturation of 10% or greater of the base pairs.

In preferred forms of the disclosed method, the nucleic acid sample or template nucleic acid is not subjected to denaturing conditions and/or no denaturation step is used. In some forms of the disclosed method, the nucleic acid sample or template nucleic acid is not subjected to heat denaturing conditions and/or no heat denaturation
25 step is used. It should be understood that while sample preparation (for example, cell lysis and processing of cell extracts) may involve conditions that might be considered denaturing (for example, treatment with alkali), the denaturation conditions or step eliminated in some forms of the disclosed method refers to denaturation steps or conditions intended and used to make nucleic acid strands accessible to primers. Such
30 denaturation is commonly a heat denaturation, but can also be other forms of denaturation such as chemical denaturation. It should be understood that in the disclosed method where the nucleic acid sample or template nucleic acid is not

subjected to denaturing conditions, the template strands are accessible to the primers (since amplification occurs). However, the template stands are not made accessible via general denaturation of the sample or template nucleic acids.

The efficiency of a DNA amplification procedure may be described for individual loci as the percent representation, where the percent representation is 100% for a locus in genomic DNA as purified from cells. For 10,000-fold amplification, the average representation frequency was 141% for 8 loci in DNA amplified without heat denaturation of the template, and 37% for the 8 loci in DNA amplified with heat denaturation of the template. The omission of a heat denaturation step results in a 3.8-fold increase in the representation frequency for amplified loci. Amplification bias may be calculated between two samples of amplified DNA or between a sample of amplified DNA and the template DNA it was amplified from. The bias is the ratio between the values for percent representation for a particular locus. The maximum bias is the ratio of the most highly represented locus to the least represented locus. For 10,000-fold amplification, the maximum amplification bias was 2.8 for DNA amplified without heat denaturation of the template, and 50.7 for DNA amplified with heat denaturation of the template. The omission of a heat denaturation step results in an 18-fold decrease in the maximum bias for amplified loci.

In another form of the method, the primers can be hexamer primers. It was discovered that such short, 6 nucleotide primers can still prime multiple strand displacement replication efficiently. Such short primers are easier to produce as a complete set of primers of random sequence (random primers) than longer primers at least because there are fewer to make. In another form of the method, the primers can each contain at least one modified nucleotide such that the primers are nuclease resistant. In another form of the method, the primers can each contain at least one modified nucleotide such that the melting temperature of the primer is altered relative to a primer of the same sequence without the modified nucleotide(s). In another form of the method, the DNA polymerase can be ϕ 29 DNA polymerase. It was discovered that ϕ 29 DNA polymerase produces greater amplification in multiple displacement amplification. The combination of two or more of the above features also yields improved results in multiple displacement amplification. In a preferred embodiment, for example, the target sample is not subjected to denaturing conditions, the primers are

hexamer primers and contain modified nucleotides such that the primers are nuclease resistant, and the DNA polymerase is ϕ 29 DNA polymerase. The above features are especially useful in whole genome strand displacement amplification (WGSDA).

5 In another form of the disclosed method, the method includes labeling of the replicated strands (that is, the strands produced in multiple displacement amplification) using terminal deoxynucleotidyl transferase. The replicated strands can be labeled by, for example, the addition of modified nucleotides, such as biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates, to the 3' ends of the replicated strands.

10 Some forms of the disclosed method provide amplified DNA of higher quality relative to previous methods due to the lack of a heat denaturation treatment of the DNA that is the target for amplification. Thus, the template DNA does not undergo the strand breakage events caused by heat treatment and the amplification that is accomplished by a single DNA polymerase extends farther along template strands of
15 increased length.

In one form of the disclosed method, a small amount of purified double-strand human genomic DNA (1 ng, for example) can be mixed with exonuclease-resistant random hexamer primers and ϕ 29 DNA polymerase under conditions that favor DNA synthesis. For example, the mixture can simply be incubated at 30°C and multiple
20 displacement amplification will take place. Thus, any single-stranded or duplex DNA may be used, without any additional treatment, making the disclosed method a simple, one-step procedure. Since so little DNA template is required, a major advantage of the disclosed method is that DNA template may be taken from preparations that contain levels of contaminants that would inhibit other DNA amplification procedures such as
25 PCR. For MDA the sample may be diluted so that the contaminants fall below the concentration at which they would interfere with the reaction. The disclosed method can be performed (and the above advantages achieved) using any type of sample, including, for example, bodily fluids such as urine, semen, lymphatic fluid, cerebrospinal fluid, and amniotic fluid.

30 The need for only small amounts of DNA template in the disclosed method means that the method is useful for DNA amplification from very small samples. In particular, the disclosed method may be used to amplify DNA from a single cell. The

ability to obtain analyzable amounts of nucleic acid from a single cell (or similarly small sample) has many applications in preparative, analytical, and diagnostic procedures such as prenatal diagnostics. Other examples of biological samples containing only small amounts of DNA for which amplification by the disclosed method would be useful are material excised from tumors or other archived medical samples, needle aspiration biopsies, clinical samples arising from infections, such as nosocomial infections, forensic samples, or museum specimens of extinct species.

More broadly, the disclosed method is useful for applications in which the amounts of DNA needed are greater than the supply. For example, procedures that analyze DNA by chip hybridization techniques are limited by the amounts of DNA that can be purified from typically sized blood samples. As a result many chip hybridization procedures utilize PCR to generate a sufficient supply of material for the high-throughput procedures. The disclosed method presents a useful technique for the generation of plentiful amounts of amplified DNA that faithfully reproduces the locus representation frequencies of the starting material.

Whole genome amplification by MDA can be carried out directly from blood or cells bypassing the need to isolate pure DNA. For example, blood or other cells can be lysed by dilution with an equal volume of 2X Alkaline Lysis Buffer (400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA), an example of a lysis solution, and incubated 10 minutes on ice. The lysed cells can be neutralized with the same volume of Neutralization Buffer (800 mM Tris-HCl, pH 4.1), an example of a stabilization solution. Preparations of lysed blood or cells (for example, 1 ml) can be used directly as template in MDA reactions (for example, 100 ml). If desired, prior to lysis, blood can be diluted 3-fold in phosphate buffered saline (PBS) and tissue culture cells can be diluted to 30,000 cells/ml in PBS.

A specific embodiment of the disclosed method is described in Example 1, wherein whole genome amplification is performed by MDA without heat treatment of the human template DNA. As shown in the example, the disclosed method produces a DNA amplification product with improved performance in genetic assays compared to amplification performed with heat treatment of the template DNA. The longer DNA products produced without heat treatment of the template yield larger DNA fragments in Southern blotting and genetic analysis using RFLP.

The breakage of DNA strands by heat treatment is demonstrated directly in Example 2, while the decreased rate and yield of DNA amplification from heat-treated DNA is depicted in Example 3. The decrease in DNA product strand length resulting from heat treatment of the DNA template is demonstrated in Example 4.

5 A specific form of the disclosed method is described in Example 5, wherein purified human genomic DNA is amplified by MDA without heat treatment of the template. As shown in the example, the disclosed method produces for a DNA amplification product with no loss of locus representation when used as a substrate in quantitative PCR assays compared to DNA amplified with heat treatment of the
10 template.

Another specific form of the disclosed method is described in Example 6, wherein purified human genomic DNA is amplified by MDA without heat treatment of the template. As shown in the example, the disclosed method produces a DNA amplification product with a low amplification bias, with the variation in representation
15 among eight different loci varying by less than 3.0. In contrast, the amplification bias of DNA products amplified by two PCR-based amplification methods, PEP and DOP-PCR, varies between two and six orders of magnitude.

Another specific form of the disclosed method is described in Example 7, wherein the amplification of c-jun sequences using specific, nested primers from a
20 human genomic DNA template is enhanced by omission of a DNA template heat denaturation step.

Another specific form of the disclosed method is described in Example 8, wherein human genomic DNA is amplified in the absence of a heat treatment step directly from whole blood or from tissue culture cells with the same efficiency as from
25 purified DNA. The DNA amplified directly from blood or cells has substantially the same locus representation values as DNA amplified from purified human DNA template. This represents an advantage over other amplification procedures such as PCR, since components such as heme in whole blood inhibit PCR and necessitate a purification step before DNA from blood can be used as a PCR template.

30 Another specific form of the disclosed method is described in Example 9, wherein purified human genomic DNA is amplified by MDA without heat treatment of the template in the presence of 70% AA-dUTP / 30% dTTP. As shown in the example,

the disclosed method provides for a DNA amplification product with the same low amplification bias as for DNA amplified in the presence of 100% dTTP.

Also disclosed is a method for amplifying and repairing damaged DNA. This method is useful, for example, for amplifying degraded genomic DNA. The method involves substantially denaturing a damaged DNA sample (generally via exposure to heat and alkaline conditions), removal or reduction of the denaturing conditions (such as by reduction of the pH and temperature of the denatured DNA sample), and replicating the DNA. The damaged DNA is repaired during replication and the average length of DNA fragments is increased. For example, the average length of DNA fragments can be increase from, for example, 2 kb in the damaged DNA sample to, for example, 10 kb or greater for the replicated DNA. The amplified and repaired DNA is in better condition for analysis and testing than the damaged DNA sample. For example, this technique can provide consistent improvements in allele representation from damaged DNA samples. This repair method can result in an overall improvement in amplification of damaged DNA by increasing the average length of the product, increasing the quality of the amplification products by 3-fold (by, for example, increasing the marker representation in the sample), and improving the genotyping of amplified products by lowering the frequency of allelic dropout; all compared to the results when amplifying damaged DNA by other methods. The replication can be multiple displacement amplification. Denaturation of the DNA sample generally is carried out such that the DNA is not further damaged. This method can generally be combined or used with any of the disclosed amplification methods. Another form of this method can involve substantially denaturing a damaged DNA sample (generally via exposure to heat and alkaline conditions), reduction of the pH of the denatured DNA sample, mixing the denatured DNA sample with an undenatured DNA sample from the same source such that the ends of DNA in the undenatured DNA sample is transiently denatured, slowly cooling the mixture of DNA samples to allow the transiently denatured ends to anneal to the denatured DNA, and replicating the annealed DNA.

A. Whole Genome Strand Displacement Amplification

In one form of the method, referred to as whole genome strand displacement amplification (WGSDA), a random or partially random set of primers is used to

randomly prime a sample of genomic nucleic acid (or another sample of nucleic acid of high complexity). By choosing a sufficiently large set of primers of random or mostly random sequence, the primers in the set will be collectively, and randomly, complementary to nucleic acid sequences distributed throughout nucleic acid in the sample. Amplification proceeds by replication with a processive polymerase initiated at each primer and continuing until spontaneous termination. A key feature of this method is the displacement of intervening primers during replication by the polymerase. In this way, multiple overlapping copies of the entire genome can be synthesized in a short time.

Whole genome strand displacement amplification can be performed by (a) mixing a set of random or partially random primers with a genomic sample (or other nucleic acid sample of high complexity), to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the genomic DNA in the primer-target sample mixture, and (b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the genomic DNA. Strand displacement replication is preferably accomplished by using a strand displacing DNA polymerase or a DNA polymerase in combination with a compatible strand displacement factor.

The method has advantages over the polymerase chain reaction since it can be carried out under isothermal conditions. Other advantages of whole genome strand displacement amplification include a higher level of amplification than whole genome PCR, amplification is less sequence-dependent than PCR, a lack of re-annealing artifacts or gene shuffling artifacts as can occur with PCR (since there are no cycles of denaturation and re-annealing), and a lower amplification bias than PCR-based genome amplification (bias of 3-fold for WGSDA versus 20- to 60-fold for PCR-based genome amplification).

Following amplification, the amplified sequences can be used for any purpose, such as uses known and established for PCR amplified sequences. For example, amplified sequences can be detected using any of the conventional detection systems for nucleic acids such as detection of fluorescent labels, enzyme-linked detection

systems, antibody-mediated label detection, and detection of radioactive labels. A key feature of the disclosed method is that amplification takes place not in cycles, but in a continuous, isothermal replication. This makes amplification less complicated and much more consistent in output. Strand displacement allows rapid generation of multiple copies of a nucleic acid sequence or sample in a single, continuous, isothermal reaction.

It is preferred that the set of primers used for WGSDA be used at concentrations that allow the primers to hybridize at desired intervals within the nucleic acid sample. For example, by using a set of primers at a concentration that allows them to hybridize, on average, every 4000 to 8000 bases, DNA replication initiated at these sites will extend to and displace strands being replicated from adjacent sites. It should be noted that the primers are not expected to hybridize to the target sequence at regular intervals. Rather, the average interval will be a general function of primer concentration. Primers for WGSDA can also be formed from RNA present in the sample. By degrading endogenous RNA with RNase to generate a pool of random oligomers, the random oligomers can then be used by the polymerase for amplification of the DNA. This eliminates any need to add primers to the reaction. Alternatively, DNase digestion of biological samples can generate a pool of DNA oligo primers for RNA dependent DNA amplification.

As in multiple strand displacement amplification, displacement of an adjacent strand makes it available for hybridization to another primer and subsequent initiation of another round of replication. The interval at which primers in the set of primers hybridize to the target sequence determines the level of amplification. For example, if the average interval is short, adjacent strands will be displaced quickly and frequently. If the average interval is long, adjacent strands will be displaced only after long runs of replication.

In the disclosed method, the DNA polymerase catalyzes primer extension and strand displacement in a processive strand displacement polymerization reaction that proceeds as long as desired. Preferred strand displacing DNA polymerases are bacteriophage ϕ 29 DNA polymerase (U.S. Patent Nos. 5,198,543 and 5,001,050 to Blanco *et al.*), large fragment Bst DNA polymerase (Exo(-) Bst), exo(-)Bca DNA polymerase, and Sequenase. During strand displacement replication one may

additionally include radioactive, or modified nucleotides such as bromodeoxyuridine triphosphate, in order to label the DNA generated in the reaction. Alternatively, one may include suitable precursors that provide a binding moiety such as biotinylated nucleotides (Langer *et al.*, *Proc. Natl. Acad. Sci. USA* **78**:6633 (1981)).

5 Genome amplification using PCR, and uses for the amplified DNA, is described in Zhang *et al.*, *Proc. Natl. Acad. Sci. USA* **89**:5847-5851 (1992), Telenius *et al.*, *Genomics* **13**:718-725 (1992), Cheung *et al.*, *Proc. Natl. Acad. Sci. USA* **93**:14676-14679 (1996), and Kukasjaarvi *et al.*, *Genes, Chromosomes and Cancer* **18**:94-101 (1997). The uses of the amplified DNA described in these publications are also
10 generally applicable to DNA amplified using the disclosed methods. Whole Genome Strand Displacement Amplification, unlike PCR-based whole genome amplification, is suitable for haplotype analysis since WGSDA yields longer fragments than PCR-based whole genome amplification. PCR-based whole genome amplification is also less
15 suitable for haplotype analysis since each cycle in PCR creates an opportunity for priming events that result in the association of distant sequences (in the genome) to be put together in the same fragment.

B. Multiple Strand Displacement Amplification

 In one preferred form of the method, referred to as multiple strand displacement amplification (MSDA), two sets of primers are used, a right set and a left set. Primers
20 in the right set of primers each have a portion complementary to nucleotide sequences flanking one side of a target nucleotide sequence and primers in the left set of primers each have a portion complementary to nucleotide sequences flanking the other side of the target nucleotide sequence. The primers in the right set are complementary to one strand of the nucleic acid molecule containing the target nucleotide sequence and the
25 primers in the left set are complementary to the opposite strand. The 5' end of primers in both sets are distal to the nucleic acid sequence of interest when the primers are hybridized to the flanking sequences in the nucleic acid molecule. Preferably, each member of each set has a portion complementary to a separate and non-overlapping nucleotide sequence flanking the target nucleotide sequence. Amplification proceeds
30 by replication initiated at each primer and continuing through the target nucleic acid sequence. A key feature of this method is the displacement of intervening primers during replication. Once the nucleic acid strands elongated from the right set of

primers reaches the region of the nucleic acid molecule to which the left set of primers hybridizes, and vice versa, another round of priming and replication will take place. This allows multiple copies of a nested set of the target nucleic acid sequence to be synthesized in a short period of time.

5 Multiple strand displacement amplification can be performed by (a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the target sequence in the primer-target sample mixture, and (b) mixing DNA polymerase with the primer-target sample mixture, to
10 produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the target sequence. Strand displacement replication is preferably accomplished by using a strand displacing DNA polymerase or a DNA polymerase in combination with a compatible strand displacement factor.

15 By using a sufficient number of primers in the right and left sets, only a few rounds of replication are required to produce hundreds of thousands of copies of the nucleic acid sequence of interest. For example, it can be estimated that, using right and left primer sets of 26 primers each, 200,000 copies of a 5000 nucleotide amplification target can be produced in 10 minutes (representing just four rounds of priming and
20 replication). It can also be estimated that, using right and left primer sets of 26 primers each, 200,000 copies of a 47,000 nucleotide amplification target can be produced in 60 minutes (again representing four rounds of priming and replication). These calculations are based on a polymerase extension rate of 50 nucleotides per second. It is emphasized that reactions are continuous and isothermal -- no cycling is required.

25 The disclosed method has advantages over the polymerase chain reaction since it can be carried out under isothermal conditions. No thermal cycling is needed because the polymerase at the head of an elongating strand (or a compatible strand-displacement factor) will displace, and thereby make available for hybridization, the strand ahead of it. Other advantages of multiple strand displacement amplification
30 include the ability to amplify very long nucleic acid segments (on the order of 50 kilobases) and rapid amplification of shorter segments (10 kilobases or less). Long nucleic acid segments can be amplified in the disclosed method since there is no

cycling which could interrupt continuous synthesis or allow the formation of artifacts due to rehybridization of replicated strands. In multiple strand displacement amplification, single priming events at unintended sites will not lead to artifactual amplification at these sites (since amplification at the intended site will quickly outstrip
5 the single strand replication at the unintended site).

In another form of the method, referred to as gene specific strand displacement amplification (GS-MSDA), target DNA is first digested with a restriction endonuclease. The digested fragments are then ligated end-to-end to form DNA circles. These circles can be monomers or concatemers. Two sets of primers are used for amplification, a
10 right set and a left set. Primers in the right set of primers each have a portion complementary to nucleotide sequences flanking one side of a target nucleotide sequence and primers in the left set of primers each have a portion complementary to nucleotide sequences flanking the other side of the target nucleotide sequence. The primers in the right set are complementary to one strand of the nucleic acid molecule
15 containing the target nucleotide sequence and the primers in the left set are complementary to the opposite strand. The primers are designed to cover all or part of the sequence needed to be amplified. Preferably, each member of each set has a portion complementary to a separate and non-overlapping nucleotide sequence flanking the target nucleotide sequence. Amplification proceeds by replication initiated at each
20 primer and continuing through the target nucleic acid sequence. In one form of GS-MSDA, referred to as linear GS-MSDA, amplification is performed with a set of primers complementary to only one strand, thus amplifying only one of the strands. In another form of GS-MSDA, cDNA sequences can be circularized to form single stranded DNA circles. Amplification is then performed with a set of primers
25 complementary to the single-stranded circular cDNA.

C. Multiple Strand Displacement Amplification of Concatenated DNA

In another form of the method, referred to as multiple strand displacement amplification of concatenated DNA (MSDA-CD), concatenated DNA is amplified. A preferred form of concatenated DNA is concatenated cDNA. Concatenated DNA can
30 be amplified using a random or partially random set of primers, as in WGSDA, or using specific primers complementary to specific hybridization targets in the concatenated DNA. MSDA-CD is preferred for amplification of a complex mixture or sample of

relatively short nucleic acid samples (that is, fragments generally in the range of 100 to 6,000 nucleotides). Messenger RNA is the most important example of such a complex mixture. MSDA-CD provides a means for amplifying all cDNAs in a cell in equal fashion. Because the concatenated cDNA can be amplified up to 5,000-fold, MSDA-
5 CD will permit RNA profiling analysis based on just a few cells. To perform MSDA-CD, DNA must first be subjected to a concatenation step. If an RNA sample (such as mRNA) is to be amplified, the RNA is first converted to a double-stranded cDNA using standard methods. The cDNA, or any other set of DNA fragments to be amplified, is then converted into a DNA concatenate, preferably with incorporation of linkers.

10 **D. Multiple Strand Displacement Amplification of Damaged DNA**

Other forms of the disclosed method can involve amplification and repair of damaged DNA. Amplification of damaged DNA can be both difficult and provide unreliable results. For example, amplification of degraded or fragmented DNA will produce truncated products and can result in allele dropout. Preparation of genomic
15 DNA samples in particular can result in damage to the genomic DNA (for example, degradation and fragmentation). Damaged DNA and damaged DNA samples can be amplified and repaired in the disclosed method of amplifying damaged DNA. The method generally works by hybridizing the ends of some DNA molecules in a damaged DNA sample to complementary sequences in the sample. Because the DNA molecules
20 providing the newly associated ends will have damage at different locations, priming from the annealed ends can result in replication of more complete fragments and can mediate repair of the damaged DNA (in the form of less damaged or undamaged replicated strands). Replication of the undamaged replicated strands by continued multiple displacement amplification produces less damaged or undamaged amplified
25 nucleic acids.

The method generally involves substantially denaturing a damaged DNA sample (generally via exposure to heat and alkaline conditions), removal or reduction of the denaturing conditions (such as by reduction of the pH and temperature of the denatured DNA sample), and replicating the DNA. The damaged DNA is repaired
30 during replication and the average length of DNA fragments is increased. For example, the average length of DNA fragments can be increased from, for example, 2 kb in the damaged DNA sample to, for example, 10 kb or greater for the replicated DNA. The

amplified and repaired DNA is in better condition for analysis and testing than the damaged DNA sample. For example, this technique can provide consistent improvements in allele representation from damaged DNA samples. This repair method can result in an overall improvement in amplification of damaged DNA by
5 increasing the average length of the product, increasing the quality of the amplification products by 3-fold (by, for example, increasing the marker representation in the sample), and improving the genotyping of amplified products by lowering the frequency of allelic dropout; all compared to the results when amplifying damaged DNA by other methods. The replication can be multiple displacement amplification.
10 Denaturation of the DNA sample generally is carried out such that the DNA is not further damaged. This method can generally be combined or used with any of the disclosed amplification methods.

In some embodiments, the method of amplifying damaged DNA can involve exposing a damaged DNA sample to conditions that promote substantial denaturation
15 of damaged DNA in the damaged DNA sample, thereby forming a denatured damaged DNA sample; altering the conditions to conditions that do not promote substantial denaturation of damaged DNA in the damaged DNA sample to form a stabilized denatured damaged DNA sample; and incubating the annealed damaged DNA under conditions that promote replication of the damaged DNA. The annealed ends of the
20 damaged DNA prime replication and replication of the damaged DNA results in repair of the replicated strands. The conditions that promote substantial denaturation of damaged DNA in the damaged DNA sample can be, for example, raising the pH of the damaged DNA sample and heating the damaged DNA sample. The altering conditions can be, for example, reducing the pH of the denatured damaged DNA sample and
25 cooling the damaged DNA sample. Raising the pH can be accomplished by exposing the damaged DNA sample to alkaline conditions. The altering conditions generally can be conditions that promote annealing of the ends of the transiently denatured damaged DNA to the substantially denatured damaged DNA. The damaged DNA sample, the denatured damaged DNA sample, or both can also be exposed to ionic conditions by,
30 for example, mixing the damaged DNA sample or denatured damaged DNA sample with an ionic solution or including salt(s) or other ions in the denaturing solution, the stabilization solution, or both.

In the method, the damaged DNA sample can be exposed to conditions that promote substantial denaturation by, for example, mixing the damaged DNA sample with a denaturing solution and by heating the damaged DNA sample to a temperature and for a length of time that substantially denatures the damaged DNA in the damaged DNA sample. The temperature can be, for example, about 25°C to about 50°C and the length of time can be, for example, about 5 minutes or more. The pH of the denatured damaged DNA sample can be reduced, for example, by mixing the denatured damaged DNA sample with a stabilization solution. The damaged DNA samples can be, for example, degraded DNA fragments of genomic DNA. Replication and repair of the damaged DNA can be accomplished by incubating the damaged DNA in the presence of a DNA polymerase, such as ϕ 29 DNA polymerase.

The damaged DNA sample can generally be slowly cooled in order to achieve the required annealing. For example, the damaged DNA sample can be cooled at a rate of, for example, about 0.1°C per minute or less, about 0.2°C per minute or less, about 0.3°C per minute or less, about 0.4°C per minute or less, about 0.5°C per minute or less, about 0.6°C per minute or less, about 0.7°C per minute or less, about 0.8°C per minute or less, about 0.9°C per minute or less, about 1°C per minute or less, about 1.0°C per minute or less, about 1.1°C per minute or less, about 1.2°C per minute or less, about 1.3°C per minute or less, about 1.4°C per minute or less, about 1.5°C per minute or less, about 1.6°C per minute or less, about 1.8°C per minute or less, about 2°C per minute or less, about 2.0°C per minute or less, about 2.2°C per minute or less, about 2.4°C per minute or less, about 2.6°C per minute or less, about 2.8°C per minute or less, about 3°C per minute or less, about 3.0°C per minute or less, about 3.5°C per minute or less, about 4°C per minute or less, about 4.0°C per minute or less, about 5.0°C per minute or less, about 0.1°C per minute, about 0.2°C per minute, about 0.3°C per minute, about 0.4°C per minute, about 0.5°C per minute, about 0.6°C per minute, about 0.7°C per minute, about 0.8°C per minute, about 0.9°C per minute, about 1°C per minute, about 1.0°C per minute, about 1.1°C per minute, about 1.2°C per minute, about 1.3°C per minute, about 1.4°C per minute, about 1.5°C per minute, about 1.6°C per minute, about 1.8°C per minute, about 2°C per minute, about 2.0°C per minute, about 2.2°C per minute, about 2.4°C per minute, about 2.6°C per minute, about 2.8°C per minute, about 3°C per minute, about 3.0°C per minute, about 3.5°C per minute, about

4°C per minute, about 4.0°C per minute, about 5.0°C per minute, 0.1°C per minute, 0.2°C per minute, 0.3°C per minute, 0.4°C per minute, 0.5°C per minute, 0.6°C per minute, 0.7°C per minute, 0.8°C per minute, 0.9°C per minute, 1°C per minute, 1.0°C per minute, 1.1°C per minute, 1.2°C per minute, 1.3°C per minute, 1.4°C per minute, 1.5°C per minute, 1.6°C per minute, 1.8°C per minute, 2°C per minute, 2.0°C per minute, 2.2°C per minute, 2.4°C per minute, 2.6°C per minute, 2.8°C per minute, 3°C per minute, 3.0°C per minute, 3.5°C per minute, 4°C per minute, 4.0°C per minute, or 5.0°C per minute.

The rate of cooling of the damaged DNA sample can also be described in terms of the percent drop in temperature. Thus, cooling a damaged DNA sample that starts at 70°C at a rate of 1% per minute or less would be cooled by 0.7°C (or less) in the first minute and 1% (or less) of the resulting temperature in the next minute. The damaged DNA sample can be cooled at a rate of, for example, about 0.1% per minute or less, about 0.2% per minute or less, about 0.3% per minute or less, about 0.4% per minute or less, about 0.5% per minute or less, about 0.6% per minute or less, about 0.7% per minute or less, about 0.8% per minute or less, about 0.9% per minute or less, about 1% per minute or less, about 1.0% per minute or less, about 1.1% per minute or less, about 1.2% per minute or less, about 1.3% per minute or less, about 1.4% per minute or less, about 1.5% per minute or less, about 1.6% per minute or less, about 1.8% per minute or less, about 2% per minute or less, about 2.0% per minute or less, about 2.2% per minute or less, about 2.4% per minute or less, about 2.6% per minute or less, about 2.8% per minute or less, about 3% per minute or less, about 3.0% per minute or less, about 3.5% per minute or less, about 4% per minute or less, about 4.0% per minute or less, about 5.0% per minute or less, about 0.1% per minute, about 0.2% per minute, about 0.3% per minute, about 0.4% per minute, about 0.5% per minute, about 0.6% per minute, about 0.7% per minute, about 0.8% per minute, about 0.9% per minute, about 1% per minute, about 1.0% per minute, about 1.1% per minute, about 1.2% per minute, about 1.3% per minute, about 1.4% per minute, about 1.5% per minute, about 1.6% per minute, about 1.8% per minute, about 2% per minute, about 2.0% per minute, about 2.2% per minute, about 2.4% per minute, about 2.6% per minute, about 2.8% per minute, about 3% per minute, about 3.0% per minute, about 3.5% per minute, about 4% per minute, about 4.0% per minute, about 5.0% per minute, 0.1% per minute, 0.2% per

minute, 0.3% per minute, 0.4% per minute, 0.5% per minute, 0.6% per minute, 0.7% per minute, 0.8% per minute, 0.9% per minute, 1% per minute, 1.0% per minute, 1.1% per minute, 1.2% per minute, 1.3% per minute, 1.4% per minute, 1.5% per minute, 1.6% per minute, 1.8% per minute, 2% per minute, 2.0% per minute, 2.2% per minute, 2.4% per minute, 2.6% per minute, 2.8% per minute, 3% per minute, 3.0% per minute, 3.5% per minute, 4% per minute, 4.0% per minute, or 5.0% per minute.

The damaged DNA sample, the denatured damaged DNA sample, or both can also be exposed to ionic conditions by, for example, mixing the damaged DNA sample or denatured damaged DNA sample with an ionic solution or including salt(s) or other ions in the denaturing solution, the stabilization solution, or both. As used herein, ionic conditions refers to a state of increased ionic strength. Thus, exposure to ionic conditions refers to exposure to a higher ionic strength than existed in the sample or solution before exposure. This will be the result when, for example, a buffer or salt is added. A solution used to make such an addition can be referred to as an ionic solution. The ionic solution can be a salt solution and can comprise one or more salts or other ions. Any suitable salt or ion can be used. The salt can be, for example, Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination. The Tris-HCl can be, for example, from pH 7.0 to 8.0. The salt can be Tris-EDTA. The ionic solution can be diluted, for example, 2 to 5 fold when mixed with the damaged DNA sample. The ionic solution can be mixed with the denatured damaged DNA sample prior to or during altering of the conditions.

Ionic conditions and the composition of ionic solutions generally can be specified by specifying a concentration of a buffer, salt, or other ion-forming compound. A combination of compounds can be used in an ionic solution or to create ionic conditions. The salt solution can comprise, for example, about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA. Ionic solutions can have a salt, buffer or ion concentration of from about 1 mM to about 2 M, from about 1 mM to about 1 M, from about 1 mM to about 900 mM, from about 1 mM to about 800 mM, from about 1 mM to about 700 mM, from about 1 mM to about 600 mM, from about 1 mM to about 500 mM, from about 1 mM to about 400 mM, from about 1 mM to about 300 mM, from about 1 mM to about 250 mM, from about 1 mM to about 200 mM,

from about 1 mM to about 150 mM, from about 1 mM to about 100 mM, from about 1 mM to about 80 mM, from about 1 mM to about 60 mM, from about 1 mM to about 50 mM, from about 1 mM to about 40 mM, from about 1 mM to about 30 mM, from about 1 mM to about 20 mM, from about 1 mM to about 10 mM, from about 1 mM to about 5 mM, from about 1 mM to about 2 mM, from about 2 mM to about 2 M, from about 2 mM to about 1 M, from about 2 mM to about 900 mM, from about 2 mM to about 800 mM, from about 2 mM to about 700 mM, from about 2 mM to about 600 mM, from about 2 mM to about 500 mM, from about 2 mM to about 400 mM, from about 2 mM to about 300 mM, from about 2 mM to about 250 mM, from about 2 mM to about 200 mM, from about 2 mM to about 150 mM, from about 2 mM to about 100 mM, from about 2 mM to about 80 mM, from about 2 mM to about 60 mM, from about 2 mM to about 50 mM, from about 2 mM to about 40 mM, from about 2 mM to about 30 mM, from about 2 mM to about 20 mM, from about 2 mM to about 10 mM, from about 2 mM to about 5 mM, from about 5 mM to about 2 M, from about 5 mM to about 1 M, from about 5 mM to about 900 mM, from about 5 mM to about 800 mM, from about 5 mM to about 700 mM, from about 5 mM to about 600 mM, from about 5 mM to about 500 mM, from about 5 mM to about 400 mM, from about 5 mM to about 300 mM, from about 5 mM to about 250 mM, from about 5 mM to about 200 mM, from about 5 mM to about 150 mM, from about 5 mM to about 100 mM, from about 5 mM to about 80 mM, from about 5 mM to about 60 mM, from about 5 mM to about 50 mM, from about 5 mM to about 40 mM, from about 5 mM to about 30 mM, from about 5 mM to about 20 mM, from about 5 mM to about 10 mM, from about 10 mM to about 2 M, from about 10 mM to about 1 M, from about 10 mM to about 900 mM, from about 10 mM to about 800 mM, from about 10 mM to about 700 mM, from about 10 mM to about 600 mM, from about 10 mM to about 500 mM, from about 10 mM to about 400 mM, from about 10 mM to about 300 mM, from about 10 mM to about 250 mM, from about 10 mM to about 200 mM, from about 10 mM to about 150 mM, from about 10 mM to about 100 mM, from about 10 mM to about 80 mM, from about 10 mM to about 60 mM, from about 10 mM to about 50 mM, from about 10 mM to about 40 mM, from about 10 mM to about 30 mM, from about 10 mM to about 20 mM, from about 20 mM to about 2 M, from about 20 mM to about 1 M, from about 20 mM to about 900 mM, from about 20 mM to about 800 mM, from about 20 mM to about 700 mM, from about

20 mM to about 600 mM, from about 20 mM to about 500 mM, from about 20 mM to about 400 mM, from about 20 mM to about 300 mM, from about 20 mM to about 250 mM, from about 20 mM to about 200 mM, from about 20 mM to about 150 mM, from about 20 mM to about 100 mM, from about 20 mM to about 80 mM, from about 20 mM to about 60 mM, from about 20 mM to about 50 mM, from about 20 mM to about 40 mM, from about 20 mM to about 30 mM, from about 30 mM to about 2 M, from about 30 mM to about 1 M, from about 30 mM to about 900 mM, from about 30 mM to about 800 mM, from about 30 mM to about 700 mM, from about 30 mM to about 600 mM, from about 30 mM to about 500 mM, from about 30 mM to about 400 mM, from about 30 mM to about 300 mM, from about 30 mM to about 250 mM, from about 30 mM to about 200 mM, from about 30 mM to about 150 mM, from about 30 mM to about 100 mM, from about 30 mM to about 80 mM, from about 30 mM to about 60 mM, from about 30 mM to about 50 mM, from about 30 mM to about 40 mM, from about 40 mM to about 2 M, from about 40 mM to about 1 M, from about 40 mM to about 900 mM, from about 40 mM to about 800 mM, from about 40 mM to about 700 mM, from about 40 mM to about 600 mM, from about 40 mM to about 500 mM, from about 40 mM to about 400 mM, from about 40 mM to about 300 mM, from about 40 mM to about 250 mM, from about 40 mM to about 200 mM, from about 40 mM to about 150 mM, from about 40 mM to about 100 mM, from about 40 mM to about 80 mM, from about 40 mM to about 60 mM, from about 40 mM to about 50 mM, from about 50 mM to about 2 M, from about 50 mM to about 1 M, from about 50 mM to about 900 mM, from about 50 mM to about 800 mM, from about 50 mM to about 700 mM, from about 50 mM to about 600 mM, from about 50 mM to about 500 mM, from about 50 mM to about 400 mM, from about 50 mM to about 300 mM, from about 50 mM to about 250 mM, from about 50 mM to about 200 mM, from about 50 mM to about 150 mM, from about 50 mM to about 100 mM, from about 50 mM to about 80 mM, from about 50 mM to about 60 mM, from about 60 mM to about 2 M, from about 60 mM to about 1 M, from about 60 mM to about 900 mM, from about 60 mM to about 800 mM, from about 60 mM to about 700 mM, from about 60 mM to about 600 mM, from about 60 mM to about 500 mM, from about 60 mM to about 400 mM, from about 60 mM to about 300 mM, from about 60 mM to about 250 mM, from about 60 mM to about 200 mM, from about 60 mM to about 150 mM, from about 60 mM to about 100

mM, from about 60 mM to about 80 mM, from about 80 mM to about 2 M, from about 80 mM to about 1 M, from about 80 mM to about 900 mM, from about 80 mM to about 800 mM, from about 80 mM to about 700 mM, from about 80 mM to about 600 mM, from about 80 mM to about 500 mM, from about 80 mM to about 400 mM, from about 80 mM to about 300 mM, from about 80 mM to about 250 mM, from about 80 mM to about 200 mM, from about 80 mM to about 150 mM, from about 80 mM to about 100 mM, from about 100 mM to about 2 M, from about 100 mM to about 1 M, from about 100 mM to about 900 mM, from about 100 mM to about 800 mM, from about 100 mM to about 700 mM, from about 100 mM to about 600 mM, from about 100 mM to about 500 mM, from about 100 mM to about 400 mM, from about 100 mM to about 300 mM, from about 100 mM to about 250 mM, from about 100 mM to about 200 mM, from about 100 mM to about 150 mM, from about 150 mM to about 2 M, from about 150 mM to about 1 M, from about 150 mM to about 900 mM, from about 150 mM to about 800 mM, from about 150 mM to about 700 mM, from about 150 mM to about 600 mM, from about 150 mM to about 500 mM, from about 150 mM to about 400 mM, from about 150 mM to about 300 mM, from about 150 mM to about 250 mM, from about 150 mM to about 200 mM, from about 200 mM to about 2 M, from about 200 mM to about 1 M, from about 200 mM to about 900 mM, from about 200 mM to about 800 mM, from about 200 mM to about 700 mM, from about 200 mM to about 600 mM, from about 200 mM to about 500 mM, from about 200 mM to about 400 mM, from about 200 mM to about 300 mM, from about 200 mM to about 250 mM, from about 250 mM to about 2 M, from about 250 mM to about 1 M, from about 250 mM to about 900 mM, from about 250 mM to about 800 mM, from about 250 mM to about 700 mM, from about 250 mM to about 600 mM, from about 250 mM to about 500 mM, from about 250 mM to about 400 mM, from about 250 mM to about 300 mM, from about 300 mM to about 2 M, from about 300 mM to about 1 M, from about 300 mM to about 900 mM, from about 300 mM to about 800 mM, from about 300 mM to about 700 mM, from about 300 mM to about 600 mM, from about 300 mM to about 500 mM, from about 300 mM to about 400 mM, from about 400 mM to about 2 M, from about 400 mM to about 1 M, from about 400 mM to about 900 mM, from about 400 mM to about 800 mM, from about 400 mM to about 700 mM, from about 400 mM to about 600 mM, from about 400 mM to about 500 mM, from about 500 mM to about 2 M, from about

500 mM to about 1 M, from about 500 mM to about 900 mM, from about 500 mM to about 800 mM, from about 500 mM to about 700 mM, from about 500 mM to about 600 mM, from about 600 mM to about 2 M, from about 600 mM to about 1 M, from about 600 mM to about 900 mM, from about 600 mM to about 800 mM, from about 600 mM to about 700 mM, from about 700 mM to about 2 M, from about 700 mM to about 1 M, from about 700 mM to about 900 mM, from about 700 mM to about 800 mM, from about 800 mM to about 2 M, from about 800 mM to about 1 M, from about 800 mM to about 900 mM, from about 900 mM to about 2 M, from about 900 mM to about 1 M, from about 1 M to about 2 M, about 2 M, about 1 M, about 900 mM, about 800 mM, about 700 mM, about 600 mM, about 500 mM, about 400 mM, about 300 mM, about 250 mM, about 200 mM, about 150 mM, about 100 mM, about 80 mM, about 60 mM, about 50 mM, about 40 mM, about 30 mM, about 20 mM, about 10 mM, about 9 mM, about 8 mM, about 7 mM, about 6 mM, about 5 mM, about 4 mM, about 3 mM, about 2 mM, about 1 mM, 2 M, 1 M, 900 mM, 800 mM, 700 mM, 600 mM, 500 mM, 400 mM, 300 mM, 250 mM, 200 mM, 150 mM, 100 mM, 80 mM, 60 mM, 50 mM, 40 mM, 30 mM, 20 mM, 10 mM, 9 mM, 8 mM, 7 mM, 6 mM, 5 mM, 4 mM, 3 mM, 2 mM, or 1 mM.

The disclosed method of repairing and amplifying DNA can result in an increase in the average length of DNA fragment in a DNA sample. This increase can be referred to in any suitable terms. For example, the increase in average fragment length can be referred to by the average fragment length of the replicated DNA fragments, the increase in average fragment length from the average fragment length of the damaged DNA sample, and the percent increase in average fragment length. The increase in average fragment length can be, for example, 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 260%, 280%, 300%, 350%, 400%, 450%, 500%, 600%, 700%, 800%, 900%, 1,000%, 5% or more, 10% or more, 15% or more, 20% or more, 25% or more, 30% or more, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, 90% or more, 100% or more, 110% or more, 120% or more, 130% or more, 140% or more, 150% or more, 160% or more, 180% or more, 200% or more, 220% or more, 240% or more, 260% or more, 280% or more, 300% or more, 350% or more, 400% or more, 450% or more, 500% or more, 600% or more, 700% or more,

800% or more, 900% or more, 1,000% or more, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 140%, about 150%, about 160%, about 180%, about 200%, about 220%, about 240%,
 5 about 260%, about 280%, about 300%, about 350%, about 400%, about 450%, about 500%, about 600%, about 700%, about 800%, about 900%, about 1,000%, about 5% or more, about 10% or more, about 15% or more, about 20% or more, about 25% or more, about 30% or more, about 40% or more, about 50% or more, about 60% or more, about 70% or more, about 80% or more, about 90% or more, about 100% or more, about
 10 110% or more, about 120% or more, about 130% or more, about 140% or more, about 150% or more, about 160% or more, about 180% or more, about 200% or more, about 220% or more, about 240% or more, about 260% or more, about 280% or more, about 300% or more, about 350% or more, about 400% or more, about 450% or more, about 500% or more, about 600% or more, about 700% or more, about 800% or more, about
 15 900% or more, or about 1,000% or more relative to the average fragment length of the damaged DNA sample before the method.

Following the repair method, the average fragment length can be, for example, 2 kilobases (kb), 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, 5 kb, 5.5 kb, 6 kb, 7 kb, 8 kb, 9 kb, 10 kb, 11 kb, 12 kb, 13 kb, 14 kb, 15 kb, 16 kb, 18 kb, 20 kb, 22 kb, 24 kb, 26 kb, 28
 20 kb, 30 kb, 2 kb or more, 2.5 kb or more, 3 kb or more, 3.5 kb or more, 4 kb or more, 4.5 kb or more, 5 kb or more, 5.5 kb or more, 6 kb or more, 7 kb or more, 8 kb or more, 9 kb or more, 10 kb or more, 11 kb or more, 12 kb or more, 13 kb or more, 14 kb or more, 15 kb or more, 16 kb or more, 18 kb or more, 20 kb or more, 22 kb or more, 24 kb or more, 26 kb or more, 28 kb or more, 30 kb or more, about 2 kb, about 2.5 kb,
 25 about 3 kb, about 3.5 kb, about 4 kb, about 4.5 kb, about 5 kb, about 5.5 kb, about 6 kb, about 7 kb, about 8 kb, about 9 kb, about 10 kb, about 11 kb, about 12 kb, about 13 kb, about 14 kb, about 15 kb, about 16 kb, about 18 kb, about 20 kb, about 22 kb, about 24 kb, about 26 kb, about 28 kb, about 30 kb, about 2 kb or more, about 2.5 kb or more, about 3 kb or more, about 3.5 kb or more, about 4 kb or more, about 4.5 kb or more,
 30 about 5 kb or more, about 5.5 kb or more, about 6 kb or more, about 7 kb or more, about 8 kb or more, about 9 kb or more, about 10 kb or more, about 11 kb or more, about 12 kb or more, about 13 kb or more, about 14 kb or more, about 15 kb or more,

about 16 kb or more, about 18 kb or more, about 20 kb or more, about 22 kb or more, about 24 kb or more, about 26 kb or more, about 28 kb or more, or about 30 kb or more.

5 The disclosed method of amplifying damaged DNA can be combined with the disclosed amplification of cell lysates. Thus, for example, the damaged DNA samples can be a cell lysate, where the cell lysate is produced by exposing cells to alkaline conditions. Some forms of the method can include exposing cells to alkaline conditions to form a cell lysate; exposing the cell lysate to conditions that promote substantial denaturation of damaged DNA in the cell lysate; reducing the pH of the cell lysate to
10 form a stabilized cell lysate; cooling the stabilized cell lysate under conditions that promote annealing of the ends of the denatured damaged DNA; and incubating the stabilized cell lysate under conditions that promote replication of the damaged DNA. During replication, the annealed ends of the damaged DNA prime replication and replication of the damaged DNA results in repair of the replicated strands and an
15 increase in the average length of DNA fragment. The cell lysate can be a whole genome. Replication of the genome results in replicated strands, where during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

In another form, the method works by hybridizing the ends of some DNA
20 molecules in a sample to complementary sequences in a damaged DNA sample. Generally, the damaged DNA sample and the DNA sample providing the annealed ends or from the same source or even the same sample. Because the DNA molecules providing the newly associated ends and damaged DNA molecules will have damage at different locations, priming from the annealed ends can result in replication of more
25 complete fragments and can mediate repair of the damaged DNA (in the form of less damaged or undamaged replicated strands). Replication of the undamaged replicated strands by continued multiple displacement amplification produces less damaged or undamaged amplified nucleic acids.

The method generally involves substantially denaturing a damaged DNA
30 sample (generally via exposure to heat and alkaline conditions), reduction of the pH of the denatured DNA sample, mixing the denatured DNA sample with an undenatured DNA sample from the same source such that the ends of DNA in the undenatured DNA

sample is transiently denatured, slowly cooling the mixture of DNA samples to allow the transiently denatured ends to anneal to the denatured DNA, and replicating the annealed DNA. The damaged DNA is repaired during replication. The replication can be multiple displacement amplification. Substantial denaturation and transient
5 denaturation of the DNA samples generally is carried out such that the DNA is not further damaged. This method can generally be combined or used with any of the disclosed amplification methods.

In some embodiments, the method of amplifying damaged DNA can involve exposing a first damaged DNA sample to conditions that promote substantial
10 denaturation of damaged DNA in the first damaged DNA sample, thereby forming a denatured damaged DNA sample; reducing the pH of the denatured damaged DNA sample to form a stabilized denatured damaged DNA sample; mixing a second damaged DNA sample with the stabilized denatured damaged DNA sample under conditions that promote transient denaturation of the ends of damaged DNA in the
15 second sample and that maintain substantial denaturation of the damaged DNA in the stabilized denatured damaged DNA sample, thereby forming a damaged DNA mixture; cooling the damaged DNA mixture under conditions that promote annealing of the ends of the transiently denatured damaged DNA to the substantially denatured damaged DNA; and incubating the annealed damaged DNA under conditions that promote
20 replication of the damaged DNA. The annealed ends of the damaged DNA prime replication and replication of the damaged DNA results in repair of the replicated strands.

In the method, the first damaged DNA sample can be exposed to conditions that promote substantial denaturation by, for example, mixing the first damaged DNA
25 sample with a denaturing solution and by heating the first damaged DNA sample to a temperature and for a length of time that substantially denatures the damaged DNA in the first damaged DNA sample. The temperature can be, for example, about 25°C to about 50°C and the length of time can be, for example, about 5 minutes or more. The pH of the denatured damaged DNA sample can be reduced, for example, by mixing the
30 denatured damaged DNA sample with a stabilization solution. The damaged DNA samples can be, for example, degraded DNA fragments of genomic DNA. The first and second damaged DNA samples can be from the same source, and in particular can

be a portion of the same damaged DNA sample. The second damaged DNA sample can be mixed with the stabilized denatured damaged DNA sample at a temperature and for a length of time that transiently denatures the damaged DNA in the second damaged DNA sample. For example, the temperature can be about 70°C or less and the length
5 of time can be about 30 seconds or less. The desired effect can also be achieved by maintaining the mixture at the temperature to which the first damaged DNA sample is exposed for denaturation. Replication and repair of the damaged DNA can be accomplished by incubating the annealed damaged DNA in the presence of a DNA polymerase, such as ϕ 29 DNA polymerase.

10 The disclosed method of amplifying damaged DNA can be combined with the disclosed amplification of cell lysates. Thus, for example, the first and second damaged DNA samples can be portions of a cell lysate, where the cell lysate is produced by exposing cells to alkaline conditions. The pH of the second damaged DNA sample can be reduced prior to mixing with the stabilized denatured damaged
15 DNA. Some forms of the method can include exposing cells to alkaline conditions to form a cell lysate; exposing a first portion of the cell lysate to conditions that promote substantial denaturation of damaged DNA in the first portion of the cell lysate; reducing the pH of the first portion of the cell lysate to form a first stabilized cell lysate and reducing the pH of a second portion of the cell lysate to form a second stabilized
20 cell lysate; mixing the second stabilized cell lysate with the first stabilized cell lysate under conditions that promote transient denaturation of the ends of damaged DNA in the second stabilized cell lysate and that maintain substantial denaturation of the damaged DNA in the first stabilized cell lysate, thereby forming a stabilized cell lysate mixture; cooling the stabilized cell lysate mixture under conditions that promote
25 annealing of the ends of the transiently denatured damaged DNA to the substantially denatured damaged DNA; and incubating the stabilized cell lysate mixture under conditions that promote replication of the damaged DNA. During replication, the annealed ends of the damaged DNA prime replication and replication of the damaged DNA results in repair of the replicated strands. The cell lysate can be a whole genome.
30 Replication of the genome results in replicated strands, where during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

E. Modifications And Additional Operations

1. Detection of Amplification Products

Amplification products can be detected directly by, for example, primary labeling or secondary labeling, as described below.

5 i. Primary Labeling

Primary labeling consists of incorporating labeled moieties, such as fluorescent nucleotides, biotinylated nucleotides, digoxigenin-containing nucleotides, or bromodeoxyuridine, during strand displacement replication. For example, one may incorporate cyanine dye deoxyuridine analogs (Yu *et al.*, *Nucleic Acids Res.*, **22**:3226-3232 (1994)) at a frequency of 4 analogs for every 100 nucleotides. A preferred method for detecting nucleic acid amplified *in situ* is to label the DNA during amplification with BrdUrd, followed by binding of the incorporated BrdU with a biotinylated anti-BrdU antibody (Zymed Labs, San Francisco, CA), followed by binding of the biotin moieties with Streptavidin-Peroxidase (Life Sciences, Inc.), and finally development of fluorescence with Fluorescein-tyramide (DuPont de Nemours & Co., Medical Products Dept.). Other methods for detecting nucleic acid amplified *in situ* include labeling the DNA during amplification with 5-methylcytosine, followed by binding of the incorporated 5-methylcytosine with an antibody (Sano *et al.*, *Biochim. Biophys. Acta* **951**:157-165 (1988)), or labeling the DNA during amplification with aminoallyl-deoxyuridine, followed by binding of the incorporated aminoallyl-deoxyuridine with an Oregon Green[®] dye (Molecular Probes, Eugene, OR) (Henegariu *et al.*, *Nature Biotechnology* **18**:345-348 (2000)).

Another method of labeling amplified nucleic acids is to incorporate 5-(3-aminoallyl)-dUTP (AAdUTP) in the nucleic acid during amplification followed by chemical labeling at the incorporated nucleotides. Incorporated 5-(3-aminoallyl)-deoxyuridine (AAdU) can be coupled to labels that have reactive groups that are capable of reacting with amine groups. AAdUTP can be prepared according to Langer *et al.* (1981). *Proc. Natl. Acad. Sci. USA.* **78**: 6633-37. Other modified nucleotides can be used in analogous ways. That is, other modified nucleotides with minimal modification can be incorporated during replication and labeled after incorporation.

Examples of labels suitable for addition to AAdUTP are radioactive isotopes, fluorescent molecules, phosphorescent molecules, enzymes, antibodies, and ligands.

Examples of suitable fluorescent labels include fluorescein isothiocyanate (FITC), 5,6-carboxymethyl fluorescein, Texas red, nitrobenz-2-oxa-1,3-diazol-4-yl (NBD), coumarin, dansyl chloride, rhodamine, amino-methyl coumarin (AMCA), Eosin, Erythrosin, BODIPY[®], Cascade Blue[®], Oregon Green[®], pyrene, lissamine, xanthenes, acridines, oxazines, phycoerythrin, macrocyclic chelates of lanthanide ions such as quantum dye[™], fluorescent energy transfer dyes, such as thiazole orange-ethidium heterodimer, and the cyanine dyes Cy3, Cy3.5, Cy5, Cy5.5 and Cy7. Examples of other specific fluorescent labels include 3-Hydroxypyrene 5,8,10-Tri Sulfonic acid, 5-Hydroxy Tryptamine (5-HT), Acid Fuchsin, Alizarin Complexon, Alizarin Red, Allophycocyanin, Aminocoumarin, Anthroyl Stearate, Astrazon Brilliant Red 4G, Astrazon Orange R, Astrazon Red 6B, Astrazon Yellow 7 GLL, Atabrine, Auramine, Aurophosphine, Aurophosphine G, BAO 9 (Bisaminophenyloxadiazole), BCECF, Berberine Sulphate, Bisbenzamide, Blancophor FFG Solution, Blancophor SV, Bodipy F1, Brilliant Sulphoflavin FF, Calcien Blue, Calcium Green, Calcofluor RW Solution, Calcofluor White, Calcophor White ABT Solution, Calcophor White Standard Solution, Carbostyryl, Cascade Yellow, Catecholamine, Chinacrine, Coriphosphine O, Coumarin-Phalloidin, CY3.1 8, CY5.1 8, CY7, Dans (1-Dimethyl Amino Naphaline 5 Sulphonic Acid), Dansa (Diamino Naphtyl Sulphonic Acid), Dansyl NH-CH₃, Diamino Phenyl Oxydiazole (DAO), Dimethylamino-5-Sulphonic acid, Dipyrrometheneboron Difluoride, Diphenyl Brilliant Flavine 7GFF, Dopamine, Erythrosin ITC, Euchrysin, FIF (Formaldehyde Induced Fluorescence), Flazo Orange, Fluo 3, Fluorescamine, Fura-2, Genacryl Brilliant Red B, Genacryl Brilliant Yellow 10GF, Genacryl Pink 3G, Genacryl Yellow 5GF, Gloxalic Acid, Granular Blue, Haematoporphyrin, Indo-1, Intrawhite Cf Liquid, Leucophor PAF, Leucophor SF, Leucophor WS, Lissamine Rhodamine B200 (RD200), Lucifer Yellow CH, Lucifer Yellow VS, Magdala Red, Marina Blue, Maxilon Brilliant Flavin 10 GFF, Maxilon Brilliant Flavin 8 GFF, MPS (Methyl Green Pyronine Stilbene), Mithramycin, NBD Amine, Nitrobenzoxadidole, Noradrenaline, Nuclear Fast Red, Nuclear Yellow, Nylosan Brilliant Flavin E8G, Oxadiazole, Pacific Blue, Pararosanine (Feulgen), Phorwite AR Solution, Phorwite BKL, Phorwite Rev, Phorwite RPA, Phosphine 3R, Phthalocyanine, Phycoerythrin R, Polyazaindacene Pontochrome Blue Black, Porphyrin, Primuline, Procion Yellow, Pyronine, Pyronine B, Pyrozal Brilliant Flavin 7GF, Quinacrine Mustard, Rhodamine

123, Rhodamine 5 GLD, Rhodamine 6G, Rhodamine B, Rhodamine B 200, Rhodamine B Extra, Rhodamine BB, Rhodamine BG, Rhodamine WT, Serotonin, Sevron Brilliant Red 2B, Sevron Brilliant Red 4G, Sevron Brilliant Red B, Sevron Orange, Sevron Yellow L, SITS (Primuline), SITS (Stilbene Isothiosulphonic acid), Stilbene, Snarf 1, sulpho Rhodamine B Can C, Sulpho Rhodamine G Extra, Tetracycline, Thiazine Red R, Thioflavin S, Thioflavin TCN, Thioflavin 5, Thiolyte, Thiozol Orange, Tinopol CBS, True Blue, Ultralite, Uranine B, Uvitex SFC, Xylene Orange, and XRITC.

Preferred fluorescent labels are fluorescein (5-carboxyfluorescein-N-hydroxysuccinimide ester), rhodamine (5,6-tetramethyl rhodamine), and the cyanine dyes Cy3, Cy3.5, Cy5, Cy5.5 and Cy7. The absorption and emission maxima, respectively, for these fluors are: FITC (490 nm; 520 nm), Cy3 (554 nm; 568 nm), Cy3.5 (581 nm; 588 nm), Cy5 (652 nm; 672 nm), Cy5.5 (682 nm; 703 nm) and Cy7 (755 nm; 778 nm), thus allowing their simultaneous detection. Other examples of fluorescein dyes include 6-carboxyfluorescein (6-FAM), 2',4',1,4,- tetrachlorofluorescein (TET), 2',4',5',7',1,4-hexachlorofluorescein (HEX), 2',7'-dimethoxy-4', 5'-dichloro-6-carboxyrhodamine (JOE), 2'-chloro-5'-fluoro-7',8'-fused phenyl-1,4-dichloro-6-carboxyfluorescein (NED), and 2'-chloro-7'-phenyl-1,4-dichloro-6-carboxyfluorescein (VIC). Fluorescent labels can be obtained from a variety of commercial sources, including Amersham Pharmacia Biotech, Piscataway, NJ; Molecular Probes, Eugene, OR; and Research Organics, Cleveland, Ohio.

ii. Secondary Labeling with Detection Probes

Secondary labeling consists of using suitable molecular probes, referred to as detection probes, to detect the amplified nucleic acids. For example, a primer may be designed to contain, in its non-complementary portion, a known arbitrary sequence, referred to as a detection tag. A secondary hybridization step can be used to bind detection probes to these detection tags. The detection probes may be labeled as described above with, for example, an enzyme, fluorescent moieties, or radioactive isotopes. By using three detection tags per primer, and four fluorescent moieties per each detection probe, one may obtain a total of twelve fluorescent signals for every replicated strand.

iii. Multiplexing and Hybridization Array Detection

Detection of amplified nucleic acids can be multiplexed by using sets of different primers, each set designed for amplifying different target sequences. Only those primers that are able to find their targets will give rise to amplified products.

5 There are two alternatives for capturing a given amplified nucleic acid to a fixed position in a solid-state detector. One is to include within the non-complementary portion of the primers a unique address tag sequence for each unique set of primers. Nucleic acid amplified using a given set of primers will then contain sequences corresponding to a specific address tag sequence. A second and preferred alternative is
10 to use a sequence present in the target sequence as an address tag.

iv. Enzyme-linked Detection

Amplified nucleic acid labeled by incorporation of labeled nucleotides can be detected with established enzyme-linked detection systems. For example, amplified nucleic acid labeled by incorporation of biotin using biotin-16-UTP (Roche Molecular
15 Biochemicals) can be detected as follows. The nucleic acid is immobilized on a solid glass surface by hybridization with a complementary DNA oligonucleotide (address probe) complementary to the target sequence (or its complement) present in the amplified nucleic acid. After hybridization, the glass slide is washed and contacted with alkaline phosphatase-streptavidin conjugate (Tropix, Inc., Bedford, MA). This
20 enzyme-streptavidin conjugate binds to the biotin moieties on the amplified nucleic acid. The slide is again washed to remove excess enzyme conjugate and the chemiluminescent substrate CSPD (Tropix, Inc.) is added and covered with a glass cover slip. The slide can then be imaged in a Biorad Fluorimager.

2. Linear Strand Displacement Amplification

25 A modified form of multiple strand displacement amplification can be performed which results in linear amplification of a target sequence. This modified method is referred to as linear strand displacement amplification (LSDA) and is accomplished by using a set of primers where all of the primers are complementary to the same strand of the target sequence. In LSDA, as in MSDA, the set of primers
30 hybridize to the target sequence and strand displacement amplification takes place. However, only one of the strands of the target sequence is replicated. LSDA requires thermal cycling between each round of replication to allow a new set of primers to

hybridize to the target sequence. Such thermal cycling is similar to that used in PCR. Unlike linear, or single primer, PCR, however, each round of replication in LSDA results in multiple copies of the target sequence. One copy is made for each primer used. Thus, if 20 primers are used in LSDA, 20 copies of the target sequence will be made in each cycle of replication.

DNA amplified using MSDA and WGSDA can be further amplified by transcription. For this purpose, promoter sequences can be included in the non-complementary portion of primers used for strand displacement amplification, or in linker sequences used to concatenate DNA for MSDA-CD.

10 **3. Reverse Transcription Multiple Displacement Amplification**

Multiple displacement amplification can be performed on RNA or on DNA strands reverse transcribed from RNA. A useful form of the disclosed method, referred to as reverse transcription multiple displacement amplification (RT-MDA) involves reverse transcribing RNA, removal of the RNA (preferably by nuclease digestion using an RNA-specific nuclease such as RNase H), and multiple displacement amplification of the reverse transcribed DNA. RT-MDA can be performed using either double-stranded cDNA or using just the first cDNA strand. In the latter case, the second cDNA strand need not be, and preferably is not, synthesized. RT-MDA is useful for quantitative analysis of mRNA or general amplification of mRNA sequences for any other purpose.

4. Repeat Multiple Displacement Amplification

The disclosed multiple displacement amplification operations can also be sequentially combined. For example, the product of MDA can itself be amplified in another multiple displacement amplification. This is referred to herein as repeat multiple displacement amplification (RMDA). This can be accomplished, for example, by diluting the replicated strands following MDA and subjecting them to a new MDA. This can be repeated one or more times. Each round of MDA will increase the amplification. Different forms of MDA, such as WGSDA and MSDA on particular target sequences can be combined. In general, repeat MDA can be accomplished by first bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. Replication of the target sequence results in replicated strands, wherein

during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand; and then diluting the replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the replicated strands under

5 conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, wherein during replication at least one of the additional replicated strands is displaced from the target sequence by strand displacement replication of another additional replicated strand. This form of the method can be extended by performing the following operation one or more times:

10 diluting the additional replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the replicated strands under conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, wherein during replication at least one of the additional replicated strands is displaced from the target sequence by

15 strand displacement replication of another additional replicated strand.

5. Using Products of Multiple Displacement Amplification

The nucleic acids produced using the disclosed method can be used for any purpose. For example, the amplified nucleic acids can be analyzed (such as by sequencing or probe hybridization) to determine characteristics of the amplified

20 sequences or the presence or absence of certain sequences. The amplified nucleic acids can also be used as reagents for assays or other methods. For example, nucleic acids produced in the disclosed method can be coupled or adhered to a solid-state substrate. The resulting immobilized nucleic acids can be used as probes or indexes of sequences in a sample. Nucleic acids produced in the disclosed method can be coupled or adhered

25 to a solid-state substrate in any suitable way. For example, nucleic acids generated by multiple strand displacement can be attached by adding modified nucleotides to the 3' ends of nucleic acids produced by strand displacement replication using terminal deoxynucleotidyl transferase, and reacting the modified nucleotides with a solid-state substrate or support thereby attaching the nucleic acids to the solid-state substrate or

30 support.

Nucleic acids produced in the disclosed method also can be used as probes or hybridization partners. For example, sequences of interest can be amplified in the

disclosed method and provide a ready source of probes. The replicated strands (produced in the disclosed method) can be cleaved prior to use as hybridization probes. For example, the replicated strands can be cleaved with DNase I. The hybridization probes can be labeled as described elsewhere herein with respect to labeling of nucleic acids produce in the disclosed method.

Nucleic acids produced in the disclosed method also can be used for subtractive hybridization to identify sequences that are present in only one of a pair or set of samples. For example, amplified cDNA from different samples can be annealed and the resulting double-stranded material can be separated from single-stranded material. Unhybridized sequences would be indicative of sequences expressed in one of the samples but not others.

Specific Embodiments

Disclosed is a method of amplifying a target nucleic acid sequence where the method can comprise bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The target sample is not subjected to denaturing conditions. The method can further comprise labeling the replicated strands using terminal deoxynucleotidyl transferase.

The method can further comprise diluting the replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the replicated strands under conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, where during replication at least one of the additional replicated strands is displaced from the target sequence by strand displacement replication of another additional replicated strand. The method can further comprise performing the following operation one or more times: diluting the additional replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the replicated strands under conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, where during

replication at least one of the additional replicated strands is displaced from the target sequence by strand displacement replication of another additional replicated strand. The method can further comprise incubating the polymerase-target sample mixture under conditions that promote strand displacement. The method can further comprise
5 bringing into contact a set of primers, DNA polymerase, and a second target sample, and incubating the second target sample under conditions that promote replication of the target sequence. The second target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target
10 sequence by strand displacement replication of another replicated strand.

The primers can be 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides long. The primers can be 5, 6, 7, 8, 9, or 10 nucleotides long. The primers can be 5, 6, 7, or 8 nucleotides long. The primers can be 6, 7, or 8 nucleotides long. The primers can be 6 nucleotides long. The primers each can contain at least one
15 modified nucleotide such that the primers are resistant to 3'-5' exonuclease. The DNA polymerase can be bacteriophage ϕ 29 DNA polymerase, Tts DNA polymerase, phage M2 DNA polymerase, VENT™ DNA polymerase, Klenow fragment of DNA polymerase I, T5 DNA polymerase, PRD1 DNA polymerase, T4 DNA polymerase holoenzyme, T7 native polymerase T7 Sequenase®, or Bst DNA polymerase. The
20 DNA polymerase can be ϕ 29 DNA polymerase.

The primers can be 6 nucleotides long, the primers each can contain at least one modified nucleotide such that the primers are nuclease resistant, and the DNA polymerase can be ϕ 29 DNA polymerase. The replicated strands can be labeled by the addition of modified nucleotides to the replicated strands. The modified nucleotides
25 can be biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates. Modified nucleotides can be incorporated into the replicated strands during replication. The modified nucleotides can be biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates. The modified nucleotides can be 5-(3-
30 aminoallyl)-2'-deoxyuridine 5'-triphosphates and the replicated strands can be labeled by reacting labels with the incorporated 5-(3-aminoallyl)-2'-deoxyuridines. The labels can be fluorescein isothiocyanate, 5,6-carboxymethyl fluorescein, Texas red, nitrobenz-

2-oxa-1,3-diazol-4-yl, coumarin, dansyl chloride, rhodamine, amino-methyl coumarin, Eosin, Erythrosin, BODIPY[®], Cascade Blue[®], Oregon Green[®], pyrene, lissamine, xanthene, acridine, oxazines, phycoerythrin, Cy3, Cy3.5, Cy5, Cy5.5, Cy7, or a combination thereof.

- 5 The target sample need not be subjected to heat denaturing conditions. The target sequence can comprise two strands, and the set of primers can have 3 or more primers complementary to one of the strands of the target sequence and at least one primer complementary to the other strand of the target sequence. The set of primers can have 3 or more primers complementary to the same strand of the target sequence.
- 10 The set of primers can have 4 or more primers complementary to the same strand of the target sequence. The set of primers can have 4 or more primers. The set of primers can have 5 or more primers. The conditions that promote replication of the target sequence can be substantially isothermic. The conditions that promote replication of the target sequence need not involve thermal cycling. The conditions need not include thermal
- 15 cycling. The target sequence can comprise an amplification target and a hybridization target, the hybridization target can flank the amplification target, the set of primers can comprise a plurality of primers, each primer can comprise a complementary portion, where the complementary portions of the primers each can be complementary to a different portion of the hybridization target.
- 20 The set of primers can comprise a right set of primers and a left set of primers, the target sequence can be double-stranded, having a first and a second strand, the hybridization target can comprise a right and left hybridization target, where the right hybridization target can flank the amplification target on one end and the left hybridization target can flank the amplification target on the other end. The
- 25 complementary portions of the right set primers can be (i) all complementary to the first strand of the target sequence and (ii) each complementary to a different portion of the right hybridization target, and the complementary portions of the left set primers can be (i) all complementary to the second strand of the target sequence and (ii) each complementary to a different portion of the left hybridization target.
- 30 The right and left set of primers each can have 3 or more primers. The right and left set of primers each can have 4 or more primers. The right and left set of primers each can have 5 or more primers. The right and left set of primers each can have the

same number of primers. The target sequence can be a nucleic acid sample of substantial complexity, and the set of primers can comprise primers having random nucleotide sequences. The target sequence can be a sample of genomic nucleic acid. The primers can be from 5 to 20 nucleotides in length. The primers can be from 5 to 10
5 nucleotides in length. The primers can be 6, 7, or 8 nucleotides in length. The primers can be 6 nucleotides in length. The primers can be all of the same length. Each primer can comprise a constant portion and a random portion, where the constant portion of each primer can have the same nucleotide sequence and the random portion of each primer can have a random nucleotide sequence.

10 The target sequence can be concatenated DNA. The concatenated DNA can be concatenated with linkers. Each linker can comprise a primer complement portion, each primer can comprise a complementary portion, and the complementary portion of each primer can be complementary to the complementary portion of the linkers. The set of primers can comprise primers having random nucleotide sequences. Each primer
15 can comprise a constant portion and a random portion, the constant portion of each primer can have the same nucleotide sequence and the random portion of each primer can have a random nucleotide sequence. The concatenated DNA can be formed by ligating DNA fragments together. The DNA fragments can be cDNA made from mRNA. The mRNA can comprise a mixture of mRNA isolated from cells. The target
20 sequence need not be a nucleic acid molecule made up of multiple tandem repeats of a single sequence that was synthesized by rolling circle replication.

The primers can comprise nucleotides, where one or more of the nucleotides can be ribonucleotides. From about 10% to about 50% of the nucleotides can be ribonucleotides. About 50% or more of the nucleotides can be ribonucleotides. All of
25 the nucleotides can be ribonucleotides. The primers can comprise nucleotides, where one or more of the nucleotides can be 2'-O-methyl ribonucleotides. From about 10% to about 50% of the nucleotides can be 2'-O-methyl ribonucleotides. About 50% or more of the nucleotides can be 2'-O-methyl ribonucleotides. All of the nucleotides can be 2'-O-methyl ribonucleotides. The primers can comprise nucleotides and the nucleotides
30 can be a mixture of ribonucleotides and 2'-O-methyl ribonucleotides. The primers can comprise nucleotides, the nucleotides can comprise bases, and one or more of the bases can be universal bases. At least one of the universal bases can be 3-nitropyrrole. The

universal base can be 5-nitroindole. From about 10% to about 50% of the bases can be universal bases. About 50% or more of the bases can be universal bases. All of the bases can be universal bases.

5 The target sample can be a biopsy sample, a blood sample, a urine sample, a cell sample, or a tissue sample. The target sample is a needle aspiration biopsy sample. Nucleic acids in the target sample need not be separated from other material in the target sample. The target sample can be a crude cell lysate. The target sample need not be processed beyond cell lysis. The replicated strands can be analyzed. The replicated strands can be analyzed using one or more DNA chips. The replicated strands can be
10 analyzed by hybridization. The replicated strands can be analyzed by nucleic acid sequencing. The replicated strands can be stored prior to, following, or both prior to and following their analysis. The target sample can be a blood sample, a urine sample, a semen sample, a lymphatic fluid sample, a cerebrospinal fluid sample, amniotic fluid sample, a biopsy sample, a needle aspiration biopsy sample, a cancer sample, a tumor
15 sample, a tissue sample, a cell sample, a cell lysate sample, a crude cell lysate sample, a forensic sample, an archeological sample, an infection sample, a nosocomial infection sample, a production sample, a drug preparation sample, a biological molecule production sample, a protein preparation sample, a lipid preparation sample, a carbohydrate preparation sample, or a combination thereof.

20 The target sample can be a blood sample. The target sample can be a needle aspiration biopsy sample. The target sample can be a crude cell lysate sample. The target sample can be a nosocomial infection sample. The sample can be derived from a patient. The target sample can be a biological molecule production sample. Production of replicated strands can indicate the presence of nucleic acids in the sample. The
25 amount of replicated strands produced can indicate the amount of nucleic acids in the sample. The target sample can be a drug preparation sample. The target sample can be a tumor sample. The target sample can be an amniotic fluid sample. The replicated strands produced from the target sample can represent a nucleic acid fingerprint of the sample.

30 The second target sample can be a sample from the same type of organism as the first target sample. The second target sample can be a sample from the same type of tissue as the first target sample. The second target sample can be a sample from the

same organism as the first target sample. The second target sample can be obtained at a different time than the first target sample. The second target sample can be a sample from a different organism than the first target sample. The second target sample can be a sample from a different type of tissue than the first target sample. The second target sample can be a sample from a different species of organism than the first target sample. The second target sample can be a sample from a different strain of organism than the first target sample. The second target sample can be a sample from a different cellular compartment than the first target sample.

A circular nucleic acid molecule can comprise the target sequence. The circular nucleic acid molecule can be produced by digesting genomic DNA with a restriction endonuclease, and circularizing the digested DNA. The digested DNA can be circularized with DNA or RNA ligase. The digested DNA can be circularized with a splint or adaptor. The target sequence can comprise an amplification target and a hybridization target, the hybridization target can flank the amplification target, the set of primers can comprise a plurality of primers, each primer can comprise a complementary portion, and the complementary portions of the primers each can be complementary to a different portion of the hybridization target.

The set of primers can comprise a right set of primers and a left set of primers, the target sequence can be double-stranded, having a first and a second strand, the hybridization target can comprise a right and left hybridization target, and the right hybridization target can flank the amplification target on one end and the left hybridization target flanks the amplification target on the other end. The complementary portions of the right set primers can be (i) all complementary to the first strand of the target sequence and (ii) each complementary to a different portion of the right hybridization target, and the complementary portions of the left set primers can be (i) all complementary to the second strand of the target sequence and (ii) each complementary to a different portion of the left hybridization target.

The circular nucleic acid molecule can be produced by circularizing cDNA. The circular nucleic acid molecule is produced by circularizing mRNA/cDNA hybrid. The mRNA/cDNA hybrid can be circularized with DNA or RNA ligase. The mRNA/cDNA hybrid can be circularized with a splint or adaptor.

Also disclosed is a method of amplifying a target nucleic acid sequence where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. Nucleic acids in the target sample are not separated
5 from other material in the target sample. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

The method can further comprise bringing into contact a set of primers, DNA
10 polymerase, and a second target sample, and incubating the second target sample under conditions that promote replication of the target sequence. The second target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another
15 replicated strand.

The target sample can be a crude cell lysate. The target sample need not be processed beyond cell lysis. The target sample can be a blood sample, a urine sample, a semen sample, a lymphatic fluid sample, a cerebrospinal fluid sample, amniotic fluid sample, a biopsy sample, a needle aspiration biopsy sample, a cancer sample, a tumor
20 sample, a tissue sample, a cell sample, a cell lysate sample, a crude cell lysate sample, a forensic sample, an archeological sample, an infection sample, a nosocomial infection sample, a production sample, a drug preparation sample, a biological molecule production sample, a protein preparation sample, a lipid preparation sample, a carbohydrate preparation sample, or a combination thereof. The target sample can be a
25 blood sample. The target sample can be a needle aspiration biopsy sample. The target sample can be a crude cell lysate sample. The target sample can be a nosocomial infection sample. The sample can be derived from a patient. The target sample can be a biological molecule production sample.

Production of replicated strands can indicate the presence of nucleic acids in the
30 sample. The amount of replicated strands produced can indicate the amount of nucleic acids in the sample. The target sample can be a drug preparation sample. The target sample can be a tumor sample. The target sample can be an amniotic fluid sample.

The replicated strands produced from the target sample can represent a nucleic acid fingerprint of the sample.

The second target sample can be a sample from the same type of organism as the first target sample. The second target sample can be a sample from the same type of tissue as the first target sample. The second target sample can be a sample from the same organism as the first target sample. The second target sample can be obtained at a different time than the first target sample. The second target sample can be a sample from a different organism than the first target sample. The second target sample can be a sample from a different type of tissue than the first target sample. The second target sample can be a sample from a different species of organism than the first target sample. The second target sample can be a sample from a different strain of organism than the first target sample. The second target sample can be a sample from a different cellular compartment than the first target sample.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is a crude cell lysate. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The primers are 5, 6, 7, 8, 9, or 10 nucleotides long. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The primers each contain at least one modified nucleotide such that the primers are nuclease resistant. Replication of the target

sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
5 the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The primer-target sample is not subjected to denaturing conditions, the primers are 6 nucleotides long, the primers each contain at least one modified nucleotides such that the primers are nuclease resistant, and the
10 DNA polymerase is ϕ 29 DNA polymerase. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
15 the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another
20 replicated strand. The method further comprises diluting the replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the replicated strands under conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, where during replication at least one of the additional replicated
25 strands is displaced from the target sequence by strand displacement replication of another additional replicated strand.

The method can further comprise performing the following operation one or more times: diluting the additional replicated strands, bringing into contact a set of primers, DNA polymerase, and the diluted replicated strands, and incubating the
30 replicated strands under conditions that promote replication of the target sequence. Replication of the target sequence results in additional replicated strands, where during

replication at least one of the additional replicated strands is displaced from the target sequence by strand displacement replication of another additional replicated strand.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises,

5 (a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the target sequence in the primer-target sample mixture, where the primer-target sample is not subjected to denaturing conditions, and

10 (b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the target sequence.

The set of primers comprises a right set of primers and a left set of primers, the target sequence is double-stranded, having a first and a second strand, where the right
15 set primers are all complementary to the first strand of the target sequence and the left set primers are all complementary to the second strand of the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The right set of primers can have
20 4 or more primers and the left set of primers has 4 or more primers.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises,

(a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that
25 promote hybridization between the primers and the target sequence in the primer-target sample mixture, where the primer-target sample is not subjected to denaturing conditions, and

(b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample
30 mixture under conditions that promote replication of the target sequence.

Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by

strand displacement replication of another replicated strand. The target sequence is a nucleic acid sample of substantial complexity, and the set of primers comprises primers having random nucleotide sequences.

Also disclosed is a method of amplifying a target nucleic acid sequence, the
5 method comprising,

(a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the target sequence in the primer-target sample mixture, where the primer-target sample is not subjected to denaturing
10 conditions, and

(b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the target sequence.

All of the primers in the set of primers are complementary to the same strand in
15 the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The set of primers can have 3 or more primers.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
20 the method comprises,

(a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the target sequence in the primer-target sample mixture, where the primer-target sample is not subjected to denaturing
25 conditions, and

(b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the target sequence.

Replication of the target sequence results in replicated strands, where during
30 replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The target sequence is a nucleic acid sample of substantial complexity, and the set of primers comprises primers

having random nucleotide sequences. Each primer comprises a constant portion and a random portion, where the constant portion of each primer has the same nucleotide sequence and the random portion of each primer has a random nucleotide sequence.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
5 the method comprises,

(a) mixing a set of primers with a target sample, to produce a primer-target sample mixture, and incubating the primer-target sample mixture under conditions that promote hybridization between the primers and the target sequence in the primer-target sample mixture, where the primer-target sample is not subjected to denaturing
10 conditions, and

(b) mixing DNA polymerase with the primer-target sample mixture, to produce a polymerase-target sample mixture, and incubating the polymerase-target sample mixture under conditions that promote replication of the target sequence.

Replication of the target sequence results in replicated strands, where during
15 replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The conditions that promote replication of the target sequence do not involve thermal cycling and the target sequence is concatenated DNA.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
20 the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where
25 during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The set of primers can have 3 or more primers complementary to the same strand of the target sequence.

Also disclosed is a method of amplifying a target nucleic acid sequence, where
the method comprises, bringing into contact a set of primers, DNA polymerase, and a
30 target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where

during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The target sequence is a nucleic acid sample of substantial complexity, and the set of primers comprises primers having random nucleotide sequences.

5 Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to denaturing conditions. All of the primers in the set of primers are complementary to the same
10 strand in the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The set of primers can have 3 or more primers.

 Also disclosed is a method of amplifying a target nucleic acid sequence, where
15 the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where
20 during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The target sequence is a nucleic acid sample of substantial complexity, and the set of primers comprises primers having random nucleotide sequences. Each primer comprises a constant portion and a random portion, where the constant portion of each primer has the same nucleotide sequence and the random portion of each primer has a random
25 nucleotide sequence.

 Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to denaturing
30 conditions. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand. The

conditions that promote replication of the target sequence do not involve thermal cycling, and the target sequence is concatenated DNA.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. The target sample is not subjected to heat denaturing conditions. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed is a method of labeling nucleic acids produced by strand displacement replication, where the method comprises, labeling nucleic acids produced by strand displacement replication using terminal deoxynucleotidyl transferase. The replicated strands can be labeled by the addition of modified nucleotides to the 3' ends of the nucleic acids. The modified nucleotides can be biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates.

Also disclosed is a method of labeling nucleic acids produced by strand displacement replication, where the method comprises, incorporating modified nucleotides into nucleic acids produced by strand displacement replication during replication.

Also disclosed is a method of attaching nucleic acids produced by strand displacement replication, where the method comprises, adding modified nucleotides to the 3' ends of nucleic acids produced by strand displacement replication using terminal deoxynucleotidyl transferase, and reacting the modified nucleotides with a solid-state support thereby attaching the nucleic acids to the solid-state support.

Also disclosed is a microarray comprising nucleic acids produced by strand displacement replication coupled or adhered to a solid-state substrate.

Also disclosed is a method of generating probes based on a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is

displaced from the target sequence by strand displacement replication of another replicated strand. The replicated strands are used as hybridization probes. The method can further comprise labeling the hybridization probes using terminal deoxynucleotidyl transferase.

5 Also disclosed is a method of amplifying messenger RNA, where the method comprises, reverse transcribing messenger RNA to produce a first strand cDNA, bringing into contact a set of primers, DNA polymerase, and the first strand cDNA, and incubating under conditions that promote replication of the first strand cDNA. Replication of the first strand cDNA results in replicated strands, where during
10 replication at least one of the replicated strands is displaced from the first strand cDNA by strand displacement replication of another replicated strand. The method can further comprise, prior to replication, degrading the messenger RNA using RNase H.

 The method can further comprise bringing into contact a set of primers, DNA polymerase, and a second target sample, and incubating the second target sample under
15 conditions that promote replication of the target sequence. The second target sample is not subjected to denaturing conditions. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

20 Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, partially degrading RNA in a target sample, bringing into contact DNA polymerase, and the target sample, and incubating the target sample under conditions that promote replication of the target sequence. Replication of the target
25 sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

 Also disclosed is a method of comparative genome hybridization, where the method comprises, hybridizing nucleic acids produced by strand displacement replication of a first sample with nucleic acids produced by strand displacement
30 replication of a second sample. Hybridization can be carried out in the absence of *CotI* DNA.

Also disclosed is a method of amplifying a target nucleic acid sequence, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, and incubating the target sample under conditions that promote replication of the target sequence. A circular nucleic acid molecule comprises the target sequence. Replication of the target sequence results in replicated strands, where during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

The set of primers can comprise primers having random nucleotide sequences. The primers can be 5, 6, 7, or 8 nucleotides long. The primers can be 6, 7, or 8 nucleotides long. The primers can be 6 nucleotides long. The modified nucleotides can be biotinylated nucleotides, fluorescent nucleotides, 5 methyl dCTP, BrdUTP, or 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates. The modified nucleotides can be 5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphates and the replicated strands can be labeled by reacting labels with the incorporated 5-(3-aminoallyl)-2'-deoxyuridines. The labels can be fluorescein isothiocyanate, 5,6-carboxymethyl fluorescein, Texas red, nitrobenz-2-oxa-1,3-diazol-4-yl, coumarin, dansyl chloride, rhodamine, amino-methyl coumarin, Eosin, Erythrosin, BODIPY[®], Cascade Blue[®], Oregon Green[®], pyrene, lissamine, xanthene, acridine, oxazines, phycoerythrin, Cy3, Cy3.5, Cy5, Cy5.5, Cy7, or a combination thereof.

The replicated strands can be used as elements in a microarray. The replicated strands can be cleaved prior to use as hybridization probes. The replicated strands can be cleaved with DNase I. The hybridization probes can be labeled by the addition of modified nucleotides to the 3' ends of the replicated strands. The modified nucleotides can be reacted with a solid-state support thereby attaching the replicated strands to the solid-state support.

The target sample can be a blood sample, a urine sample, a semen sample, a lymphatic fluid sample, a cerebrospinal fluid sample, amniotic fluid sample, a biopsy sample, a needle aspiration biopsy sample, a cancer sample, a tumor sample, a tissue sample, a cell sample, a cell lysate sample, a crude cell lysate sample, a forensic sample, an archeological sample, an infection sample, a nosocomial infection sample, a production sample, a drug preparation sample, a biological molecule production sample, a protein preparation sample, a lipid preparation sample, a carbohydrate

preparation sample, or a combination thereof. The target sample can be a blood sample. The target sample is a needle aspiration biopsy sample. The target sample can be a crude cell lysate sample. The target sample can be a nosocomial infection sample. The sample can contain both human and non-human nucleic acids. The non-human
5 nucleic acid can be amplified preferentially by the use of primers specific for the non-human nucleic acid. The human nucleic acid can be amplified preferentially by the use of primers specific for human nucleic acid. The sample can be derived from a patient. The target sample can be a biological molecule production sample. Production of replicated strands can indicate the presence of nucleic acids in the sample. The amount
10 of replicated strands produced can indicate the amount of nucleic acids in the sample. The target sample can be a drug preparation sample. The target sample can be a tumor sample. The target sample can be an amniotic fluid sample. The replicated strands produced from the target sample can represent a nucleic acid fingerprint of the sample.

The second target sample can be a sample from the same type of organism as
15 the first target sample. The second target sample can be a sample from the same type of tissue as the first target sample. The second target sample can be a sample from the same organism as the first target sample. The second target sample can be obtained at a different time than the first target sample. The second target sample can be a sample from a different organism than the first target sample. The second target sample can be
20 a sample from a different type of tissue than the first target sample. The second target sample can be a sample from a different species of organism than the first target sample. The second target sample can be a sample from a different strain of organism than the first target sample. The second target sample can be a sample from a different cellular compartment than the first target sample.

25 The circular nucleic acid molecule can be produced by digesting genomic DNA with a restriction endonuclease, and circularizing the digested DNA. The digested DNA can be circularized with DNA or RNA ligase. The digested DNA can be circularized with a splint or adaptor. The target sequence can comprises an amplification target and a hybridization target, the hybridization target can flank the
30 amplification target, the set of primers can comprise a plurality of primers, each primer can comprise a complementary portion, and the complementary portions of the primers each can be complementary to a different portion of the hybridization target.

The set of primers can comprise a right set of primers and a left set of primers, the target sequence can be double-stranded, having a first and a second strand, the hybridization target can comprise a right and left hybridization target, the right hybridization target can flank the amplification target on one end and the left hybridization target can flank the amplification target on the other end. The complementary portions of the right set primers can be (i) all complementary to the first strand of the target sequence and (ii) each complementary to a different portion of the right hybridization target, and the complementary portions of the left set primers can be (i) all complementary to the second strand of the target sequence and (ii) each complementary to a different portion of the left hybridization target.

The circular nucleic acid molecule can be produced by circularizing cDNA. The circular nucleic acid molecule can be produced by circularizing mRNA/cDNA hybrid. The mRNA/cDNA hybrid can be circularized with DNA or RNA ligase. The mRNA/cDNA hybrid can be circularized with a splint or adaptor.

Also disclosed is a method of amplifying a whole genome, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, where the target sample comprises a whole genome, and incubating the target sample under conditions that promote replication of the genome. The target sample is not subjected to denaturing conditions. Replication of the genome results in replicated strands, where during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand. The primers can be 6 nucleotides long, the primers each can contain at least one modified nucleotide such that the primers are nuclease resistant, and the DNA polymerase can be ϕ 29 DNA polymerase. The primers can be of random nucleotide composition, the primers each can contain at least one modified nucleotide such that the primers are nuclease resistant, and the DNA polymerase can be ϕ 29 DNA polymerase.

Also disclosed is a method of amplifying a chromosome, where the method comprises, bringing into contact a set of primers, DNA polymerase, and a target sample, where the target sample comprises a chromosome, and incubating the target sample under conditions that promote replication of the chromosome, where the target sample is not subjected to denaturing conditions. Replication of the chromosome results in replicated strands, where during replication at least one of the replicated

strands is displaced from the chromosome by strand displacement replication of another replicated strand. The primers can be 6 nucleotides long, the primers each can contain at least one modified nucleotide such that the primers are nuclease resistant, and the DNA polymerase can be ϕ 29 DNA polymerase. The primers can be of random
5 nucleotide composition, the primers each can contain at least one modified nucleotide such that the primers are nuclease resistant, and the DNA polymerase can be ϕ 29 DNA polymerase.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome,
10 neutralizing the cell lysate to form a neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

15 Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, neutralizing the cell lysate, wherein the cell lysate is not purified, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of
20 the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, neutralizing the cell lysate, wherein nucleic acids in the cell lysate are not separated
25 from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

30 Furthermore, disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, neutralizing the cell lysate to form a neutralized cell lysate, and incubating the

neutralized cell lysate under conditions that promote replication of the genome, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication
5 of another replicated strand.

Also disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, neutralizing the cell lysate to form a neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein
10 the neutralized cell lysate is subjected to heat denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Methods of amplifying a whole genome are also disclosed, the methods
15 comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, neutralizing the cell lysate to form a neutralized cell lysate, wherein nucleic acids in the neutralized cell lysate are not separated from other material in the cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein the neutralized cell lysate is not subjected to
20 denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell
25 lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, wherein the neutralized cell lysate comprises a whole genome, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand
30 displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, bringing into contact a set

of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, wherein the cell lysate is not purified, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the
5 replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a
10 whole genome, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, wherein the neutralized cell lysate comprises a whole genome, and
15 incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.
20

Also disclosed are methods of amplifying a whole genome, the methods
25 comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, wherein the neutralized cell lysate comprises a whole genome, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein the neutralized cell lysate is subjected to heat
30 denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the method comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, wherein the neutralized cell lysate comprises a whole genome, wherein
5 nucleic acids in the neutralized cell lysate are not separated from other material in the neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of the genome, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced
10 from the genome by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the method comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and incubating the neutralized cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a
15 target sequence, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the
20 methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the cell lysate is not purified, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target
25 sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein nucleic acids in
30 the cell lysate are not separated from other material in the cell lysate, wherein replication of the cell lysate results in replicated strands, wherein during replication at

least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to
5 form a neutralized cell lysate, and incubating the neutralized cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced
10 from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and incubating the neutralized cell lysate in the presence
15 of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the neutralized cell lysate is subjected to denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated
20 strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and incubating the neutralized cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication
25 of a target sequence, wherein nucleic acids in the neutralized cell lysate are not separated from other material in the neutralized cell lysate, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement
30 replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to

form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of a target sequence, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein the cell lysate is not purified, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of a target sequence, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, and incubating the neutralized cell lysate
5 under conditions that promote replication of a target sequence, wherein the neutralized cell lysate is subjected to heat denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

10 Disclosed are methods of amplifying a target nucleic acid sequence, the method comprising, lysing cells to form a cell lysate, neutralizing the cell lysate to form a neutralized cell lysate, and bringing into contact a set of primers, DNA polymerase, and the neutralized cell lysate, and incubating the neutralized cell lysate under conditions that promote replication of a target sequence, wherein nucleic acids in the neutralized
15 cell lysate are not separated from other material in the neutralized cell lysate, wherein the neutralized cell lysate is not subjected to denaturing conditions, wherein replication of the neutralized cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

20 Disclosed are methods of amplifying a whole genome, the method comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during
25 replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein the cell lysate is not purified, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome
30 results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, and incubating the cell lysate under conditions that promote replication of the genome, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, and incubating the cell lysate under conditions that promote replication of the genome, wherein the cell lysate is subjected to heat denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a whole genome, the method comprising, lysing cells to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the genome, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a whole genome, the method comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole

genome, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

5 Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, wherein the cell lysate is not purified, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in
10 replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

 Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA
15 polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand
20 displacement replication of another replicated strand.

 Also disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, and incubating the cell lysate under conditions that promote replication of the
25 genome, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

 Disclosed are methods of amplifying a whole genome, the methods comprising,
30 lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, and incubating the cell lysate under conditions that promote replication of the genome,

wherein the cell lysate is subjected to heat denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

5 Disclosed are methods of amplifying a whole genome, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, wherein the cell lysate comprises a whole genome, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, and incubating the cell lysate under conditions that promote replication of the
10 genome, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

 Also disclosed are methods of amplifying a target nucleic acid sequence, the
15 methods comprising, lysing cells to form a cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another
20 replicated strand.

 Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the cell lysate is not purified, wherein
25 replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

 Also disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, and incubating the cell lysate in
30 the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, wherein replication of the cell lysate

results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the
5 methods comprising, lysing cells to form a cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the
10 target sequence by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the method comprising, lysing cells to form a cell lysate, and incubating the cell lysate in the presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein the cell lysate is subjected to heat
15 denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the method comprising, lysing cells to form a cell lysate, and incubating the cell lysate in the
20 presence of a set of primers and DNA polymerase and under conditions that promote replication of a target sequence, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced
25 from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the method comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions
30 that promote replication of a target sequence, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated

strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein the cell lysate is not purified, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Also disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein the cell lysate is subjected to heat denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated strands is

displaced from the target sequence by strand displacement replication of another replicated strand.

Disclosed are methods of amplifying a target nucleic acid sequence, the methods comprising, lysing cells to form a cell lysate, bringing into contact a set of
5 primers, DNA polymerase, and the cell lysate, and incubating the cell lysate under conditions that promote replication of a target sequence, wherein nucleic acids in the cell lysate are not separated from other material in the cell lysate, wherein the cell lysate is not subjected to denaturing conditions, wherein replication of the cell lysate results in replicated strands, wherein during replication at least one of the replicated
10 strands is displaced from the target sequence by strand displacement replication of another replicated strand.

Any of the disclosed methods can be performed wherein the neutralized cell lysate is not subjected to denaturing conditions or wherein the neutralized cell lysate is subjected to heat denaturing conditions.

15 Disclosed are kits for amplifying a whole genome, the kits comprising a solution for lysis, a solution for neutralization, a set of primers, and a DNA polymerase.

Also disclosed are kits for amplifying a whole genome, the kits comprising a solution for cell lysis, a solution for neutralization of a cell lysate, a set of primers, and a DNA polymerase.

20 Disclosed are kits for amplifying a whole genome, the kits comprising a solution for lysing cells, a solution for neutralizing lysed cells, a set of primers, and a DNA polymerase.

Also disclosed are kits for amplifying a whole genome, the kits comprising a composition for lysis, a composition for neutralization, a set of primers, and a DNA
25 polymerase.

Disclosed are kits for amplifying a whole genome, the kits comprising a composition for cell lysis, a composition for neutralization of a cell lysate, a set of primers, and a DNA polymerase.

30 Disclosed are kits for amplifying a whole genome, the kits comprising a composition for lysing cells, a composition for neutralizing lysed cells, a set of primers, and a DNA polymerase.

Disclosed are kits for amplifying a whole genome, the kits comprising a solution for lysis, wherein the solution for lysis is alkaline, a solution for neutralization, a set of primers, wherein the primers are 6 nucleotides long, wherein the primers each contain at least one modified nucleotide such that the primers are nuclease resistant, and a DNA polymerase, wherein the DNA polymerase is ϕ 29 DNA polymerase.

Disclosed are kits for amplifying a whole genome, the kits comprising a solution for cell lysis, wherein the solution for cell lysis is alkaline, a solution for neutralization of a cell lysate, a set of primers, wherein the primers are 6 nucleotides long, wherein the primers each contain at least one modified nucleotide such that the primers are nuclease resistant, and a DNA polymerase, wherein the DNA polymerase is ϕ 29 DNA polymerase.

Also disclosed are kits for amplifying a whole genome, the kits comprising a solution for lysing cells, wherein the solution for lysing cells is alkaline, a solution for neutralizing lysed cells, a set of primers, wherein the primers are 6 nucleotides long, wherein the primers each contain at least one modified nucleotide such that the primers are nuclease resistant, and a DNA polymerase, wherein the DNA polymerase is ϕ 29 DNA polymerase.

Disclosed are methods of amplifying a whole genome, the method comprising, exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate comprises a whole genome, reducing the pH of the cell lysate to form a stabilized cell lysate, and incubating the stabilized cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Also disclosed are methods, wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution or wherein the lysis solution comprises a base or wherein the base is an aqueous base.

Also disclosed are methods, wherein the base is potassium hydroxide, sodium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine,

methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, or DBU (1,8-diazobicyclo[5,4,0]undec-7-ene).

Disclosed are methods, wherein the base is potassium hydroxide.

5 Also disclosed are methods, wherein the lysis solution comprises 400 mM KOH and/or wherein the lysis solution comprises 100 mM dithiothreitol, and 10 mM EDTA or wherein the lysis solution consists of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

Disclosed are methods, wherein the lysis solution comprises a plurality of basic
10 agents and/or wherein the cells are mixed with an equal volume of the lysis solution.

Disclosed are methods, wherein the lysis solution comprises a buffer, as well as methods wherein the buffer is a phosphate buffer, Good buffer, BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, TRICINE, sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-
15 tris, or Bis-tris propane. Also disclosed are methods, wherein the buffer is Tris-HCl at pH 4.1. The disclosed methods can comprises a plurality of buffering agents.

Also disclosed are methods, wherein the pH of the cell lysate is reduced to the range of about pH 7.0 to about pH 6.8.

Also disclosed are methods, wherein the pH of the cell lysate is reduced by
20 mixing the cell lysate with a stabilization solution. Disclosed are methods wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1 as well as methods, wherein the stabilization solution comprises a plurality of buffering agents and/or wherein the cell lysate is mixed with an equal volume of the stabilization solution.

Disclosed are methods, wherein the stabilization solution comprises an acid as
25 well as methods wherein the acid is hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, or a carboxylic acid. Disclosed are methods wherein the carboxylic acid is ethanoic, propanoic, or butanoic acid.

30 Disclosed are methods, wherein the stabilization solution comprises a plurality of acidic agents.

Disclosed are methods, wherein the pH of the cell lysate is reduced to about pH 9.0 or below, about pH 8.5 or below about pH 8.0 or below, or about pH 7.5 or below.

Also disclosed are methods, wherein the pH of the cell lysate is reduced to the range of about pH 9.0 to about pH 6.0, about pH 9.0 to about pH 7.0, about pH 9.0 to about pH 7.5, pH 9.0 to about pH 8.0, pH 8.5 to about pH 6.0, pH 8.5 to about pH 7.0, pH 8.5 to about pH 7.5, pH 8.5 to about pH 8.0, pH 8.0 to about pH 6.0, pH 8.0 to about pH 6.5, pH 8.0 to about pH 7.0, pH 8.0 to about pH 7.5, pH 7.5 to about pH 6.0, or pH 7.5 to about pH 7.0.

Disclosed are methods, wherein nucleic acids in the cell lysate and the stabilized cell lysate are not separated from other material in the cell lysate, wherein the cell lysate and the stabilized cell lysate are not subjected to purification prior to the incubation, wherein the purification comprises separation of nucleic acids in the cell lysate from other material in the cell lysate, wherein the purification comprises centrifugation, extraction, chromatography, filtration, dialysis, or a combination of these, wherein the purification comprises precipitation other than precipitation caused by the alkaline conditions or by the reduction of the pH, wherein the purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these, wherein the cell lysate, stabilized cell lysate, or both are subjected to partial purification prior to the incubation, wherein the partial purification comprises centrifugation, extraction, chromatography, precipitation, filtration, dialysis, or a combination of these, wherein the partial purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these, wherein the cell lysate and the stabilized cell lysate are not subjected to substantial purification prior to the incubation, wherein the substantial purification does not include centrifugation, extraction, chromatography, precipitation, filtration, or dialysis, wherein the substantial purification does not include centrifugation, phenol-chloroform extraction, or column chromatography, wherein the cell lysate, stabilized cell lysate, or both are subjected to centrifugation, extraction, chromatography, precipitation, filtration, or dialysis prior to the incubation, wherein the cell lysate, stabilized cell lysate, or both are subjected to centrifugation, phenol-chloroform extraction, or column chromatography prior to the incubation, wherein the substantial purification comprises centrifugation, extraction, chromatography, filtration, dialysis, or a combination of

these, wherein the substantial purification comprises precipitation other than precipitation caused by the alkaline conditions or by the reduction of the pH, wherein the substantial purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these, wherein the cell lysate and the stabilized cell lysate are not purified prior to the incubation, wherein the cell lysate, stabilized cell lysate, or both are partially purified prior to the incubation, wherein the incubation is substantially isothermic, wherein neither the cell lysate nor the stabilized cell lysate is heated substantially above the temperature of the incubation, wherein neither the cell lysate nor the stabilized cell lysate is subjected to substantial heating above the temperature of the incubation, wherein the cells are not heated substantially above the temperature of the incubation, wherein the cells are not subjected to substantial heating above the temperature of the incubation, wherein the cells are not heated substantially above the temperature at which the cells grow, wherein the cells are not subjected to substantial heating above the temperature at which the cells grow, wherein the cell lysate, stabilized cell lysate, and the cells are not heated substantially above the temperature of the incubation, wherein the cell lysate, stabilized cell lysate, and the cells are not subjected to substantial heating above the temperature of the incubation, wherein the cell lysate, stabilized cell lysate, and the cells are not heated, prior to or during the incubation, substantially above the temperature at which the cells grow, wherein the cell lysate, stabilized cell lysate, and the cells are not subjected to, prior to or during the incubation, substantial heating above the temperature at which the cells grow prior, wherein neither the cell lysate nor the stabilized cell lysate is heated above a temperature and for a time that would cause notable denaturation of the genome, wherein neither the cell lysate nor the stabilized cell lysate is subjected to heating above a temperature and for a time that would cause notable denaturation of the genome, wherein the cells are not lysed by heat, wherein the cells are not heated above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions, and/or wherein the cells are not subjected to heating above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions.

Disclosed are methods of amplifying a whole genome, the methods comprising, exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate

comprises a whole genome, wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution, reducing the pH of the cell lysate to form a stabilized cell lysate, wherein the pH of the cell lysate is reduced by mixing the cell lysate with a stabilization solution, and incubating the stabilized cell lysate under
5 conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand

Disclosed are methods, wherein the lysis solution comprises potassium
10 hydroxide, such as 400 mM KOH.

Also disclosed are methods wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA or wherein the lysis solution consists of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

Disclosed are methods, wherein the cells are mixed with an equal volume of the
15 lysis solution and/or an equal volume of the stabilization solution

Disclosed are methods, wherein the stabilization solution comprises Tris-HCl at pH 4.1 and wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

Also disclosed are methods wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4.1.

Disclosed are methods, wherein the lysis solution consists of 400 mM KOH and
20 10 mM EDTA, wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4, wherein the stabilized cell lysate is incubated in the presence of 37.5 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1 mM deoxynucleotide triphosphates, 50 μM primers, and φ29 DNA Polymerase.

Also disclosed are methods, wherein the stabilized cell lysate is incubated in the
25 presence of 37.5 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1 mM deoxynucleotide triphosphates, 50 μM primers, and φ29 DNA Polymerase by mixing the stabilized cell lysate with one quarter volume of reaction mix, and φ29 DNA Polymerase, wherein the reaction mix consists of 150 mM Tris-HCl, 200 mM KCl, 40
30 mM MgCl₂, 20 mM (NH₄)₂SO₄, 4 mM deoxynucleotide triphosphates, and 0.2 mM primers.

Disclosed are methods of amplifying a whole genome, the methods comprising, exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution, wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA, reducing the pH of the cell lysate to form a stabilized cell lysate, wherein the pH of the cell lysate is reduced by mixing the cell lysate with a stabilization solution, wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1, and incubating the stabilized cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

Disclosed are methods, wherein the cells are mixed with an equal volume of the lysis solution and/or wherein the cell lysate is mixed with an equal volume of the stabilization solution.

Disclosed are kits for amplifying a whole genome, the kits comprising a lysis solution, a stabilization solution, a set of primers, and a DNA polymerase.

Disclosed are kits wherein the lysis solution comprises potassium hydroxide, such as 400 mM KOH.

Also disclosed are kits wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

Disclosed are kits, wherein the stabilization solution comprises Tris-HCl at pH 4.1, wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1, or wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4.1.

Disclosed are kits comprising deoxynucleotide triphosphates.

Disclosed are kits comprising 1M dithiothreitol, 1X Phosphate-Buffered Saline, pH 7.5, and control DNA template, wherein the lysis solution comprises 400 mM KOH and 10 mM EDTA, wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4, wherein the set of primers comprises a reaction mix, wherein the reaction mix comprises 150 mM Tris-HCl, 200 mM KCl, 40 mM MgCl₂, 20 mM (NH₄)₂SO₄, 4 mM deoxynucleotide triphosphates, and 0.2 mM random hexamer primers, wherein the DNA polymerase is ϕ 29 DNA polymerase.

Disclosed are kits comprising one or more detection probes, wherein the detection probes each comprise a complementary portion, wherein the complementary portion is complementary to a nucleic acid sequence of interest.

Disclosed are kits, wherein the kit is designed to detect nucleic acid sequences of interest in the genome and/or kits wherein the kit is designed to assess a disease, condition or predisposition of an individual based on the nucleic acid sequences of interest.

Disclosed are methods of amplifying damaged DNA, the method comprising exposing a damaged DNA sample to conditions that promote substantial denaturation of damaged DNA in the damaged DNA sample, thereby forming a denatured damaged DNA sample, altering the conditions to conditions that do not promote substantial denaturation of damaged DNA in the damaged DNA sample to form a stabilized damaged DNA sample, incubating damaged DNA in the stabilized damaged DNA sample under conditions that promote replication of the damaged DNA, wherein replication of the damaged DNA results in a longer average fragment length for the replicated damaged DNA than the average fragment length in the damaged DNA sample, wherein during replication at least one of the replicated strands is displaced by strand displacement replication of another replicated strand.

Disclosed are methods, wherein the damaged DNA sample, the denatured damaged DNA sample, or both are exposed to ionic conditions, wherein the damaged DNA sample and denatured damaged DNA sample are exposed to ionic conditions by mixing an ionic solution with the damaged DNA sample, wherein the ionic solution is mixed with the damaged DNA sample prior to or during exposure of the damaged DNA sample to conditions that promote substantial denaturation of the damaged DNA, wherein the ionic solution is a salt solution, wherein the salt solution comprises one or more salts, wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination, wherein the Tris-HCl is from pH 7.0 to 8.0, wherein the salt is Tris-EDTA, wherein the salt solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA, wherein the ionic solution is diluted 2 to 5 fold when mixed with the damaged DNA sample, wherein the denatured damaged DNA sample is exposed to ionic conditions by mixing an ionic solution with the denatured

damaged DNA sample, wherein the ionic solution is mixed with the denatured
damaged DNA sample prior to or during altering of the conditions, wherein the
damaged DNA sample is exposed to conditions that promote substantial denaturation
by mixing the damaged DNA sample with a denaturing solution and by heating the
5 damaged DNA sample to a temperature and for a length of time that substantially
denatures the damaged DNA in the damaged DNA sample, wherein the damaged DNA
sample is mixed with the denaturing solution after the DNA sample is heated, wherein
the damaged DNA sample is mixed with the denaturing solution before the DNA
sample is heated, wherein the damaged DNA sample is mixed with the denaturing
10 solution at the same time the DNA sample is heated, wherein the damaged DNA
sample is mixed with the denaturing solution during heating of the DNA sample,
wherein the damaged DNA sample is mixed with the denaturing solution when heating
of the DNA sample begins, and/or wherein mixing the damaged DNA sample with a
denaturing solution produces alkaline conditions in the damaged DNA sample

15 Disclosed are methods wherein the denaturing solution comprises a base,
wherein the base is an aqueous base, wherein the base is sodium hydroxide, potassium
hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium
hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium
bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine,
20 diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine,
methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine,
aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide,
barium hydroxide, or DBU (1,8-diazobicyclo[5,4,0]undec-7-ene), wherein the base is
sodium hydroxide, wherein the denaturing solution comprises about 150 mM to about 1
25 M NaOH.

Disclosed are methods, wherein the denaturing solution is 10X concentration,
wherein the damaged DNA sample is mixed with the denaturing solution to create a 1X
concentration, and/or wherein the alkaline conditions comprise 15 to 50 mM NaOH.

Disclosed are methods, wherein the damaged DNA in the damaged DNA
30 sample is substantially denatured without further damaging the DNA, wherein the
damaged DNA sample is heated to a temperature of about 70°C or less and for a length
of time of about 5 minutes or less, wherein the temperature is about 60°C to about

70°C, wherein the temperature is about 50°C to about 60°C, wherein the temperature is about 40°C to about 50°C, wherein the temperature is about 25°C to about 40°C, wherein the temperature is about 60°C to about 70°C and the length of time is about 3 minutes, and/or wherein the temperature is about 25°C to about 50°C and the length of
5 time is about 5 minutes or more.

Disclosed are methods wherein altering the conditions comprises reducing the pH of and cooling the denatured damaged DNA sample, wherein the temperature to which the damaged DNA sample is heated is maintained during reduction of the pH of the denatured damaged DNA sample, wherein the temperature to which the damaged
10 DNA sample is heated is reduced before reduction of the pH of the denatured damaged DNA sample, and/or wherein the temperature to which the damaged DNA sample is heated is reduced during reduction of the pH of the denatured damaged DNA sample.

Disclosed are methods, wherein cooling the denatured damaged DNA sample is commenced during reduction of the pH of the denatured damaged DNA sample,
15 wherein cooling the denatured damaged DNA sample is commenced when the pH of the denatured damaged DNA sample is reduced.

Disclosed are methods, wherein the pH of the denatured damaged DNA sample is reduced by mixing the denatured damaged DNA sample with a stabilization solution.

Disclosed are methods, wherein the pH of the denatured damaged DNA sample
20 is reduced to the range of about pH 7.5 to about pH 8.0.

Disclosed are methods, wherein the pH of the denatured damaged DNA sample is reduced by mixing the denatured damaged DNA sample with a stabilization solution.

Disclosed are methods, wherein the stabilization solution comprises a buffer, and/or wherein the buffer is phosphate buffer, Good buffer, BES, BICINE, CAPS,
25 EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, TRICINE, sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, or Bis-tris propane.

Disclosed are methods, wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

30 Disclosed are methods, wherein the stabilization solution comprises an acid.

Disclosed are methods, wherein the acid is hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric

acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, or a carboxylic acid.

Disclosed are methods, wherein the carboxylic acid is ethanoic, propanoic, or butanoic.

5 Disclosed are methods, wherein the stabilization solution comprises a plurality of acidic agents.

Disclosed are methods, wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 9.0 to about pH 6.8, about pH 9.0 to about pH 7.5, about pH 9.0 to about pH 8.0, about pH 8.5 to about pH 6.8, about pH 8.5 to about pH 7.5, about pH 8.5 to about pH 8.0, about pH 8.0 to about pH 6.8, about pH 8.0 to about pH 7.5.

Disclosed are methods, wherein the pH of the denatured damaged DNA sample is reduced to about pH 9.0 or less, about pH 8.5 or less, about pH 8.0 or less, about pH 7.5 or less.

15 Disclosed are methods, wherein the stabilization solution comprises one or more salts.

Disclosed are methods, wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination.

20 Disclosed are methods, wherein the Tris-HCl is from pH 7.0 to 8.0.

Disclosed are methods, wherein the salt is Tris-EDTA.

Disclosed are methods, wherein the stabilization solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA.

25 Disclosed are methods, wherein the damaged DNA mixture is cooled at a rate of about 1°C per minute or less.

Disclosed are methods, wherein the damaged DNA mixture is cooled at a rate of about 1% per minute or less.

Disclosed are methods, wherein the damaged DNA mixture is cooled to room temperature or lower from 60°C to 70°C, from 50 to 60°C, from 40°C to 50°C, 30°C to 40°C, 50°C to 70°C.

30 Disclosed are methods, wherein the denaturing solution comprises one or more salts.

Disclosed are methods, wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination.

Disclosed are methods, wherein the Tris-HCl is from pH 7.0 to 8.0.

5 Disclosed are methods, wherein the salt is Tris-EDTA.

Disclosed are methods, wherein the denaturing solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA.

Disclosed are methods, wherein the damaged DNA sample is comprised of degraded DNA fragments of genomic DNA.

10 Disclosed are methods, wherein replication and repair of the damaged DNA is accomplished by incubating the damaged DNA in the presence of a DNA polymerase.

Disclosed are methods, wherein the polymerase is a DNA polymerase that can extend the 3'-ends of the damaged DNA.

Disclosed are methods, wherein the DNA polymerase is ϕ 29 DNA polymerase, 15 BST DNA polymerase, Taq DNA polymerase, a modified form of Taq DNA polymerase, a Reverse Transcriptase, T4 DNA polymerase, T7 DNA polymerase, Pol I DNA polymerase, or a modified form of DNA Polymerase I.

Disclosed are methods, wherein the DNA polymerase is ϕ 29 DNA Polymerase.

Disclosed are methods, wherein the damaged DNA is amplified using a kit, 20 wherein the kit comprises a denaturing solution, a stabilization solution, a set of primers, and a DNA polymerase.

Disclosed are methods, wherein the damaged DNA sample is a cell lysate, wherein the cell lysate is produced by exposing cells to alkaline condition, wherein the cell lysate comprises a whole genome,.

25 Disclosed are methods, comprising following exposure of the cells to alkaline conditions, exposing a first portion of the cell lysate to conditions that promote substantial denaturation of damaged DNA in the first portion of the cell lysate, wherein reducing the pH of the cell lysate comprises reducing the pH of the first portion of the cell lysate to form a first stabilized cell lysate and reducing the pH of a second portion 30 of the cell lysate to form a second stabilized cell lysate, following reducing the pH of the cell lysate, mixing the second stabilized cell lysate with the first stabilized cell lysate under conditions that promote transient denaturation of the ends of damaged

DNA in the second stabilized cell lysate and that maintain substantial denaturation of the damaged DNA in the first stabilized cell lysate, thereby forming a stabilized cell lysate mixture, and prior to incubating the stabilized cell lysate, cooling the stabilized cell lysate mixture under conditions that promote annealing of the ends of the
5 transiently denatured damaged DNA to the substantially denatured damaged DNA, wherein incubating the stabilized cell lysate under conditions that promote replication of the genome also promotes replication of the damaged DNA, wherein the annealed ends of the damaged DNA prime replication, wherein replication of the damaged DNA results in repair of the replicated strands.

10 Disclosed are methods of amplifying damaged DNA, the method comprising exposing a first damaged DNA sample to conditions that promote substantial denaturation of damaged DNA in the first damaged DNA sample, thereby forming a denatured damaged DNA sample, reducing the pH of the denatured damaged DNA sample to form a stabilized denatured damaged DNA sample, mixing a second
15 damaged DNA sample with the stabilized denatured damaged DNA sample under conditions that promote transient denaturation of the ends of damaged DNA in the second sample and that maintain substantial denaturation of the damaged DNA in the stabilized denatured damaged DNA sample, thereby forming a damaged DNA mixture, cooling the damaged DNA mixture under conditions that promote annealing of the ends
20 of the transiently denatured damaged DNA to the substantially denatured damaged DNA, incubating the annealed damaged DNA under conditions that promote replication of the damaged DNA, wherein the annealed ends of the damaged DNA prime replication, wherein replication of the damaged DNA results in repair of the replicated strands, wherein during replication at least one of the replicated strands is
25 displaced by strand displacement replication of another replicated strand.

Disclosed are methods, wherein the temperature to which the first damaged DNA sample is heated is maintained during mixing of the second damaged DNA sample with the stabilized denatured damaged DNA sample.

Disclosed are methods, wherein the pH of the stabilized denatured damaged
30 DNA sample is not high enough nor low enough to cause further substantial denaturation upon mixing second damaged DNA sample with the stabilized denatured damaged DNA sample.

Disclosed are methods, wherein the first damaged DNA sample is a portion of a damaged DNA sample, wherein the second damaged DNA sample is a portion of the same damaged DNA sample.

Disclosed are methods, wherein the first damaged DNA sample is from the same source as the second damaged DNA sample.

Disclosed are methods, wherein the first damaged DNA sample is from the same organism as the second damaged DNA sample.

Disclosed are methods, wherein the first damaged DNA sample is from the same tissue as the second damaged DNA sample.

Disclosed are methods, wherein the second damaged DNA sample is mixed with the stabilized denatured damaged DNA sample at a temperature and for a length of time that transiently denatures the damaged DNA in the second damaged DNA sample.

Disclosed are methods, wherein the temperature is about 70°C or less and the length of time is about 30 seconds or less.

Disclosed are methods, wherein the second damaged DNA sample is mixed with the stabilized denatured damaged DNA sample at a temperature that does not further damaging the DNA.

Disclosed are methods, wherein the temperature is about 60°C to about 70°C, about 50°C to about 60°C, about 40°C to about 50°C, about 25°C to about 40°C, about 25°C to about 70°C. Disclosed are methods wherein the length of time is about 30 seconds.

Examples

A. Example 1: Whole Genome Amplification using Nuclease Resistant Hexamer Primers

This example describes a demonstration of an embodiment of the disclosed method and analysis and comparison of the results. The exemplified method is the disclosed multiple displacement amplification form of whole genome amplification using nuclease resistant random hexamer primers. Some reactions in this example were performed without subjecting the sample to denaturing conditions, a preferred form of the disclosed method. In other reactions, the template DNA was subjected to denaturation prior to amplification. MDA was performed using ϕ 29 DNA polymerase.

1. Materials and Methods

i. DNA and Enzymes.

A panel of human genomic DNA samples, the Human Variation Panel-Caucasian Panel of 100 (reference number HD100CAU) was obtained from Coriell
5 Cell Repositories. Human genomic DNA was also obtained from Promega Corp. Thiophosphate-modified random hexamer (5'-NpNpNpNp^sNp^sN-3') was synthesized at Molecular Staging, ϕ 29 DNA polymerase was from Amersham Pharmacia Biotech, and yeast pyrophosphatase was from Boehringer-Mannheim. DNA size markers (100 bp DNA ladder, 1 kb DNA ladder) were from Gibco BRL.

10 ii. Amplification of Human Genomic DNA.

Human genomic DNA (300 ng to 0.03 ng, as indicated) was placed into 0.2 ml tubes in a total volume of 50 μ l, yielding final concentrations of 25 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, and 100 μ M exonuclease-resistant hexamer. A heat-treatment step (that is, exposure to denaturing conditions) to increase primer annealing
15 was included or omitted, as indicated, for individual experiments. Annealing reactions were heated to 95°C for 3 minutes and chilled to 4°C in a PCR System Thermocycler (Perkin Elmer). Reactions were then brought to a final volume of 100 μ l, containing final concentrations of 37 mM Tris-HCl, pH 8.0, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1.0 mM dNTPs, 1 unit/ml of yeast pyrophosphatase, 50 μ M exonuclease-
20 resistant hexamer, and 800 units/ml ϕ 29 DNA polymerase. Radioactively labeled α -[³²P] dCTP, approximately 60 cpm/pmol total dNTPs, was added as indicated. Reactions were incubated for 18 hours at 30°C. Incorporation of acid-precipitable radioactive deoxyribonucleotide product was determined with glass fiber filters. After the reactions were terminated, 3 μ l aliquots were cleaved with restriction endonuclease
25 AluI and analyzed by electrophoresis through a 1.0% agarose gel in Tris-borate-EDTA buffer, stained with GelStar (Molecular Probes) or SYBR Green (Molecular Probes), and imaged with a Storm 860 PhosphorImager (APB). Denaturing gel analysis was carried out by electrophoresis through a 1.0% agarose gel in 30 mM NaOH, 1 mM EDTA. The radioactive products in the dried gel were visualized with the Storm 860
30 PhosphorImager.

iii. Southern Analysis.

10 µg of whole genome amplified DNA or human genomic DNA controls were digested with EcoRI restriction endonuclease and separated through a 1% agarose gel in 1x TBE buffer. Standard Southern analysis procedure (Southern, Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol.* 98:503-517 (1975)) was performed using a Hybond-N+ membrane (Amersham Pharmacia Biotech, Piscataway, NJ). An exon fragment probe of parathyroid hormone (p20.36) and RFLP marker probes for the D13S12 (p9D11) and Thyroglobulin (pCHT.16/8) loci were obtained from American Type Culture Collection. Probes were radiolabeled using the NEBlot random primer labeling method (New England Biolabs, Beverly, MA). The membrane was prehybridized for 1 hr and hybridized to the radiolabeled probe overnight in a Membrane Hybridization Buffer (Amersham Pharmacia Biotech, Piscataway, NJ). The hybridized membrane was washed in 2x SSC and 0.1% SDS twice for 5 min at room temperature, 1x SSC and 0.1%SDS for 15 min at 42°C, and 0.1x SSC twice for 15 min at 65°C. The membrane was then exposed overnight and analyzed using the Storm 860 PhosphorImager.

iv. Quantitative PCR Analysis.

TaqMan analysis was performed using the ABI 7700 according to the manufacturer's specifications (Applied Biosystems, Foster City, CA) using 1 µg of amplified DNA as template. TaqMan assay reagents for the 8 loci tested were obtained from ABI. The 8 loci and their chromosome assignments were, acidic ribosomal protein (1p36.13); connexin 40 (1q21.1); chemokine (C-C motif) receptor 1 (3p21); chemokine (C-C motif) receptor 6 (6q27); chemokine (C-C) receptor 7 (17q21); CXCR5 Burkitt lymphoma receptor 1 (chr. 11); c-Jun (1p32-p31); and MKP1 dual specificity phosphatase 1 (5q34). Connexin 40 is located near the centromere and chemokine (C-C motif) receptor 6 is located near the telomere. A standard curve for input template was generated to determine the loci copy number in amplified DNA relative to that of genomic DNA. The standard curve was generated from 0, 0.001, 0.01, 0.1, 0.5, and 1 µg of genomic DNA.

v. Amplification of Human Genomic DNA by Degenerate Oligonucleotide PCR (DOP-PCR).

Human genomic DNA (ranging from 300 ng to 0.03 ng) was amplified as described (Telenius *et al.*, *Genomics*. 13:718-725 (1992)). Radioactively labeled α -
5 [32P] dCTP, approximately 60 cpm/pmol total dNTPs, was added to the reaction for the quantitation of PCR product yield. Taq DNA Polymerase was from Invitrogen Life Technologies, Carlsbad, CA. DOP-PCR amplifications were carried out using a GeneAmp 9700 PCR System thermocycler (Applied Biosystems, Foster City, CA). Locus representation in the DOP-PCR product was quantitatively analysed using the
10 TaqMan assay (Invitrogen Life Technologies, Carlsbad, CA).

vi. Amplification of Human Genomic DNA by Primer Extension Preamplification (PEP).

Human genomic DNA (ranging from 300 ng to 0.03 ng) was placed into 0.2 ml tubes in a total volume of 60 μ l, yielding final concentrations of 33 μ M PEP random
15 primer (5'-NNN NNN NNN NNN NNN-3') as described (Zhang *et al.*, Whole genome amplification from a single cell: implications for genetic analysis. *Proc Natl Acad Sci U S A*. 89:5847-5851 (1992)). Radioactively labeled α -[32P] dCTP, approximately 60 cpm/pmol total dNTPs, was added to the reaction for the quantitation of PCR product. PEP reactions were carried out using a GeneAmp 9700 PCR System thermocycler
20 (Applied Biosystems, Foster City, CA). Locus representation in the PEP product was quantitatively analysed using the TaqMan assay (Invitrogen Life Technologies, Carlsbad, CA).

vii. Genotyping of Single Nucleotide Polymorphisms.

The SNPs analyzed here had the following chromosomal locations; 1822, 251,
25 and 221, 13q32; 465, 458, and 474, 19q13; VCAM, 1p31; IL-8, 4q13; PDK2-2, 17p; SNP21, not known. Assays were carried out as described by Faruqi *et al.*, High-Throughput Genotyping of Single Nucleotide Polymorphisms with Rolling Circle Amplification. *BMC Genomics*, 2:4 (2001). Briefly, DNA denaturation, annealing and ligation reactions were carried out in an Eppendorf Master Cycler (Eppendorf
30 Scientific, Germany). Exponential RCA reactions were performed in the Real-Time ABI 7700 Sequence Detector (Perkin Elmer). Two controls lacking ligase were also carried out for each SNP, confirming the specificity of the assays. The DNA samples

were digested with the restriction endonuclease AluI before being used as template in the ligation reaction. Ligation reactions were set up in 96-well MicroAmp Optical plates (Perkin Elmer) in a 10 μ l reaction volume containing 1 unit Ampligase (Epicentre Technologies), 20 mM Tris-HCl (pH 8.3), 25 mM KCl, 10 mM MgCl₂, 0.5 mM NAD, and 0.01% Triton® X-100. Standard reactions contained 0.5 pM open circle padlock and 100 ng of Alu I digested genomic DNA. DNA was denatured by heating the reactions at 95°C for 3 min followed by annealing and ligation at 60°C for 20 min. 20 μ l of ERCA mix was added to the 10 μ l ligation reaction. The ERCA mix contained 5% TMA oxalate, 400 μ M dNTP mix, 1 μ M each of the two primers, 8 units of Bst polymerase (New England Biolabs, MA), and 1X modified ThermoPol II reaction buffer containing 20 mM Tris-HCl (pH 8.8), 10 mM KCl, 10 mM (NH₄)₂SO₄ and 0.1% Triton® X-100.

viii. Comparative Genome Hybridization.

Genomic DNA preparations were nick-translated to incorporate nucleotides modified with biotin for amplified samples or digoxigenin for unamplified control samples. Equimolar amounts of amplified and unamplified DNA were co-hybridized in the presence or absence of CotI DNA to suppress repetitive DNA cross-hybridization. Specific hybridization signals were detected by avidin-FITC and anti-digoxigenin rhodamine. Captured images of metaphase chromosomes were analyzed using the Applied Imaging CGH software program and fluorescence profiles were generated. As controls, differentially labeled amplified or unamplified DNAs were mixed, hybridized, detected and subjected to ratio analysis as outlined above.

2. Results

Using the embodiment of the disclosed method, 30 pg (approximately 9 genomic copies) of human genomic DNA was amplified to approximately 30 μ g within 4 hours. The average fragment length was greater than 10 kb. The amplified human DNA exhibited normal representation for 10 single nucleotide polymorphisms (SNPs). Maximum bias among 8 chromosomal loci was less than 3-fold in contrast to four to six orders of magnitude for PCR-based WGA methods. Human DNA amplified with the disclosed method is useful for several common methods of genetic analysis, including genotyping of single nucleotide polymorphisms (SNP), chromosome painting, Southern blotting and RFLP analysis, subcloning, and DNA sequencing.

It has been discovered that the use of random hexamer primers and ϕ 29 DNA polymerase in multiple displacement amplification, a cascading, strand displacement reaction (U.S. Patent No. 6,124,120 to Lizardi), will readily amplify linear, human genomic DNA. Amplification of genomic DNA by the disclosed form of MDA at
5 30°C is exponential for 4-6 hours. The effect of template concentration on amplification yield in the disclosed method is shown in Figure 1. 100 fg to 10 ng human genomic DNA was denatured at 95°C for 5 min, and then MDA was carried out at 30°C as described above. Aliquots were taken at the times indicated in Figure 1 to quantitate DNA synthesis. Amplification reactions (100 μ l) yielded approximately 25-
10 30 μ g DNA product regardless of the starting amount of genomic DNA over a 5-log range (100 fg to 10 ng; Figure 1). For some applications, this allows subsequent genetic analysis without the need to measure or adjust DNA concentration.

The products of the MDA reaction were characterized as follows. Radioactively labeled human genome amplification samples (0.6 μ g) were
15 electrophoresed through an alkaline agarose gel (1%, Tris-Borate EDTA buffer), stained, and imaged as described above. Average product length exceeded 10 kb.

To examine the integrity of amplified DNA, a restriction fragment within the human parathyroid hormone gene (chromosome 11p15.2-15.1) was analyzed on Southern blots. A 1.9 kb restriction enzyme fragment was observed for MDA-based
20 WGA products amplified from as few as 10 genomic copies (or 10^6 -fold amplification). These MDA reactions included a heat denaturation step and amplification was carried out as described above. EcoRI DNA digests were probed using a radioactively-labeled genomic fragment of the parathyroid hormone gene (p20.36) that hybridized to an approximately 1.9 kb DNA fragment. The EcoRI-cleaved DNA preparations were
25 genomic DNA, DNA amplified by MDA from varying amounts of input genomic DNA, or an MDA reaction that lacked input genomic DNA template. This demonstrates that the products of MDA are long enough to yield specific DNA fragments several kb in length after cleavage by restriction endonucleases.

While PCR-based WGA methods typically generate products of only several
30 hundred nucleotides in length (Telenius *et al.*, *Genomics*. 13:718-725 (1992); Cheung and Nelson, *Proc Natl Acad Sci U S A*. 93:14676-14679 (1996); Zhang *et al.*, *Proc Natl Acad Sci U S A*. 89:5847-5851 (1992)), products of the disclosed method were of

sufficient length and integrity for RFLP-based genotyping: 16 random individuals were correctly genotyped by the presence of a 2.1 kb, and 1.1 kb Pst I fragment.

Specifically, PstI DNA digests were probed using a radioactively-labeled genomic fragment of the RFLP marker D13S12 locus (p9D11) that hybridized to an invariant 3.8 kb DNA fragment and a polymorphic 2.1 kb (allele A) or 1.1 kb (allele B) DNA fragment. The PstI-cleaved DNA preparations were genomic DNA and 5 different patient DNAs amplified by MDA without any heat denaturation of the DNA template (10,000x amplification).

Omission of denaturation conditions prior to MDA was useful for detection of restriction fragments greater than 5 kb in length. MDA reactions were performed with or without heat denaturation of the genomic target DNA heterozygous for two thyroglobulin alleles. Amplification was carried out as described above. TaqI DNA digests were probed using a radioactively-labeled genomic fragment of the thyroglobulin gene (pCHT.16/8) that hybridized to invariant 1 kb and 3 kb DNA fragments and a polymorphic 5.8 kb (allele A) or 5.2 kb (allele B) DNA fragment. The TaqI-cleaved DNA preparations were genomic DNA, DNA amplified by MDA reaction (10,000x) with a 95°C preheating step, and an MDA reaction (10,000x) without the preheating step. MDA without heat denaturation gave a good yield of DNA fragments 5.2 and 5.8 kb in size, while neither of the 5.2 or 5.8 kb DNA fragments were visible using MDA with heat denaturation of the template.

The most useful results from whole genome amplification are obtained when the amplification provides complete coverage of genomic sequences and minimal amplification bias. It is also preferred that the amplification product perform similarly to unamplified genomic DNA during subsequent genetic analysis. Genome coverage after MDA with heat denaturation of the template was examined for 10 randomly distributed SNPs after amplifications of 100-, 10,000-, or 100,000- fold. The presence of all loci was confirmed in the amplified DNA with the exception of one locus (PDK2-2) in 100,000-fold amplified DNA (Table 1). MDA DNA from 72 individuals was genotyped for one of these SNPs. Following 100-fold amplification by MDA, genotyping accuracy was 97% (70 of 72 genotypes scored correctly, Table 1, SNP 1822), a result that was indistinguishable from unamplified genomic DNA genotyped by the same method. MDA-based WGA thus offers an attractive alternative to multiple

locus-specific PCR preamplifications for large SNP scoring studies, especially where sample availability is limited.

Table 1

SNP designation	Fold whole genome amplification		
	100X	10,000X	100,000X
	Correct SNP calls/total assays		
1822	70/72	4/4	3/4
251	8/8	4/4	4/4
221	3/4	3/4	4/4
465	4/4	3/4	3/4
458	4/4	3/4	3/4
474	4/4	3/4	4/4
VCAM	4/4	4/4	4/4
IL-8	4/4	4/4	3/4
SNP21	4/4	4/4	3/4
PDK2-2	4/4	4/4	0/12

5 Sequence bias can occur in amplification methods, and may result from factors such as priming efficiency, template accessibility, GC content, and proximity to telomeres and centromeres. Amplification bias of the disclosed MDA method was examined by TaqMan quantitative PCR for 8 genes, including one near the centromere of chromosome 1 (connexin 40) and one adjacent to the telomere of chromosome 6
10 (chemokine (C-C motif) receptor 6). For 100, 1,000, and 10,000-fold MDA without heat denaturation, the maximum amplification biases were only 2.7, 2.3, and 2.8 respectively, expressed as the ratio of the most highly represented gene to the least represented gene. In contrast, two WGA methods based on PCR, DOP-PCR (Telenius *et al.*, *Genomics*. 13:718-725 (1992); Cheung and Nelson, *Proc Natl Acad Sci U S A*. 93:14676-14679 (1996)) and PEP (Zhang *et al.*, *Proc Natl Acad Sci U S A*. 89:5847-
15 5851 (1992)), exhibited amplification bias of 4-6 orders of magnitude. These values were consistent with literature values for the bias of PCR-mediated WGA methods;

PEP has been reported to generate an amplification bias of up to 50-fold even between two alleles of the same gene (Paunio et al., *Clin. Chem.* 42:1382-1390 (1996)).

Significantly, the 3-fold amplification bias of MDA remained almost constant between 100 and 100,000-fold amplification. An absence of significant sequence bias observed
5 in MDA-based WGA may be explained by the extraordinary processivity of Φ 29 DNA polymerase (Blanco et al., *J. Biol. Chem.* 264:8935-8940 (1989)). Tight binding of the polymerase to the template assures replication through obstacles caused by DNA primary or secondary structure.

Surprisingly, TaqMan quantification indicated that certain gene loci were
10 enriched by MDA in amplified DNA; locus representation was >100% of the representation in genomic DNA. Representation of mitochondrial DNA was the same between starting gDNA and amplified DNA. Southern blots and chromosome painting experiments indicated that repetitive sequences were underrepresented in MDA product, conferring an effective enrichment of genes. Thus, amplified DNA contained
15 between 100-300% copy numbers of 8 genes relative to genomic DNA. Additional studies will be necessary to identify the extent and type of repetitive element under-representation; one hypothesis for this observation is that primers corresponding to highly repetitive elements become depleted during MDA. However, in contrast to PCR-based WGA products, which contain up to 70% amplification artifacts (Cheung and Nelson, Whole genome amplification using a degenerate oligonucleotide primer allows hundreds of genotypes to be performed on less than one nanogram of genomic DNA. *Proc Natl Acad Sci U S A.* 93:14676-14679 (1996)), MDA-based WGA products
20 appear to be entirely derived from genomic sequences.

Whole genome MDA was also tested for uniformity of chromosome coverage
25 by comparative genomic hybridization. MDA-generated DNA was cohybridized to metaphase chromosomes with an equivalent amount of unamplified genomic DNA. Amplification reactions included a heat denaturation step and amplification was carried out as described above. Amplified and unamplified DNA samples were nick-translated to incorporate biotin nucleotide and digoxigenin nucleotide, respectively. Specific
30 signals were detected by avidin FITC and anti-digoxigenin rhodamine. With *CotI* suppression, the hybridization patterns of MDA probes and unamplified probes were indistinguishable even after 100,000-fold amplification. Without *CotI* suppression,

however, the 100,000-fold amplified probe gave reduced centromeric signals, indicating some loss of repetitive centromeric sequences. The uniformity of signal along the length of the chromosome arms was further evidence that MDA-based WGA does not induce significant amplification bias. These results indicated that MDA compared favorably with DOP PCR for preparation of chromosome painting probes (Kim et al., Whole genome amplification and molecular genetic analysis of DNA from paraffin-embedded prostate adenocarcinoma tumor tissue. *J Urol.* 162:1512-1518 (1999); Klein et al., Comparative genomic hybridization, loss of heterozygosity, and DNA sequence analysis of single cells. *Proc Natl Acad Sci U S A* 96:4494-4499 (1999); Wells et al., Detailed chromosomal and molecular genetic analysis of single cells by whole genome amplification and comparative genomic hybridization. *Nucleic Acids Res* 27:1214-1218 (1999)), but that unlike the latter, suppression hybridization may be unnecessary for detection of single copy sequences. This method should be invaluable for DNA probe preparation for comparative genome hybridization, karyotyping and chip-based genetic analysis from limited patient DNA sources such as needle biopsy material or amniocentesis samples.

For genome subcloning and sequencing MDA appears to have several intrinsic advantages over PCR-based methods. Product size of >10kb is compatible with genome subcloning. Since no in vivo propagation of amplified material is necessary, MDA may represent an efficient method for amplifying “poisonous” genomic sequences. In addition, ϕ 29 DNA polymerase used for MDA has an error rate of 1 in $10^6 - 10^7$ (Esteban et al., Fidelity of Phi29 DNA polymerase. Comparison between protein-primed initiation and DNA polymerization. *J. Biol. Chem.* 268:2719-2726 (1993)) in contrast to approximately 3 in 10^4 for PCR with Taq DNA polymerase (Eckert and Kunkel, DNA polymerase fidelity and the polymerase chain reaction. *PCR Methods and Applications.* 1:17-24 (1991)). Therefore, PCR accumulates about one mutation per 900 bases (Saiki et al., Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239:487-491 (1988)) after 20 cycles. Sequence error rates of cloned MDA-based WGA products appear to be similar to those for cloned genomic DNA. Furthermore, minimal amplification bias, uniform yields, and assay simplicity make MDA amenable to the automated, high-density microwell formats used for genome subcloning and sequencing.

In summary, the utility of MDA-based WGA was demonstrated for a variety of uses including quantitative PCR, SNP genotyping, Southern blot analysis of restriction fragments, and chromosome painting. Several situations may be contemplated where MDA may represent the method of choice for WGA: Firstly, applications where faithful replication during WGA is necessary, such as molecular cloning, or single cell analysis; Secondly, applications where adequate genome representation during WGA is critical, such as genome-wide SNP genotyping studies; And thirdly, where minimization of bias during WGA is important, particularly cytogenetic testing such as pre-natal diagnosis (Harper and Wells, Recent advances and future developments in PGD. *Prenat Diagn.* 19:1193-1199 (1999)), comparative genome hybridization (Wells and Delhanty, Comprehensive chromosomal analysis of human preimplantation embryos using whole genome amplification and single cell comparative genomic hybridization. *Mol Hum Reprod.* 6:1055-1062 (2000)), and assessment of loss of heterozygosity (Paulson et al., Loss of heterozygosity analysis using whole genome amplification, cell sorting, and fluorescence-based PCR. *Genome Res.* 9:482-491 (1999)). Additional situations where there is a significant need for WGA include amplification of DNA from micro-dissected tissues (Kim et al., Whole genome amplification and molecular genetic analysis of DNA from paraffin-embedded prostate adenocarcinoma tumor tissue. *J Urol.* 162:1512-1518 (1999)), buccal smears (Gillespie et al., HLA class II typing of whole genome amplified mouth swab DNA. *Tissue Antigens.* 56:530-538 (2000)), and archival, anthropological samples (Buchanan et al., Long DOP-PCR of rare archival anthropological samples. *Hum Biol.* 72:911-925 (2000)). Finally, DNA amplified from sorted, individual chromosomes may be used for the generation of whole chromosome-specific painting probes (Guillier-Gencik et al., Generation of whole-chromosome painting probes specific to each chicken macrochromosome. *Cytogenet Cell Genet.* 87:282-285 (1999)).

B. Example 2: Increasing Time of Incubation at 95°C Causes Increasing Template DNA Strand Breakage.

This example demonstrates that significant template DNA strand breakage is generated by incubation at 95°C (which is used in typical amplification reactions to denature the DNA), and that strand breakage is reduced by decreasing the duration of heat treatment. As with most nucleic acid amplification techniques, the integrity of the

starting DNA template can have an important effect on the rate and yield of the amplified product. In reactions where the nucleic acid to be amplified is degraded, the yield of amplified product may be reduced both in quality and quantity. This example demonstrates the reduction of template DNA strand breakage by decreasing time of incubation at 95°C.

Six reactions were carried out under the conditions used for DNA template strand separation with heat-denaturation in order to illustrate the degradation of template DNA. Human genomic DNA (10 µg) was placed into a 0.2 ml tube in a total volume of 50 µl, yielding final concentrations of 25 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂. Reactions were heated to 95°C in a PCR System Thermocycler (Perkin Elmer), and an aliquot of 8 µl was taken and put on ice at indicated times. For each time point, a 2 µl aliquot was analyzed by electrophoresis through a 0.8% alkaline agarose gel (30 mM NaOH, 1 mM EDTA). After electrophoresis, the gel was neutralized with 1X TBE, stained with SYBR Green II (Molecular Probes), and imaged with a Storm 860 PhosphorImager (APB). The total amount of DNA imaged was determined for each lane of the gel and the fragment size at which 50% of the DNA was larger, and 50% was smaller, was determined for the samples drawn at each time point. The results are shown in Figure 2.

As can be seen, significant breakage of template DNA occurs after incubation at 95°C for longer than three minutes. Such DNA breakage is substantially reduced when incubation is limited to one minute.

C. Example 3: Increasing Time of DNA Template Incubation at 95°C Causes Decreased Rate and Yield of DNA Amplification.

This example demonstrates that increased time of template DNA incubation at 95°C (which is used in typical amplification reactions to denature the DNA) causes a reduction in both the rate and the yield with which DNA is amplified by MDA. Omission of DNA template incubation at 95°C results in the greatest rate and yield of DNA amplification.

Six MDA reactions were carried out using template DNA treated at 95°C under the conditions described in Example 2. Purified human genomic DNA (3 ng) was placed into 0.2 ml tubes in a total volume of 50 µl, containing 25 mM Tris-HCl, pH 7.5, 50 mM KCl, and 10 mM MgCl₂. Reactions were heated to 95°C for the time

indicated and chilled to 4°C in a PCR System Thermocycler (Perkin Elmer). These reactions were then brought to a final volume of 100 µl, containing final concentrations of 37 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, 50 µM exonuclease-resistant hexamer, 5 mM (NH₄)₂SO₄, 1.0 mM dNTPs, 1 unit/ml of yeast pyrophosphatase, and 800 units/ml φ29 DNA polymerase. Radioactively labeled α-[³²P] dCTP, approximately 60 cpm/pmol total dNTPs, was added and reactions were incubated for 22 hours at 30°C. Aliquots were taken at the times indicated and incorporation of acid-precipitable radioactive deoxyribonucleotide product was determined with glass fiber filters. The results are shown in Figure 3.

As can be seen, omission of heat treatment of the DNA template results in the optimal rate and yield of DNA synthesis (see 0 min curve). Increasing duration of DNA template heat treatment resulted in progressively reduced rate and yield of DNA synthesis (see 1 min, 3 min, 5 min, 10 min, and 20 min curves).

D. Example 4: Increasing Time of DNA Template Incubation at 95°C Results in Decreasing DNA Product Strand Size.

This example demonstrates that increased time of template DNA incubation at 95°C (which is used in typical amplification reactions to denature the DNA) causes a reduction in the length of the DNA products amplified by MDA. Omission of DNA template incubation at 95°C results in the greatest size of DNA amplification products.

Six MDA reactions were carried out using template DNA treated at 95°C under the conditions described in Example 3. Radioactively labeled α-[³²P] dCTP was added, approximately 60 cpm/pmol total dNTPs. Reactions were incubated for 22 hours at 30°C and aliquots were taken at the times indicated. For each time point, a 2 µl aliquot was analyzed by electrophoresis through a 0.8% alkaline agarose gel (30 mM NaOH, 1 mM EDTA). After electrophoresis, the gel was neutralized and imaged as described in Example 2. The total amount of DNA imaged was determined for each lane of the gel and the fragment size at which 50% of the DNA was larger, and 50% was smaller, was determined for the samples from each time point. The results are shown in Figure 4.

As can be seen, omission of heat treatment of the DNA template results in the synthesis of the largest DNA products. Increasing duration of DNA template heat treatment resulted in progressively reduced DNA product size.

E. Example 5: Omission of DNA Template Incubation at 95°C Results in Increased Locus Representation in DNA Products Amplified by MDA.

This example demonstrates that omission of template DNA incubation at 95°C (which is used in typical amplification reactions to denature the DNA) results in no loss
5 in the representation of eight randomly selected loci in DNA products amplified by MDA. Omission of DNA template incubation at 95°C actually results in an increase in locus representation of DNA amplification products relative to template genomic DNA.

Two MDA reactions were carried out using template DNA either treated or not treated at 95°C. Purified human genomic DNA (3 ng) was placed into a 0.2 ml tube in
10 a total volume of 50 µl, containing 25 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, and 100 µM exonuclease-resistant hexamer. The annealing reaction was heated to 95°C for 3 minutes and chilled to 4°C in a PCR System Thermocycler (Perkin Elmer). The reaction was then brought to a final volume of 100 µl, containing final concentrations of 37 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, 50 µM
15 exonuclease-resistant hexamer, 5 mM (NH₄)₂SO₄, 1.0 mM dNTPs, 1 unit/ml of yeast pyrophosphatase, and 800 units/ml φ29 DNA polymerase. For amplification lacking the heat denaturation step, DNA template (3 ng) was placed directly into a 0.2 ml tube in a total volume of 100 µl containing 37 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, and 50 µM exonuclease-resistant hexamer, 5 mM (NH₄)₂SO₄, 1.0 mM dNTPs,
20 1 unit/ml of yeast pyrophosphatase, and 800 units/ml φ29 DNA polymerase. Reactions were incubated for 18 hours at 30°C.

TaqMan® quantitative PCR analysis was performed using the ABI 7700 according to the manufacturer's specifications (Applied Biosystems, Foster City, CA) using 1 µg of MDA-amplified DNA as template. TaqMan® assay reagents for the 8
25 loci tested were obtained from ABI. The 8 loci and their chromosome assignments were, acidic ribosomal protein (1p36.13); connexin 40 (1q21.1); c-Jun (1p32-p31); MKP1 dual specificity phosphatase 1 (5q34); chemokine (C-C motif) receptor 7 (17q21); chemokine (C-C motif) receptor 1 (3p21); CXCR5 Burkitt lymphoma receptor 1 (chr. 11); and chemokine (C-C motif) receptor 6 (6q27). Connexin 40 is located near
30 the centromere and chemokine (C-C motif) receptor 6 is located near the telomere. A standard curve for input template was generated to determine the loci copy number in

amplified DNA relative to that of genomic DNA. The standard curve was generated from 0, 0.001, 0.01, 0.1, 0.5, and 1 μ g of genomic DNA. The locus representation was expressed as a percent, relative to the locus representation in the input genomic DNA, and was calculated as the yield of quantitative PCR product from 1 μ g amplified DNA divided by the yield from 1 μ g genomic DNA control. The results are shown in Figure 5.

As can be seen, there is no reduction of locus representation in DNA amplified from template DNA without heat treatment at 95°C. However, significant loss of locus representation was observed from template DNA heat-denatured for 3 min at 95°C.

F. Example 6: Amplification Bias of Loci Amplified by MDA is Significantly Lower than Amplification Bias of DNA Amplified by PEP or DOP-PCR.

This example demonstrates that, for 100-, 1,000-, and 10,000-fold MDA-amplified DNA, omission of template DNA incubation at 95°C (which is used in typical amplification reactions to denature the DNA) results in low bias in the representation of eight randomly selected loci in DNA products. In contrast, two whole genome amplification (WGA) methods based on PCR, DOP-PCR and PEP, exhibit amplification biases of 2-6 orders of magnitude.

MDA reactions were carried out using template DNA not treated at 95°C as described in Example 5. Reactions (100 μ l) contained 300, 30, 3, or 0.3 ng DNA, resulting in approximately 100-, 1000-, 10,000-, and 100,000-fold DNA amplification.

Amplification of human genomic DNA by degenerate oligonucleotide PCR (DOP-PCR; Telenius *et al.*, *Genomics*. 13:718-725 (1992); Cheung and Nelson, *Proc Natl Acad Sci U S A*. 93:14676-14679 (1996)) was carried out as follows. Human genomic DNA (ranging from 300ng to 0.03ng) was placed into 0.2 ml tubes in a total volume of 50 μ l, yielding final concentrations of 2 μ M DOP Primer (5'-CCG ACT CGA GNN NNN NAT GTG G-3' (SEQ ID NO:20); N = A, G, C, or T in approximately equal proportions), 200 μ M dNTPs, 10 mM Tris Cl (pH 8.3), 0.005% (v/v) BRIJ 35, 1.5 mM MgCl₂, and 50 mM KCl. Radioactively labeled α -[³²P] dCTP, approximately 60 cpm/pmol total dNTPs was added to the reaction for quantitation of DNA synthesis as described for MDA product quantitation in Example 3. After the initial denaturation of the template at 95°C for 5 min, 2.5 units Taq DNA Polymerase

(Invitrogen Life Technologies, Carlsbad, CA) was added, followed by 5 cycles of 94°C for 1 min, 30°C for 1.5 min, ramping up to 72°C in 3 min and elongation at 72°C for 3 min, and then 35 cycles of 94°C for 1 min, 62°C for 1 min, and 72°C for 2 min (+14 extra seconds/ cycle). A final elongation was done at 72°C for 7 min. Amplification reactions were carried out using a GeneAmp 9700 PCR Systems Thermocycler (Applied Biosystems, Foster City, CA).

Amplification of human genomic DNA by primer extension preamplification (PEP; Zhang *et al.*, *Proc Natl Acad Sci U S A.* 89:5847-5851 (1992)) was carried out as follows. Human genomic DNA (ranging from 300ng to 0.03ng) was placed into 0.2 ml tubes in a total volume of 60 µl, yielding final concentrations of 33uM PEP random primer (5'-NNN NNN NNN NNN NNN-3'), 100uM dNTPs, 10 mM Tris-HCl, pH 8.3 (20°C), 1.5 mM MgCl₂, 50 mM KCl, and 5 units of Taq DNA Polymerase (Invitrogen Life Technologies, Carlsbad, CA). Radioactively labeled α-[³²P] dCTP, approximately 60 cpm/pmol total dNTPs was added to the reaction for quantitation of PCR product yield as described for MDA product quantitation in Example 3. The PEP reaction was performed for 50 cycles at 92°C for 1 min, 37°C for 2 min, ramping up to 55°C at 10 sec/degree, and elongation at 55°C for 4 min. PEP reactions were carried out using a GeneAmp 9700 PCR Systems Thermocycler (Applied Biosystems, Foster City, CA).

TaqMan® quantitative PCR analysis was performed as described in Example 5, and maximum amplification bias between loci was calculated by dividing the high locus representation value by the low value for each level of fold amplification. The results are shown in Figure 6.

The relative representation of eight loci is depicted in Figure 6 for amplification reactions carried out by three different WGA procedures. The X-axis represents the fold amplification in the amplified DNA used as template for quantitative PCR; the Y-axis is the locus representation, expressed as a percent, relative to input genomic DNA, which is calculated as the yield of quantitative PCR product from 1 µg amplified DNA divided by the yield from 1 µg genomic DNA control. The results for eight loci are indicated as follows; CXCR5, open diamonds; connexin40, open triangles; MKP1, open squares; CCR6, open circles; acidic ribosomal protein, filled diamonds; CCR1, filled triangles; cJUN, filled squares; CCR7, filled circles. Figure 6A depicts the percent representation for eight loci derived from MDA-amplified DNA. Figure 6B

depicts the percent representation for eight loci present in DNA amplified using DOP-PCR. Figure 6C depicts the percent representation for eight loci present in PEP-amplified DNA.

As can be seen, for 100-, 1,000-, and 10,000-fold amplified MDA, the maximum amplification biases were only 2.7, 2.3, and 2.8 respectively, expressed as the ratio of the most highly represented gene to the least represented gene. Significantly, the 3-fold amplification bias of MDA remained almost constant between 100- and 100,000-fold amplification (Figure 6A). In contrast, amplification by the DOP-PCR method exhibited an amplification bias ranging between 4 and 6 orders of magnitude (Figure 6B). In addition, the PEP method exhibited an amplification bias spanning 2-4 orders of magnitude (Figure 6C).

G. Example 7: Amplification using nested primers yields specific amplification of c-jun sequences.

This example demonstrates the amplification of a specific DNA region from a complex mixture of DNA sequences using ϕ 29 DNA polymerase with sequence-specific, nested primers.

Amplification reactions were carried out either with or without a heat denaturing/annealing step using either exonuclease-resistant hexamers or a nested set of 19 exonuclease-resistant sequence-specific primers. The nested primers used in this example are listed in Table 2. The presence of an asterisk in the nucleotide sequence indicates the presence of a phosphorothioate bond. The nested primers are designed to hybridize to opposite strands on each side of the human c-jun gene, and the closest left and right primers encompass a 3420 bp fragment containing the c-jun gene. On each side of the c-jun gene, these nested primers are spaced 150-400 nucleotides between each other. The region of human DNA encompassing the recognition sites for these primers can be accessed from Genbank using Accession Number AL136985 and spans positions 65001 to 73010 of the nucleotide sequence.

Table 2

Left Primers (5' to 3')

c-Jun L9	TCC ATC ACG AGT TAT GC*A* C	(SEQ ID NO:1)
c-Jun L8	TGG AGT TAC TAA GGG AA*G* C	(SEQ ID NO:2)

c-Jun L7	ACT GAG TTC ATG AAC CC*T* C	(SEQ ID NO:3)
c-Jun L6	ATT AAC TCA TTG AAG GC*C* C	(SEQ ID NO:4)
c-Jun L5	TCT GTG CTG TAC TGT TG*T* C	(SEQ ID NO:5)
c-Jun L4	AGT TTG GCA AAC TGG GC*T* C	(SEQ ID NO:6)
c-Jun L3	TGG CTC TTG GTA TGA AA*A* G	(SEQ ID NO:7)
c-Jun L2	ACT GTT AGT TTC CAT AG*G* C	(SEQ ID NO:8)
c-Jun L1	TGA ATA CAT TTA TTG TG*A* C	(SEQ ID NO:9)

Right Primers (5' to 3')

c-Jun R1	CGA CTG TAG GAG GGC AG*C* G	(SEQ ID NO:10)
c-Jun R2	CGT CAG CCC ACA ATG CA*C* C	(SEQ ID NO:11)
c-Jun R3	GTA CTT GGA TTC TCA GC*C* T	(SEQ ID NO:12)
c-Jun R4	CAA ATC TCT CGG CTT CT*A* C	(SEQ ID NO:13)
c-Jun R5	CGT GTT GTG TTA AGC GT*G* T	(SEQ ID NO:14)
c-Jun R6	CCG CGG AAA AGG AAC CA*C* T	(SEQ ID NO:15)
c-Jun R7	CTC CTG GCA GCC CAG TG*A* G	(SEQ ID NO:16)
c-Jun R8	CTC CTC CCC TCG ATG CT*T* C	(SEQ ID NO:17)
c-Jun R9	CAG TTA CCC TCT GCA GA*T* C	(SEQ ID NO:18)
c-Jun R10	CTA TTT CCT CTG CAG AT*A* A	(SEQ ID NO:19)

Four reactions were carried out under the following conditions. Human

5 genomic DNA (50 ng) was placed into a 0.2 ml tube in a total volume of 50 μ l, yielding final concentrations of 25 mM Tris-HCl, pH 7.5, 50 mM KCl, 10 mM MgCl₂, and 100 μ M exonuclease-resistant hexamer or 1 μ M each of exonuclease-resistant nested primers. A heat-treatment step to increase primer annealing was included or omitted, as indicated, for individual reactions. Annealing reactions were heated to 95°C for 3

10 minutes and slowly cooled down to 37°C in a PCR System Thermocycler (Perkin Elmer). Reactions were divided into two, each had 25 μ l and then was brought to a final volume of 50 μ l, containing final concentrations of 37 mM Tris-HCl, pH 8.0, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1.0 mM dNTPs, 1 unit/ml of yeast pyrophosphatase, 50 μ M exonuclease-resistant hexamer or 0.5 μ M exonuclease-

resistant nested primers, and 800 units/ml ϕ 29 DNA polymerase. Radioactively labeled α -[32 P] dCTP, approximately 60 cpm/pmol total dNTPs, was added to one of two parallel reactions for quantification of DNA synthesis. Reactions were incubated for 18 hours at 37°C. Incorporation of acid-precipitable radioactive deoxyribonucleotide product was determined with glass fiber filters. The reactions that did not contain α -[32 P] dCTP were analyzed by TaqMan® quantitative PCR analysis as described in Example 5. A standard curve for input template was generated to determine the locus copy number of the amplified DNA sample relative to that of genomic DNA. The standard curve was generated from 0, 0.001, 0.01, 0.1, 0.5, and 1 μ g of genomic DNA. The results are shown in Figure 7.

The Y-axis is the locus representation relative to input genomic DNA. It is calculated as the yield of quantitative PCR product from 1 μ g amplified DNA divided by the yield from 1 μ g genomic DNA control, expressed as a percent.

As can be seen, the representation of c-jun sequences amplified with random hexamers from DNA heated to 95°C was 69% (see RH (heat) bar). The representation of c-jun sequences amplified with nested primers from DNA heated to 95°C was only 3% (see NP (heat) bar). The representation of c-jun sequences in DNA amplified with random hexamers without template DNA heat treatment was 211% (see RH (no heat) bar). The representation of c-jun sequences in DNA amplified with nested primers without template DNA heat treatment was 2828% (see NP (no heat) bar).

H. Example 8: Amplification Bias of Loci Amplified by MDA from Whole Blood

This example demonstrates that genomic DNA can be amplified using MDA directly from whole blood or from tissue culture cells and that the locus representation is substantially the same as for DNA amplified from purified genomic DNA template. This example illustrates lysis by subjecting cells to alkaline conditions (by addition of a lysis solution) without any lysis by heating, stabilization of the cell lysate (by addition of a stabilization solution), and multiple displacement amplification of genomic DNA in the stabilized cell lysate. The stabilized cell lysate is used for amplification without purification of the genomic DNA. As a control, DNA amplified in the absence of added template was tested and contains no detectable sequence representation for these loci.

DNA was prepared from blood or a tissue culture cell line as follows. Human blood samples were obtained from Grove Hill Laboratory. U266, a myeloma cell line, was obtained from ATCC and passaged according to the accompanying protocol. Cells were lysed in an alkaline lysis solution by a modification of Zhang et al. (Zhang, L. *et al.* Whole genome amplification from a single cell: implications for genetic analysis. *Proc Natl Acad Sci U S A.* 89, 5847-5851 (1992)). Blood was diluted 3-fold in PBS (137 mM NaCl, 2.7 mM KCl, 9.5 mM Na, KPO₄, pH 7.4), while tissue culture cells were diluted to 30,000 cells/ml in PBS. Blood or cells were lysed by dilution with an equal volume of Alkaline Lysis Buffer (400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA) and incubated 10 min on ice. The lysed cells were neutralized with the same volume of Neutralization Buffer (400 mM HCl, 600 mM Tris-HCl, pH 7.5). Preparations of lysed blood or cells (1 µl) were used directly as template in MDA reactions (100 µl) as described.

MDA reactions (100 µl) were carried out without denaturation at 95°C as described in Example 5. Reactions using purified human genomic DNA template contained 300 or 30 ng DNA, resulting in approximately 100- or 1000-fold DNA amplification. Reactions using DNA from lysed blood or cells contained 1 µl of the stabilized cell lysates as template. Control amplification reactions contained no added template DNA.

TaqMan® quantitative PCR analysis was performed on amplified DNA samples as described in Example 5, and the results are shown in Figure 8. The relative representation of eight loci for DNA from five different amplification reactions is depicted in Figure 8. The Y-axis is the locus representation, expressed as a percent, relative to input genomic DNA, which is calculated as the yield of quantitative PCR product from 1 µg of amplified DNA divided by the yield from 1 µg of genomic DNA control. Bars with declining diagonals depict the locus representation for DNA amplified from whole blood. Solid gray bars depict the locus representation for DNA amplified from 30 ng purified genomic DNA (9,000 genome copies). Solid white bars depict the locus representation for DNA amplified from 300 ng purified genomic DNA (90,000 genome copies). Bars with rising diagonals depict the locus representation for DNA amplified from tissue culture cells (10 cell equivalents of DNA). Solid black bars depict the locus representation for DNA amplified from reactions with no added

template (the values for the data represented by the black bars are so small that the bars are not visible on the graph).

As can be seen, DNA amplified directly from whole blood or from tissue culture cells without purification has substantially the same values for locus representation as DNA amplified from purified genomic DNA template.

I. Example 9: The Amplification Bias of Loci Amplified by MDA in Reactions Containing AAdUTP is the Same as the Amplification Bias of DNA Amplified in Reactions Containing 100% dTTP.

This example demonstrates that genomic DNA can be amplified using MDA in reactions containing AAdUTP (5-(3-aminoallyl)-2'-deoxyuridine 5'-triphosphate, Sigma-Aldrich Co.) and that the locus representation is substantially the same as for DNA amplified in reactions containing only dTTP.

MDA reactions (100 μ l) containing 100% dTTP were carried out without denaturation at 95°C as described in Example 5. Reactions containing 70% AAdUTP were carried out under the same conditions as reactions containing 100% dTTP, with the exception that they contained 0.7 mM AAdUTP and 0.3 mM dTTP, instead of the standard 1.0 mM dTTP. Reactions contained 1 ng human genomic DNA template, resulting in approximately 30,000-fold DNA amplification.

TaqMan® quantitative PCR analysis was performed on amplified DNA samples as described in Example 5, and the relative representation of eight loci for DNA from two amplification reactions is depicted in Figure 9. The Y-axis is the locus representation, expressed as a percent, relative to input genomic DNA, which is calculated as the yield of quantitative PCR product from 1 μ g of amplified DNA divided by the yield from 1 μ g of genomic DNA control. Black bars depict the locus representation for DNA amplified in a reaction containing 100% dTTP. White bars depict the locus representation for DNA amplified in a reaction containing 30% dTTP / 70% AAdUTP.

As can be seen, DNA amplified in reactions containing 30% dTTP / 70% AAdUTP has substantially the same values for locus representation as DNA amplified in reactions containing 100% dTTP.

J. Example 10: Amplification of c-jun sequences from human genomic DNA using sequence specific primers and target circularization.

This example describes an embodiment of the disclosed method and analysis of the resulting DNA products. The exemplified method is the disclosed gene-specific multiple displacement amplification form of multiple strand displacement amplification using nuclease-resistant sequence-specific primers and circularized DNA template.

Amplification reactions were carried out without a heat denaturing/annealing step under the conditions described in Example 5 using exonuclease-resistant sequence-specific primers that hybridize to sequences within a 5.5kb EcoRI fragment that contains the c-jun sequence. The sequence-specific primers used in this example are listed in Table 3. An asterisk in the nucleotide sequence indicates the presence of a phosphorothioate bond. The sequence-specific primers are designed to hybridize to opposite strands on each side of the c-jun gene sequence, and the primers encompass a 2025 bp fragment containing the c-jun gene sequence. The primers are spaced 150-400 nucleotides between each other on each side of the c-jun gene sequence. The region of human DNA encompassing the recognition sites for these primers can be accessed from Genbank using Accession Number AL136985 and spans positions 66962 to 68987 of the nucleotide sequence.

Table 3

Left Primers (5' to 3')

c-Jun D1	CTG AAA CAT CGC ACT AT*C *C	(SEQ ID NO:21)
c-Jun D2	CCA AAC TTT GAA ATG TT*T *G	(SEQ ID NO:22)
c-Jun D3	CTG CCA CCA ATT CCT GC*T *T	(SEQ ID NO:23)
c-Jun D4	CAT AAG CAA AGG CCA TC*T *T	(SEQ ID NO:24)
c-Jun D5	GGA AGC AAT TCA AGA TC*T *G	(SEQ ID NO:25)
c-Jun D6	CTT CAG ATT GCA GCA AT*G *T	(SEQ ID NO:26)
c-Jun D7	GAA TTA ATG AAA TTG GG*A *G	(SEQ ID NO:27)
c-Jun D8	ACT GTT AGT TTC CAT AG*G *C	(SEQ ID NO:28)
c-Jun D9	CAA GGT TGA TTA TTT TA*G *A	(SEQ ID NO:29)
c-Jun D10	AGT ACT AGT TCA TGT TT*T *C	(SEQ ID NO:30)

Right Primers (5' to 3')

c-Jun U1	TAG TAC TCC TTA AGA AC*A *C	(SEQ ID NO:31)
c-Jun U2	CTA ACA TTC GAT CTC AT*T *C	(SEQ ID NO:32)
c-Jun U3	GCG GAC GGG CTG TCC CC*G *C	(SEQ ID NO:33)
c-Jun U4	GGA AGG ACT TGG CGC GC*C *C	(SEQ ID NO:34)
c-Jun U5	AAC TAA AGC CAA GGG TA*T *C	(SEQ ID NO:35)
c-Jun U6	ATA ACA CAG AGA GAC A*G *A	(SEQ ID NO:36)
c-Jun U7	CAA CTC ATG CTA ACG CA*G *C	(SEQ ID NO:37)
c-Jun U8	GGA AGC TGG AGA GAA TC*G *C	(SEQ ID NO:38)
c-Jun U9	GAC ATG GAG TCC CAG GA*G *C	(SEQ ID NO:39)
c-Jun U10	AGG CCC TGA AGG AGG AG*C *C	(SEQ ID NO:40)

Four reactions were carried out under the following conditions. Human genomic DNA (5 µg, Coriell Cell Repositories) was digested with 100 units of EcoRI for 3 hours in 50 µl according to the manufacturers conditions. The EcoRI endonuclease was inactivated by incubation at 65°C for 30 min. Digested DNA fragments (0.5 µg) were circularized in an 840 µl volume using 1.7 units (Weiss units) of T4 DNA ligase. The reaction was incubated for 16 h at 4°C. A mock ligation was carried out under identical conditions except for the omission of DNA ligase. Two amplification reactions utilizing a portion of the ligated mixture as template (16.8 µl; 10 ng DNA) were carried out, one with sequence-specific primers at a concentration of 2.5 µM each and the other with the primers at a concentration of 0.5 µM each. Two more amplification reactions were carried out utilizing the mock-ligated DNA as template with sequence-specific primer concentrations of 2.5 µM and 0.5 µM.

Radioactively labeled α -[³²P] dCTP, approximately 60 cpm/pmol total dNTPs, was added to parallel reactions for quantification of DNA synthesis. Reactions were incubated for 18 hours at 37°C. The reactions that did not contain α -[³²P] dCTP were analyzed by TaqMan® quantitative PCR analysis as described in Example 5. A standard curve for input template was generated to determine the locus copy number of the amplified DNA sample relative to that of genomic DNA. The standard curve was generated from 0, 0.001, 0.01, 0.1, 0.5, and 1 µg of genomic DNA. The results are

shown in Figure 10. The Y-axis is the locus representation relative to input genomic DNA. It is calculated as the yield of quantitative PCR product from 1 μ g amplified DNA divided by the yield from 1 μ g genomic DNA control, expressed as a percent.

As can be seen, the representation of c-jun sequences amplified with 2.5 μ M Gene-Specific primers from ligated DNA was below detection (see GS 2.5 +L bar). The representation of c-jun sequences amplified with 2.5 μ M Gene-Specific primers from mock-ligated DNA was also not detected (see GS 2.5 -L bar). The representation of c-jun sequences in DNA amplified with 0.5 μ M Gene-Specific primers from ligated DNA was 32000% (see GS 0.5 +L bar). The representation of c-jun sequences in DNA amplified with 0.5 μ M Gene-Specific primers from mock-ligated DNA was also not detected (see GS 0.5 -L bar). These results demonstrate a 320-fold amplification of the c-jun sequences using gene-specific primers and circularized template DNA. Digested DNA that was not ligated did not show any appreciable amplification. Only ligated DNA that was amplified with 0.5 μ M primers was amplified, while 2.5 μ M primers did not yield any amplification.

The specificity of the DNA amplification reaction carried out with 2.5 μ M Gene-Specific primers and circularized DNA template was tested by comparing it to the amount of DNA amplification observed at seven other loci. TaqMan® quantitative PCR analysis was performed as described in Example 5, and the results are shown in Figure 11. The relative representation of the c-jun locus and seven other loci is depicted and the seven loci showed only low levels of amplification, indicating that the c-jun sequences were specifically amplified using the c-jun specific primers.

K. Example 11: Increasing the quality of Multiple Displacement Amplification of degraded genomic DNA by repairing the damaged sample.

This example demonstrates that the quality of amplified DNA in WGA is increased when the damaged genomic DNA is pretreated with a repair method by denatured with 70°C heat in the presence of 15 mM NaOH, adding Tris-HCl pH 4.0 and original sample back to the denatured sample, then slow cooling to hybridize the damaged DNA. The damaged gDNA was created by sonicating intact genomic DNA to an average length of 1 kb. This degraded sample was then pretreated with the repair method and placed into a multiple displacement amplification reaction. The quality of

the amplification products were assessed by a Taqman assay which compares the representation of a particular locus on a chromosome of 1 μ g of amplification product to a 1 μ g standard genomic DNA that has not been amplified. This example demonstrates that damaged genomic DNA with the repair treatment has 3-fold
5 improvement in locus representation of the amplified products compared to amplified products of damaged genomic DNA without the repair method.

The damaged genomic DNA was created as follows: Promega's genomic DNA sample was diluted to 100 ngs/ μ l in dH₂O. The sample was sonicated with a 550 Sonic Dismembrator by Fisher Scientific setting 40% for 9 seconds. The sample was diluted
10 further to 5 ng/ μ l and 25 ngs total was used to denature with 15 mM NaOH final and heated to 70°C for 5 minutes. At 70°C, Tris-HCl pH 4.0 was added to create a final pH of 8.0 and 25 ng of native original denatured samples was also added. Amplification by multiple displacement synthesis was performed on repaired damaged DNA and damaged DNA, and a Control intact DNA. The results in Figure 12 show that the
15 repair of the damaged DNA has significantly improved the amplification of the damaged sample compared to amplification of damaged DNA without repair.

L. Example 12: Increasing the quality of Multiple Displacement Amplification of real genomic DNA samples damaged by prolonged storage by repairing the damaged sample.

20 This example demonstrates that the quality of amplified product by multiple displacement amplification of genomic DNA samples damaged by prolonged storage was improved by the repair method.

The damaged genomic DNA samples were pretreated with a repair method by denaturing with 70°C heat in the presence of 15 mM NaOH, adding Tris-HCl pH 4.0
25 and original sample back to the denatured sample, slow cooling to hybridize the damaged DNA, then placing into a multiple displacement amplification reaction. The quality of the amplification products were assessed by a Taqman assay as described in Example 11.

As seen in Figure 13, the quality of the amplification products of the damaged
30 genomic DNA samples once repaired is significantly improved. On average, a 3-fold improvement in the quality of the amplification products was observed.

M. Example 13: Optimization of the denaturation step: ionic strength, heat and NaOH is results in improved repair.

This example demonstrates that ionic strength, heat and NaOH are useful for the denaturation step in the disclosed method of repairing and amplifying damaged DNA.

5 Damaged DNA samples were prepared as follows: DNA samples was diluted in dH₂O to 5 ng/μl and were heated to 70°C for 5 minutes which degrades the samples into smaller fragments. The heated DNA samples were slow cooled and multiple displacement amplification was performed on these samples. A Taqman analysis of the amplification products from the heated DNA substrate showed that use of an ionic
10 solution led to further degradation of the DNA, ultimately leading to poor amplification of these samples.

To determine if heating the damaged DNA in the presence of salt can improve the amplification of this substrate, damaged DNA (25 ngs) was heated to 70°C in the presence of a salt (Tris-EDTA (10 mM Tris pH 8.0 and 1 mM EDTA)) for 3 minutes,
15 25 ngs of native damaged sample was added back, and then the sample was cooled. This method showed some improvement on the repair of the damaged DNA. When 25 mM final of NaOH was added to the reaction before the heat step, this improved the repair method giving consistent repair of the damaged DNA. This was done by diluting damaged DNA in Tris-EDTA (TE) to 5 ng/μl, 25 ngs of DNA was added to NaOH
20 final concentration 25 mM, the sample was heated to 70°C for 3 minutes, Tris-HCl pH 4.0 was added to reduce the pH to 8.0, 25 ngs of the original native damaged DNA sample was added back, and the mixture was cooled to room temperature. Repair with heat, Tris, and NaOH gave better locus representation than the no repair and repair with heat and Tris only treatments.

25 N. Example 14: Neutralization step of the repair method improves amplification of damaged DNA sample.

This example demonstrates that the neutralization of the repair method is important for improved quality of amplification products. A damaged DNA sample was prepared as follows: a DNA sample was diluted in dH₂O to 10 ng/μl and was
30 heated to 70°C for 5 minutes to create damaged DNA samples. The heated DNA sample was diluted to 5 ngs/μl in TE, 25 ngs of damaged DNA was added to final 25 mM NaOH, heated to 70°C for 3 minutes and was either neutralized with Tris-HCl pH

4.0 to a pH of 8.0 and slow cooled or was slow cooled without neutralization. Amplification reactions of these samples were performed and the quality of products were analyzed by Taqman analysis. Repair with neutralization gave a better locus representation than the no repair and repair without neutralization treatments.

5

It is understood that the disclosed invention is not limited to the particular methodology, protocols, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

10

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells, reference to "the antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

15

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of skill in the art to which the disclosed invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are as described. Publications cited herein and the material for which they are cited are specifically incorporated by reference. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

20

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

25

CLAIMS

We claim:

1. A method of amplifying a whole genome, the method comprising, exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate comprises a whole genome, reducing the pH of the cell lysate to form a stabilized cell lysate, and incubating the stabilized cell lysate under conditions that promote replication of the genome, wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.
2. The method of claim 1 wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution.
3. The method of claim 2 wherein the lysis solution comprises a base.
4. The method of claim 3 wherein the base is an aqueous base.
5. The method of claim 3 wherein the base is potassium hydroxide, sodium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine, methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, or DBU (1,8-diazobicyclo[5,4,0]undec-7-ene).
6. The method of claim 5 wherein the base is potassium hydroxide.
7. The method of claim 6 wherein the lysis solution comprises 400 mM KOH.
8. The method of claim 7 wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.
9. The method of claim 8 wherein the lysis solution consists of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.
10. The method of claim 3 wherein the lysis solution comprises a plurality of basic agents.

11. The method of claim 2 wherein the cells are mixed with an equal volume of the lysis solution.
12. The method of claim 11 wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.
13. The method of claim 2 wherein the lysis solution comprises a buffer.
14. The method of claim 13 wherein the buffer is a phosphate buffer, Good buffer, BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, TRICINE, sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, or Bis-tris propane.
15. The method of claim 13 wherein the lysis solution comprises a plurality of buffering agents.
16. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 7.0 to about pH 6.8.
17. The method of claim 1 wherein the pH of the cell lysate is reduced by mixing the cell lysate with a stabilization solution.
18. The method of claim 17 wherein the stabilization solution comprises a buffer.
19. The method of claim 18 wherein the buffer is phosphate buffer, Good buffer, BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES, TRICINE, sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, or Bis-tris propane.
20. The method of claim 19 wherein the buffer is Tris-HCl at pH 4.1.
21. The method of claim 20 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.
22. The method of claim 21 wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4.1.
23. The method of claim 18 wherein the stabilization solution comprises a plurality of buffering agents.
24. The method of claim 17 wherein the cell lysate is mixed with an equal volume of the stabilization solution.
25. The method claim 24 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

26. The method of claim 17 wherein the stabilization solution comprises an acid.

27. The method of claim 26 wherein the acid is hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, or a carboxylic acid.

28. The method of claim 27 wherein the carboxylic acid is ethanoic, propanoic, or butanoic.

29. The method of claim 17 wherein the stabilization solution comprises a plurality of acidic agents.

30. The method of claim 1 wherein the pH of the cell lysate is reduced to about pH 9.0 or below.

31. The method of claim 1 wherein the pH of the cell lysate is reduced to about pH 8.5 or below.

32. The method of claim 1 wherein the pH of the cell lysate is reduced to about pH 8.0 or below.

33. The method of claim 1 wherein the pH of the cell lysate is reduced to about pH 7.5 or below.

34. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 9.0 to about pH 6.0.

35. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 9.0 to about pH 7.0.

36. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 9.0 to about pH 7.5.

37. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 9.0 to about pH 8.0.

38. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.5 to about pH 6.0.

39. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.5 to about pH 7.0.

40. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.5 to about pH 7.5.

41. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.5 to about pH 8.0.
42. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.0 to about pH 6.0.
43. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.0 to about pH 6.5.
44. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.0 to about pH 7.0.
45. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 8.0 to about pH 7.5.
46. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 7.5 to about pH 6.0.
47. The method of claim 1 wherein the pH of the cell lysate is reduced to the range of about pH 7.5 to about pH 7.0.
48. The method of claim 1 wherein nucleic acids in the cell lysate and the stabilized cell lysate are not separated from other material in the cell lysate.
49. The method of claim 1 wherein the cell lysate and the stabilized cell lysate are not subjected to purification prior to the incubation.
50. The method of claim 49 wherein the purification comprises separation of nucleic acids in the cell lysate from other material in the cell lysate.
51. The method of claim 49 wherein the purification comprises centrifugation, extraction, chromatography, filtration, dialysis, or a combination of these.
52. The method of claim 49 wherein the purification comprises precipitation other than precipitation caused by the alkaline conditions or by the reduction of the pH.
53. The method of claim 49 wherein the purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these.
54. The method of claim 1 wherein the cell lysate, stabilized cell lysate, or both are subjected to partial purification prior to the incubation.
55. The method of claim 54 wherein the partial purification comprises centrifugation, extraction, chromatography, precipitation, filtration, dialysis, or a combination of these.

56. The method of claim 54 wherein the partial purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these.

57. The method of claim 1 wherein the cell lysate and the stabilized cell lysate are not subjected to substantial purification prior to the incubation.

58. The method of claim 57 wherein the substantial purification does not include centrifugation, extraction, chromatography, precipitation, filtration, or dialysis.

59. The method of claim 57 wherein the substantial purification does not include centrifugation, phenol-chloroform extraction, or column chromatography.

60. The method of claim 57 wherein the cell lysate, stabilized cell lysate, or both are subjected to centrifugation, extraction, chromatography, precipitation, filtration, or dialysis prior to the incubation.

61. The method of claim 57 wherein the cell lysate, stabilized cell lysate, or both are subjected to centrifugation, phenol-chloroform extraction, or column chromatography prior to the incubation.

62. The method of claim 57 wherein the substantial purification comprises centrifugation, extraction, chromatography, filtration, dialysis, or a combination of these.

63. The method of claim 57 wherein the substantial purification comprises precipitation other than precipitation caused by the alkaline conditions or by the reduction of the pH.

64. The method of claim 57 wherein the substantial purification comprises centrifugation, phenol-chloroform extraction, column chromatography, or a combination of these.

65. The method of claim 1 wherein the cell lysate and the stabilized cell lysate are not purified prior to the incubation.

66. The method of claim 1 wherein the cell lysate, stabilized cell lysate, or both are partially purified prior to the incubation.

67. The method of claim 1 wherein the incubation is substantially isothermic.

68. The method of claim 67 wherein neither the cell lysate nor the stabilized cell lysate is heated substantially above the temperature of the incubation.

69. The method of claim 67 wherein neither the cell lysate nor the stabilized cell lysate is subjected to substantial heating above the temperature of the incubation.

70. The method of claim 67 wherein the cells are not heated substantially above the temperature of the incubation.

71. The method of claim 67 wherein the cells are not subjected to substantial heating above the temperature of the incubation.

72. The method of claim 67 wherein the cells are not heated substantially above the temperature at which the cells grow.

73. The method of claim 67 wherein the cells are not subjected to substantial heating above the temperature at which the cells grow.

74. The method of claim 67 wherein the cell lysate, stabilized cell lysate, and the cells are not heated substantially above the temperature of the incubation.

75. The method of claim 67 wherein the cell lysate, stabilized cell lysate, and the cells are not subjected to substantial heating above the temperature of the incubation.

76. The method of claim 67 wherein the cell lysate, stabilized cell lysate, and the cells are not heated, prior to or during the incubation, substantially above the temperature at which the cells grow.

77. The method of claim 67 wherein the cell lysate, stabilized cell lysate, and the cells are not subjected to, prior to or during the incubation, substantial heating above the temperature at which the cells grow prior.

78. The method of claim 1 wherein neither the cell lysate nor the stabilized cell lysate is heated above a temperature and for a time that would cause notable denaturation of the genome.

79. The method of claim 1 wherein neither the cell lysate nor the stabilized cell lysate is subjected to heating above a temperature and for a time that would cause notable denaturation of the genome.

80. The method of claim 1 wherein the cells are not lysed by heat.

81. The method of claim 1 wherein the cells are not heated above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions.

82. The method of claim 1 wherein the cells are not subjected to heating above a temperature and for a time that would cause substantial cell lysis in the absence of the alkaline conditions.

83. A method of amplifying a whole genome, the method comprising,
exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution,

reducing the pH of the cell lysate to form a stabilized cell lysate, wherein the pH of the cell lysate is reduced by mixing the cell lysate with a stabilization solution, and
incubating the stabilized cell lysate under conditions that promote replication of the genome,

wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

84. The method of claim 83 wherein the lysis solution comprises potassium hydroxide.

85. The method of claim 84 wherein the lysis solution comprises 400 mM KOH.

86. The method of claim 85 wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

87. The method of claim 86 wherein the lysis solution consists of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

88. The method of claim 83 wherein the cells are mixed with an equal volume of the lysis solution.

89. The method of claim 88 wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

90. The method of claim 83 wherein the stabilization solution comprises Tris-HCl at pH 4.1.

91. The method of claim 90 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

92. The method of claim 91 wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4.1.

93. The method of claim 83 wherein the cell lysate is mixed with an equal volume of the stabilization solution.

94. The method claim 93 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

95. The method of claim 83 wherein the lysis solution consists of 400 mM KOH and 10 mM EDTA, wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4, wherein the stabilized cell lysate is incubated in the presence of 37.5 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1 mM deoxynucleotide triphosphates, 50 μM primers, and φ29 DNA Polymerase.

96. The method of claim 95 wherein the stabilized cell lysate is incubated in the presence of 37.5 mM Tris-HCl, 50 mM KCl, 10 mM MgCl₂, 5 mM (NH₄)₂SO₄, 1 mM deoxynucleotide triphosphates, 50 μM primers, and φ29 DNA Polymerase by mixing the stabilized cell lysate with one quarter volume of reaction mix, and φ29 DNA Polymerase, wherein the reaction mix consists of 150 mM Tris-HCl, 200 mM KCl, 40 mM MgCl₂, 20 mM (NH₄)₂SO₄, 4 mM deoxynucleotide triphosphates, and 0.2 mM primers.

97. A method of amplifying a whole genome, the method comprising,
exposing cells to alkaline conditions to form a cell lysate, wherein the cell lysate comprises a whole genome, wherein the cells are exposed to alkaline conditions by mixing the cells with a lysis solution, wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA,

reducing the pH of the cell lysate to form a stabilized cell lysate, wherein the pH of the cell lysate is reduced by mixing the cell lysate with a stabilization solution, wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1, and

incubating the stabilized cell lysate under conditions that promote replication of the genome,

wherein replication of the genome results in replicated strands, wherein during replication at least one of the replicated strands is displaced from the genome by strand displacement replication of another replicated strand.

98. The method of claim 97 wherein the cells are mixed with an equal volume of the lysis solution.

99. The method of claim 97 wherein the cell lysate is mixed with an equal volume of the stabilization solution.

100. A kit for amplifying a whole genome, the kit comprising
a lysis solution,
a stabilization solution,
a set of primers, and
a DNA polymerase.

101. The kit of claim 100 wherein the lysis solution comprises potassium hydroxide.

102. The kit of claim 101 wherein the lysis solution comprises 400 mM KOH.

103. The kit of claim 102 wherein the lysis solution comprises 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

104. The kit of claim 103 wherein the lysis solution consists of 400 mM KOH, 100 mM dithiothreitol, and 10 mM EDTA.

105. The kit of claim 100 wherein the stabilization solution comprises Tris-HCl at pH 4.1.

106. The kit of claim 105 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

107. The kit of claim 106 wherein the stabilization solution consists of 800 mM Tris-HCl, pH 4.1.

108. The kit of claim 100 further comprising deoxynucleotide triphosphates.

109. The kit of claim 100 further comprising

1M dithiothreitol,

1X Phosphate-Buffered Saline, pH 7.5, and

control DNA template,

wherein the lysis solution comprises 400 mM KOH and 10 mM EDTA,

wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4,

wherein the set of primers comprises a reaction mix, wherein the reaction mix comprises 150 mM Tris-HCl, 200 mM KCl, 40 mM MgCl₂, 20 mM (NH₄)₂SO₄, 4 mM deoxynucleotide triphosphates, and 0.2 mM random hexamer primers,

wherein the DNA polymerase is ϕ 29 DNA polymerase.

110. The kit of claim 100 further comprising one or more detection probes, wherein the detection probes each comprise a complementary portion, wherein the complementary portion is complementary to a nucleic acid sequence of interest.

111. The method of claim 110 wherein the kit is designed to detect nucleic acid sequences of interest in the genome.

112. The method of claim 111 wherein the kit is designed to assess a disease, condition or predisposition of an individual based on the nucleic acid sequences of interest.

113. A method of amplifying damaged DNA, the method comprising exposing a damaged DNA sample to conditions that promote substantial denaturation of damaged DNA in the damaged DNA sample, thereby forming a denatured damaged DNA sample,

altering the conditions to conditions that do not promote substantial denaturation of damaged DNA in the damaged DNA sample to form a stabilized damaged DNA sample,

incubating damaged DNA in the stabilized damaged DNA sample under conditions that promote replication of the damaged DNA, wherein replication of the damaged DNA results in a longer average fragment length for the replicated damaged DNA than the average fragment length in the damaged DNA sample, wherein during replication at least one of the replicated strands is displaced by strand displacement replication of another replicated strand.

114. The method of claim 113 wherein the damaged DNA sample, the denatured damaged DNA sample, or both are exposed to ionic conditions.

115. The method of claim 114 wherein the damaged DNA sample and denatured damaged DNA sample are exposed to ionic conditions by mixing an ionic solution with the damaged DNA sample.

116. The method of claim 115 wherein the ionic solution is mixed with the damaged DNA sample prior to or during exposure of the damaged DNA sample to conditions that promote substantial denaturation of the damaged DNA.

117. The method of claim 115 wherein the ionic solution is a salt solution.

118. The method of claim 117 wherein the salt solution comprises one or more salts.

119. The method of claim 118 wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination.

120. The method of claim 119 wherein the Tris-HCl is from pH 7.0 to 8.0.

121. The method of claim 119 wherein the salt is Tris-EDTA.

122. The method of claim 121 wherein the salt solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA.

123. The method of claim 122 wherein the ionic solution is diluted 2 to 5 fold when mixed with the damaged DNA sample.

124. The method of claim 114 wherein the denatured damaged DNA sample is exposed to ionic conditions by mixing an ionic solution with the denatured damaged DNA sample.

125. The method of claim 124 wherein the ionic solution is mixed with the denatured damaged DNA sample prior to or during altering of the conditions.

126. The method of claim 113 wherein the damaged DNA sample is exposed to conditions that promote substantial denaturation by mixing the damaged DNA sample with a denaturing solution and by heating the damaged DNA sample to a temperature and for a length of time that substantially denatures the damaged DNA in the damaged DNA sample.

127. The method of claim 126 wherein the damaged DNA sample is mixed with the denaturing solution after the DNA sample is heated.

128. The method of claim 126 wherein the damaged DNA sample is mixed with the denaturing solution before the DNA sample is heated.

129. The method of claim 126 wherein the damaged DNA sample is mixed with the denaturing solution at the same time the DNA sample is heated.

130. The method of claim 126 wherein the damaged DNA sample is mixed with the denaturing solution during heating of the DNA sample.

131. The method of claim 126 wherein the damaged DNA sample is mixed with the denaturing solution when heating of the DNA sample begins.

132. The method of claim 126 wherein mixing the damaged DNA sample with a denaturing solution produces alkaline conditions in the damaged DNA sample.

133. The method of claim 132 wherein the denaturing solution comprises a base.

134. The method of claim 133 wherein the base is an aqueous base.

135. The method of claim 133 wherein the base is sodium hydroxide, potassium hydroxide, potassium acetate, sodium acetate, ammonium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium carbonate, ammonia, aniline, benzylamine, n-butylamine, diethylamine, dimethylamine, diphenylamine, ethylamine, ethylenediamine, methylamine, N-methylaniline, morpholine, pyridine, triethylamine, trimethylamine, aluminum hydroxide, rubidium hydroxide, cesium hydroxide, strontium hydroxide, barium hydroxide, or DBU (1,8-diazobicyclo[5,4,0]undec-7-ene).

136. The method of claim 135 wherein the base is sodium hydroxide.

137. The method of claim 136 wherein the denaturing solution comprises about 150 mM to about 1 M NaOH.

138. The method of claim 132 wherein the denaturing solution is 10X concentration, wherein the damaged DNA sample is mixed with the denaturing solution to create a 1X concentration.

139. The method of claim 132 wherein the alkaline conditions comprise 15 to 50 mM NaOH.

140. The method of claim 126 wherein the damaged DNA in the damaged DNA sample is substantially denatured without further damaging the DNA.

141. The method of claim 126 wherein the damaged DNA sample is heated to a temperature of about 70°C or less and for a length of time of about 5 minutes or less.

142. The method of claim 126 wherein the temperature is about 60°C to about 70°C.

143. The method of claim 126 wherein the temperature is about 50°C to about 60°C.

144. The method of claim 126 wherein the temperature is about 40°C to about 50°C.

145. The method of claim 126 wherein the temperature is about 25°C to about 40°C.

146. The method of claim 126 wherein the temperature is about 60°C to about 70°C and the length of time is about 3 minutes.

147. The method of claim 126 wherein the temperature is about 25°C to about 50°C and the length of time is about 5 minutes or more.

148. The method of 126 wherein altering the conditions comprises reducing the pH of and cooling the denatured damaged DNA sample.

149. The method of claim 148 wherein the temperature to which the damaged DNA sample is heated is maintained during reduction of the pH of the denatured damaged DNA sample.

150. The method of claim 148 wherein the temperature to which the damaged DNA sample is heated is reduced before reduction of the pH of the denatured damaged DNA sample.

151. The method of claim 148 wherein the temperature to which the damaged DNA sample is heated is reduced during reduction of the pH of the denatured damaged DNA sample.

152. The method of claim 148 wherein cooling the denatured damaged DNA sample is commenced during reduction of the pH of the denatured damaged DNA sample.

153. The method of claim 148 wherein cooling the denatured damaged DNA sample is commenced when the pH of the denatured damaged DNA sample is reduced.

154. The method of claim 148 wherein the pH of the denatured damaged DNA sample is reduced by mixing the denatured damaged DNA sample with a stabilization solution.

155. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 7.5 to about pH 8.0.

156. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced by mixing the denatured damaged DNA sample with a stabilization solution.

157. The method of claim 156 wherein the stabilization solution comprises a buffer.

158. The method of claim 154 wherein the buffer is phosphate buffer, Good buffer, BES, BICINE, CAPS, EPPS, HEPES, MES, MOPS, PIPES, TAPS, TES,

TRICINE, sodium cacodylate, sodium citrate, triethylammonium acetate, triethylammonium bicarbonate, Tris, Bis-tris, or Bis-tris propane.

159. The method of claim 154 wherein the stabilization solution comprises 800 mM Tris-HCl, pH 4.1.

160. The method of claim 154 wherein the stabilization solution comprises an acid.

161. The method of claim 160 wherein the acid is hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, acetylsalicylic acid, ascorbic acid, carbonic acid, citric acid, formic acid, nitric acid, perchloric acid, HF, HBr, HI, H₂S, HCN, HSCN, HClO, monochloroacetic acid, dichloroacetic acid, trichloroacetic acid, or a carboxylic acid.

162. The method of claim 160 wherein the carboxylic acid is ethanoic, propanoic, or butanoic.

163. The method of claim 160 wherein the stabilization solution comprises a plurality of acidic agents.

164. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 9.0 to about pH 6.8.

165. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 9.0 to about pH 7.5.

166. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 9.0 to about pH 8.0.

167. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 8.5 to about pH 6.8.

168. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 8.5 to about pH 7.5.

169. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 8.5 to about pH 8.0.

170. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 8.0 to about pH 6.8.

171. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to the range of about pH 8.0 to about pH 7.5.

172. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to about pH 9.0 or less.

173. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to about pH 8.5 or less.

174. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to about pH 8.0 or less.

175. The method of claim 154 wherein the pH of the denatured damaged DNA sample is reduced to about pH 7.5 or less.

176. The method of claim 154 wherein the stabilization solution comprises one or more salts.

177. The method of claim 176 wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination.

178. The method of claim 177 wherein the Tris-HCl is from pH 7.0 to 8.0.

179. The method of claim 177 wherein the salt is Tris-EDTA.

180. The method of claim 179 wherein the stabilization solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA.

181. The method of claim 148 wherein the damaged DNA mixture is cooled at a rate of about 1°C per minute or less.

182. The method of claim 148 wherein the damaged DNA mixture is cooled at a rate of about 1% per minute or less.

183. The method of claim 182 wherein the damaged DNA mixture is cooled to room temperature or lower from 60°C to 70°C.

184. The method of claim 182 wherein the damaged DNA mixture is cooled to room temperature or lower from 50 to 60°C.

185. The method of claim 182 wherein the damaged DNA mixture is cooled to room temperature or lower from 40°C to 50°C.

186. The method of claim 182 wherein the damaged DNA mixture is cooled to room temperature or lower from 30°C to 40°C.

187. The method of claim 182 wherein the damaged DNA mixture is cooled to room temperature from 50°C to 70°C.

188. The method of claim 126 wherein the denaturing solution comprises one or more salts.

189. The method of claim 188 wherein the salt is Tris-HCl, Tris-EDTA, sodium chloride, potassium chloride, magnesium chloride, sodium acetate, potassium acetate, magnesium acetate, or a combination.

190. The method of claim 189 wherein the Tris-HCl is from pH 7.0 to 8.0.

191. The method of claim 189 wherein the salt is Tris-EDTA.

192. The method of claim 191 wherein the denaturing solution comprises about 50 mM to about 500 mM Tris and about 1 mM to about 5 mM EDTA.

193. The method of claim 113 wherein the damaged DNA sample is comprised of degraded DNA fragments of genomic DNA.

194. The method of claim 113 wherein replication and repair of the damaged DNA is accomplished by incubating the damaged DNA in the presence of a DNA polymerase.

195. The method of claim 194 wherein the polymerase is a DNA polymerase that can extend the 3'-ends of the damaged DNA.

196. The method of claim 195 wherein the DNA polymerase is ϕ 29 DNA polymerase, BST DNA polymerase, Taq DNA polymerase, a modified form of Taq DNA polymerase, a Reverse Transcriptase, T4 DNA polymerase, T7 DNA polymerase, Pol I DNA polymerase, or a modified form of DNA Polymerase I.

197. The method of claim 194 wherein the DNA polymerase is ϕ 29 DNA Polymerase.

198. The method of claim 194 wherein the damaged DNA is amplified using a kit, wherein the kit comprises

- a denaturing solution,
- a stabilization solution,
- a set of primers, and
- a DNA polymerase.

199. The method of claim 113 wherein the damaged DNA sample is a cell lysate, wherein the cell lysate is produced by exposing cells to alkaline condition, wherein the cell lysate comprises a whole genome,.

200. The method of claim 1 further comprising

following exposure of the cells to alkaline conditions, exposing a first portion of the cell lysate to conditions that promote substantial denaturation of damaged DNA in the first portion of the cell lysate, wherein reducing the pH of the cell lysate comprises reducing the pH of the first portion of the cell lysate to form a first stabilized cell lysate and reducing the pH of a second portion of the cell lysate to form a second stabilized cell lysate,

following reducing the pH of the cell lysate, mixing the second stabilized cell lysate with the first stabilized cell lysate under conditions that promote transient denaturation of the ends of damaged DNA in the second stabilized cell lysate and that maintain substantial denaturation of the damaged DNA in the first stabilized cell lysate, thereby forming a stabilized cell lysate mixture, and

prior to incubating the stabilized cell lysate, cooling the stabilized cell lysate mixture under conditions that promote annealing of the ends of the transiently denatured damaged DNA to the substantially denatured damaged DNA,

wherein incubating the stabilized cell lysate under conditions that promote replication of the genome also promotes replication of the damaged DNA, wherein the annealed ends of the damaged DNA prime replication, wherein replication of the damaged DNA results in repair of the replicated strands.

201. A method of amplifying damaged DNA, the method comprising

exposing a first damaged DNA sample to conditions that promote substantial denaturation of damaged DNA in the first damaged DNA sample, thereby forming a denatured damaged DNA sample,

reducing the pH of the denatured damaged DNA sample to form a stabilized denatured damaged DNA sample,

mixing a second damaged DNA sample with the stabilized denatured damaged DNA sample under conditions that promote transient denaturation of the ends of damaged DNA in the second sample and that maintain substantial denaturation of the damaged DNA in the stabilized denatured damaged DNA sample, thereby forming a damaged DNA mixture,

cooling the damaged DNA mixture under conditions that promote annealing of the ends of the transiently denatured damaged DNA to the substantially denatured damaged DNA,

incubating the annealed damaged DNA under conditions that promote replication of the damaged DNA, wherein the annealed ends of the damaged DNA prime replication, wherein replication of the damaged DNA results in repair of the replicated strands, wherein during replication at least one of the replicated strands is displaced by strand displacement replication of another replicated strand.

202. The method of claim 201 wherein the temperature to which the first damaged DNA sample is heated is maintained during mixing of the second damaged DNA sample with the stabilized denatured damaged DNA sample.

203. The method of claim 201 wherein the pH of the stabilized denatured damaged DNA sample is not high enough nor low enough to cause further substantial denaturation upon mixing second damaged DNA sample with the stabilized denatured damaged DNA sample.

204. The method of claim 201 wherein the first damaged DNA sample is a portion of a damaged DNA sample, wherein the second damaged DNA sample is a portion of the same damaged DNA sample.

205. The method of claim 201 wherein the first damaged DNA sample is from the same source as the second damaged DNA sample.

206. The method of claim 201 wherein the first damaged DNA sample is from the same organism as the second damaged DNA sample.

207. The method of claim 201 wherein the first damaged DNA sample is from the same tissue as the second damaged DNA sample.

208. The method of claim 201 wherein the second damaged DNA sample is mixed with the stabilized denatured damaged DNA sample at a temperature and for a length of time that transiently denatures the damaged DNA in the second damaged DNA sample.

209. The method of claim 208 wherein the temperature is about 70°C or less and the length of time is about 30 seconds or less.

210. The method of claim 208 wherein the second damaged DNA sample is mixed with the stabilized denatured damaged DNA sample at a temperature that does not further damaging the DNA.

211. The method of claim 210 wherein the temperature is about 60°C to about 70°C.

212. The method of claim 211 wherein the temperature is about 50°C to about 60°C.

213. The method of claim 212 wherein the temperature is about 40°C to about 50°C.

214. The method of claim 213 wherein the temperature is about 25°C to about 40°C.

215. The method of claim 201 wherein the temperature is about 25°C to about 70°C and the length of time is about 30 seconds.

Figure 1




Figure 2




Figure 3

Effect of time at 95 degrees on rate and yield of MDA




Figure 4

Effect of time at 95 degrees on
average product DNA strand size




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10
Amplification of c-jun sequence using Gene Specific MDA





Figure 11
Specific amplification of c-jun sequences from human
genomic DNA



Repair Method Improves Sonicated gDNA Samples In the WGA Reaction

Comparison of Multiple Displacement Amplification with or
without the Repair Method




FIG. 12

Recovery of Damaged gDNA Samples in Whole Genome Amplification Reaction

Taqman Analysis of Multiple Displacement Amplification of 40 Samples with or without the Recovery Treatment




FIG. 13

SEQUENCE LISTING

<110> Molecular Staging, Inc.

Frank B. Dean
Roger S. Lasken
Linhua Fang
A. Fawad Faruqi
Osama A. Alsmadi
Mark D. Driscoll
Seiyu Hosono
Michele Wisniewski
Wanmin Song

<120> NUCLEIC ACID AMPLIFICATION

<130> 13172.0012P1

<150> 09/997,868

<151> 2001-10-15

<160> 40

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 1

tccatcacga gttatgcac

19

<210> 2

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 2

tggagttact aaggaagc

19

<210> 3

<211> 19

<212> DNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 3
actgagttca tgaaccctc 19

<210> 4
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 4
attaactcat tgaaggccc 19

<210> 5
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 5
tctgtgctgt actgttgtc 19

<210> 6
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 6
agtttggcaa actgggctc 19

<210> 7
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 7
tggctcttgg tatgaaaag 19

<210> 8
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 8
actgttagtt tccataggc 19

<210> 9
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 9
tgaatacatt tattgtgac 19

<210> 10
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 10
cgactgtagg agggcagcg 19

<210> 11
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 11
cgtcagccca caatgcacc 19

<210> 12
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 12
gtacttggat tctcagcct 19

<210> 13
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 13
caaatctctc ggcttctac 19

<210> 14
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 14
cgtgttgtgt taagcgtgt 19

<210> 15
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 15
ccgcggaaaa ggaaccact 19

<210> 16
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 16
ctcctggcag cccagtgag 19

<210> 17
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 17
ctcctcccct cgatgcttc 19

<210> 18
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 18
cagttaccct ctgcagatc 19

<210> 19
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 19
ctatttcctc tgcagataa 19

<210> 20
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<221> misc_feature
<222> 11-16
<223> N = a, g, c, or t (u)

<400> 20
ccgactcgag nnnnnnatgt gg 22

<210> 21
<211> 19
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 21

ctgaaacatc gcactatcc

19

<210> 22

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 22

ccaaactttg aaatgtttg

19

<210> 23

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 23

ctgccaccaa ttcctgctt

19

<210> 24

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 24

cataagcaaa ggccatctt

19

<210> 25

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 25
ggaagcaatt caagatctg 19

<210> 26
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 26
cttcagattg cagcaatgt 19

<210> 27
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 27
gaattaatga aattgggag 19

<210> 28
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 28
actgttagtt tccataggc 19

<210> 29
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 29
caaggttgat tattttaga 19

<210> 30
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 30
agtactagtt catgttttc 19

<210> 31
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 31
tagtactcct taagaacac 19

<210> 32
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 32
ctaacattcg atctcattc 19

<210> 33
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 33
gcggacgggc tgtccccgc 19

<210> 34
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 34
ggaaggactt ggcgcgcc 19

<210> 35
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 35
aactaaagcc aagggtatc 19

<210> 36
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 36
ataacacaga gagacaga 18

<210> 37
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 37
caactcatgc taacgcagc 19

<210> 38
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 38
ggaagctgga gagaatcgc 19

<210> 39
<211> 19
<212> DNA
<213> Artificial Sequence

10

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 39

gacatggagt cccaggagc

19

<210> 40

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 40

aggccctgaa ggaggagcc

19